

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
1 May 2003 (01.05.2003)

PCT

(10) International Publication Number  
WO 03/035075 A1

(51) International Patent Classification<sup>7</sup>: A61K 31/505,  
31/53, C07D 473/34, 473/38, 473/24, 473/16, 473/00,  
473/40, 403/12, 487/04

(21) International Application Number: PCT/US02/27240

(22) International Filing Date: 27 August 2002 (27.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
10/027,591 19 October 2001 (19.10.2001) US

(71) Applicant: ICOS CORPORATION [US/US]; 22021  
20th Avenue S.E., Bothell, WA 98021 (US).

(72) Inventors: SADHU, Chanchal; 903 233rd Street, S.E.,  
Bothell, WA 98021 (US). DICK, Kenneth, O.; 22517 1st  
Place West, Bothell, WA 98021-8321 (US). TREIBERG,  
Jennifer; 19411 Grannis Road, Bothell, WA 98012 (US).  
SOWELL, C., Gregory; 9423 61st Avenue W., Mukilteo,  
WA 98275 (US). KESICKI, Edward, A.; 2504 208th  
Place SE, Bothell, WA 98021 (US). OLIVER, Amy;  
23123 15th Avenue, S.E., Bothell, WA 98021 (US).

(74) Agent: NAPOLI, James, J.; Marshall, Gerstein &  
Borun, Suite 6300, Sears Tower, 233 South Wacker Drive,  
Chicago, IL 60606-6357 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC,  
VN, YU, ZA, ZM, ZW.



(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: INHIBITORS OF HUMAN PHOSPHATIDYL-INOSITOL 3-KINASE DELTA



(57) Abstract: Methods of inhibiting phosphatidylinositol 3-kinase delta isoform (PI3K $\delta$ ) activity, and methods of treating diseases, such as disorders of immunity and inflammation, in which PI3K $\delta$  plays a role in leukocyte function with compounds of formula (1) are disclosed. Preferably, the methods employ active agents that selectively inhibit PI3K $\delta$ , while not significantly inhibiting activity of other PI3K isoforms. Compounds of formula (1) are provided that inhibit PI3K $\delta$  activity, including compounds that selectively inhibit PI3K $\delta$  activity. Methods of using PI3K $\delta$  inhibitory compounds to inhibit cancer cell growth or proliferation are also provided. Accordingly, the invention provides methods of using PI3K $\delta$  inhibitory compounds to inhibit PI3K $\delta$ -mediated processes *in vitro* and *in vivo*. Wherein R<sup>1</sup>-R<sup>3</sup>, X, Y and A are defined herein.

- 1 -

INHIBITORS OF HUMAN PHOSPHATIDYL-  
INOSITOL 3-KINASE DELTA5                    CROSS-REFERENCE TO RELATED APPLICATIONS

                  This application is a continuation-in-part  
of U.S. Serial No. 09/841,341, filed April 24, 2001,  
pending, which claims the benefit of U.S. provision-  
10 al application Serial No. 60/199,655, filed April  
25, 2000 and U.S. provisional application Serial No.  
60/238,057, filed October 25, 2000.

15                    FIELD OF THE INVENTION

                  The present invention relates generally to  
phosphatidylinositol 3-kinase (PI3K) enzymes, and  
more particularly to selective inhibitors of PI3K  
activity and to methods of using such materials.

20

BACKGROUND OF THE INVENTION

                  Cell signaling via 3'-phosphorylated  
phosphoinositides has been implicated in a variety  
25 of cellular processes, e.g., malignant transforma-  
tion, growth factor signaling, inflammation, and  
immunity (see Rameh et al., *J. Biol Chem*, 274:8347-  
8350 (1999) for a review). The enzyme responsible  
for generating these phosphorylated signaling prod-  
30 ucts, phosphatidylinositol 3-kinase (PI 3-kinase;  
PI3K), was originally identified as an activity  
associated with viral oncoproteins and growth factor  
receptor tyrosine kinases that phosphorylates phos-  
phatidylinositol (PI) and its phosphorylated deriva-

- 2 -

tives at the 3'-hydroxyl of the inositol ring (Panayotou et al., *Trends Cell Biol* 2:358-60 (1992)).

5           The levels of phosphatidylinositol-3,4,5-triphosphate (PIP3), the primary product of PI 3-kinase activation, increase upon treatment of cells with a variety of agonists. PI 3-kinase activation, therefore, is believed to be involved in a range of cellular responses including cell growth, differ-  
10   entiation, and apoptosis (Parker et al., *Current Biology*, 5:577-99 (1995); Yao et al., *Science*, 267:2003-05 (1995)). Though the downstream targets of phosphorylated lipids generated following PI 3-kinase activation have not been well characterized,  
15   emerging evidence suggests that pleckstrin-homology domain- and FYVE-finger domain-containing proteins are activated when binding to various phosphatidyl-inositol lipids (Sternmark et al., *J Cell Sci*, 112:4175-83 (1999); Lemmon et al., *Trends Cell Biol*,  
20   7:237-42 (1997)). *In vitro*, some isoforms of protein kinase C (PKC) are directly activated by PIP3, and the PKC-related protein kinase, PKB, has been shown to be activated by PI 3-kinase (Burgering et al., *Nature*, 376:599-602 (1995)).

25           Presently, the PI 3-kinase enzyme family has been divided into three classes based on their substrate specificities. Class I PI3Ks can phosphorylate phosphatidylinositol (PI), phosphatidyl-inositol-4-phosphate, and phosphatidylinositol-4,5-  
30   biphosphate (PIP2) to produce phosphatidylinositol-3-phosphate (PIP), phosphatidylinositol-3,4-biphosphate, and phosphatidylinositol-3,4,5-triphosphate, respectively. Class II PI3Ks phosphorylate PI and

- 3 -

phosphatidylinositol-4-phosphate, whereas Class III PI3Ks can only phosphorylate PI.

The initial purification and molecular cloning of PI 3-kinase revealed that it was a heterodimer consisting of p85 and p110 subunits (Otsu et al., *Cell*, 65:91-104 (1991); Hiles et al., *Cell*, 70:419-29 (1992)). Since then, four distinct Class I PI3Ks have been identified, designated PI3K  $\alpha$ ,  $\beta$ ,  $\delta$ , and  $\gamma$ , each consisting of a distinct 110 kDa catalytic subunit and a regulatory subunit. More specifically, three of the catalytic subunits, i.e., p110 $\alpha$ , p110 $\beta$  and p110 $\delta$ , each interact with the same regulatory subunit, p85; whereas p110 $\gamma$  interacts with a distinct regulatory subunit, p101. As described below, the patterns of expression of each of these PI3Ks in human cells and tissues are also distinct. Though a wealth of information has been accumulated in recent past on the cellular functions of PI 3-kinases in general, the roles played by the individual isoforms are largely unknown.

Cloning of bovine p110 $\alpha$  has been described. This protein was identified as related to the *Saccharomyces cerevisiae* protein: Vps34p, a protein involved in vacuolar protein processing. The recombinant p110 $\alpha$  product was also shown to associate with p85 $\alpha$ , to yield a PI3K activity in transfected COS-1 cells. See Hiles et al., *Cell*, 70, 419-29 (1992).

The cloning of a second human p110 isoform, designated p110 $\beta$ , is described in Hu et al., *Mol Cell Biol*, 13:7677-88 (1993). This isoform is said to associate with p85 in cells, and to be ubiquitously expressed, as p110 $\beta$  mRNA has been found



- 4 -

in numerous human and mouse tissues as well as in human umbilical vein endothelial cells, Jurkat human leukemic T cells, 293 human embryonic kidney cells, mouse 3T3 fibroblasts, HeLa cells, and NBT2 rat bladder carcinoma cells. Such wide expression suggests that this isoform is broadly important in signaling pathways.

Identification of the p110 $\delta$  isoform of PI 3-kinase is described in Chantry et al., *J Biol Chem*, 272:19236-41 (1997). It was observed that the human p110 $\delta$  isoform is expressed in a tissue-restricted fashion. It is expressed at high levels in lymphocytes and lymphoid tissues, suggesting that the protein might play a role in PI 3-kinase-mediated signaling in the immune system. Details concerning the P110 $\delta$  isoform also can be found in U.S. Patent Nos. 5,858,753; 5,822,910; and 5,985,589. See also, Vanhaesebroeck et al., *Proc Natl Acad Sci USA*, 94:4330-5 (1997), and international publication WO 97/46688.

In each of the PI3K $\alpha$ ,  $\beta$ , and  $\delta$  subtypes, the p85 subunit acts to localize PI 3-kinase to the plasma membrane by the interaction of its SH2 domain with phosphorylated tyrosine residues (present in an appropriate sequence context) in target proteins (Rameh et al., *Cell*, 83:821-30 (1995)). Two isoforms of p85 have been identified, p85 $\alpha$ , which is ubiquitously expressed, and p85 $\beta$ , which is primarily found in the brain and lymphoid tissues (Volinia et al., *Oncogene*, 7:789-93 (1992)). Association of the p85 subunit to the PI 3-kinase p110 $\alpha$ ,  $\beta$ , or  $\delta$  catalytic subunits appears to be required for the catalytic activity and stability of these enzymes. In

- 5 -

addition, the binding of Ras proteins also upregulates PI 3-kinase activity.

The cloning of p110 $\gamma$  revealed still further complexity within the PI3K family of enzymes (Stoyanov et al., *Science*, 269:690-93 (1995)). The p110 $\gamma$  isoform is closely related to p110 $\alpha$  and p110 $\beta$  (45-48% identity in the catalytic domain), but as noted does not make use of p85 as a targeting subunit. Instead, p110 $\gamma$  contains an additional domain termed a "pleckstrin homology domain" near its amino terminus. This domain allows interaction of p110 $\gamma$  with the  $\beta\gamma$  subunits of heterotrimeric G proteins and this interaction appears to regulate its activity.

The p101 regulatory subunit for PI3K $\gamma$  was originally cloned in swine, and the human ortholog identified subsequently (Krugmann et al., *J Biol Chem*, 274:17152-8 (1999)). Interaction between the N-terminal region of p101 with the N-terminal region of p110 $\gamma$  appears to be critical for the PI3K $\gamma$  activation through G $\beta\gamma$  mentioned above.

A constitutively active PI3K polypeptide is described in international publication WO 96/25488. This publication discloses preparation of a chimeric fusion protein in which a 102-residue fragment of p85 known as the inter-SH2 (iSH2) region is fused through a linker region to the N-terminus of murine p110. The p85 iSH2 domain apparently is able to activate PI3K activity in a manner comparable to intact p85 (Klippel et al., *Mol Cell Biol*, 14:2675-85 (1994)).

Thus, PI 3-kinases can be defined by their amino acid identity or by their activity. Addi-

- 6 -

tional members of this growing gene family include more distantly related lipid and protein kinases including Vps34 TOR1, and TOR2 of *Saccharomyces cerevisiae* (and their mammalian homologs such as FRAP and mTOR), the ataxia telangiectasia gene product (ATR) and the catalytic subunit of DNA-dependent protein kinase (DNA-PK). See generally, Hunter, *Cell*, 83:1-4 (1995).


PI 3-kinase also appears to be involved in a number of aspects of leukocyte activation. A p85-associated PI 3-kinase activity has been shown to physically associate with the cytoplasmic domain of CD28, which is an important costimulatory molecule for the activation of T-cells in response to antigen (Pages et al., *Nature*, 369:327-29 (1994); Rudd, *Immunity*, 4:527-34 (1996)). Activation of T cells through CD28 lowers the threshold for activation by antigen and increases the magnitude and duration of the proliferative response. These effects are linked to increases in the transcription of a number of genes including interleukin-2 (IL2), an important T cell growth factor (Fraser et al., *Science*, 251:313-16 (1991)). Mutation of CD28 such that it can no longer interact with PI 3-kinase leads to a failure to initiate IL2 production, suggesting a critical role for PI 3-kinase in T cell activation.

Specific inhibitors against individual members of a family of enzymes provide invaluable tools for deciphering functions of each enzyme. Two compounds, LY294002 and wortmannin, have been widely used as PI 3-kinase inhibitors. These compounds, however, are nonspecific PI3K inhibitors, as they do not distinguish among the four members of Class I PI

- 7 -


3-kinases. For example, the  $IC_{50}$  values of wortmannin against each of the various Class I PI 3-kinases are in the range of 1-10 nM. Similarly, the  $IC_{50}$  values for LY294002 against each of these PI 3-kinases is about 1  $\mu$ M (Fruman et al., *Ann Rev Biochem*, 67:481-507 (1998)). Hence, the utility of these compounds in studying the roles of individual Class I PI 3-kinases is limited.

Based on studies using wortmannin, there is evidence that PI 3-kinase function also is required for some aspects of leukocyte signaling through G-protein coupled receptors (Thelen et al., *Proc Natl Acad Sci USA*, 91:4960-64 (1994)). Moreover, it has been shown that wortmannin and LY294002 block neutrophil migration and superoxide release. However, inasmuch as these compounds do not distinguish among the various isoforms of PI3K, it remains unclear which particular PI3K isoform or isoforms are involved in these phenomena.



LY294002

- 8 -



wortmannin

In view of the above considerations, it is clear that existing knowledge is lacking with respect to structural and functional features of the PI 3-kinase enzymes, including subcellular localization, activation states, substrate affinities, and the like. Moreover, the functions that these enzymes perform in both normal and diseased tissues remains to be elucidated. In particular, the function of PI3K $\delta$  in leukocytes has not previously been characterized, and knowledge concerning its function in human physiology remains limited. The coexpression in these tissues of other PI3K isoforms has heretofore confounded efforts to segregate the activities of each enzyme. Furthermore, separation of the activities of the various PI3K isozymes may not be possible without identification of inhibitors that demonstrate selective inhibition characteristics. Indeed, Applicants are not presently aware that such selective, or better, specific, inhibitors of PI3K isozymes have been demonstrated.

Thus, there exists a need in the art for further structural characterization of the PI3K $\delta$  polypeptide. There also exists a need for func-

- 9 -

tional characterization of PI3K $\delta$ . Furthermore, our understanding of PI3K $\delta$  requires further elaboration of the structural interactions of p110 $\delta$ , both with its regulatory subunit and with other proteins in the cell. There also remains a need for selective or specific inhibitors of PI3K isozymes, in order that the functions of each isozyme can be better characterized. In particular, selective or specific inhibitors of PI3K $\delta$  are desirable for exploring the role of this isozyme and for development of pharmaceuticals to modulate the activity of the isozyme.

One aspect of the present invention is to provide compounds that can inhibit the biological activity of human PI3K $\delta$ . Another aspect of the invention is to provide compounds that inhibit PI3K $\delta$  selectively while having relatively low inhibitory potency against the other PI3K isoforms. Another aspect of the invention is to provide methods of characterizing the function of human PI3K $\delta$ . Another aspect of the invention is to provide methods of selectively modulating human PI3K $\delta$  activity, and thereby promoting medical treatment of diseases mediated by PI3K $\delta$  dysfunction. Other aspects and advantages of the invention will be readily apparent to the artisan having ordinary skill in the art.

#### SUMMARY OF THE INVENTION

It has now been discovered that these and other aspects can be achieved by the present invention, which, in one aspect, is a method for disrupting leukocyte function, comprising contacting leukocytes with a compound that selectively inhibits

- 10 -


phosphatidylinositol 3-kinase delta (PI3K $\delta$ ) activity  
in the leukocytes. According to the method, the  
leukocytes can comprise cells selected from the  
group consisting of neutrophils, B lymphocytes, T  
5 lymphocytes, and basophils.

For example, in cases in which the leuko-  
cytes comprise neutrophils, the method can comprise  
disrupting at least one neutrophil function selected  
from the group consisting of stimulated superoxide  
10 release, stimulated exocytosis, and chemotactic  
migration. Preferably, the method does not substan-  
tially disrupt bacterial phagocytosis or bacterial  
killing by the neutrophils. In cases wherein the  
leukocytes comprise B lymphocytes, the method can  
15 comprise disrupting proliferation of the B lympho-  
cytes or antibody production by the B lymphocytes.  
In cases wherein the leukocytes comprise T lympho-  
cytes, the method can comprise disrupting prolifera-  
tion of the T lymphocytes. In cases wherein the  
20 leukocytes comprise basophils, the method can com-  
prise disrupting histamine release by the basophils.

In the methods of the invention wherein a  
selective PI3K $\delta$  inhibitor is employed, it is pre-  
ferred that the compound be at least about 10-fold  
25 selective for inhibition of PI3K $\delta$  relative to other  
Type I PI3K isoforms in a cell-based assay. More  
preferably, the compound is at least about 20-fold  
selective for inhibition of PI3K $\delta$  relative to other  
Type I PI3K isoforms in a cell-based assay. Still  
30 more preferably, the compound is at least about 50-  
fold selective for inhibition of PI3K $\delta$  relative to  
other Type I PI3K isoforms in a biochemical assay.

- 11 -

Preferred selective compounds useful according to the methods include compounds having the structure (I):



(I)

wherein A is an optionally substituted monocyclic or bicyclic ring system containing at least two nitrogen atoms, and at least one ring of the system is aromatic;

X is selected from the group consisting of  $C(R^b)_2$ ,  $CH_2CHR^b$ , and  $CH=C(R^b)$ ;

Y is selected from the group consisting of null, S, SO, SO<sub>2</sub>, NH, O, C(=O), OC(=O), C(=O)O, and NHC(=O)CH<sub>2</sub>S;

R<sup>1</sup> and R<sup>2</sup>, independently, are selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, aryl, heteroaryl, halo, NHC(=O)C<sub>1-3</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, NO<sub>2</sub>, OR<sup>a</sup>, CF<sub>3</sub>, OCF<sub>3</sub>, N(R<sup>a</sup>)<sub>2</sub>, CN, OC(=O)R<sup>a</sup>, C(=O)R<sup>a</sup>, C(=O)OR<sup>a</sup>, arylOR<sup>b</sup>, Het, NR<sup>a</sup>C(=O)C<sub>1-3</sub>alkyleneC(=O)OR<sup>a</sup>, arylOC<sub>1-3</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, arylOC(=O)R<sup>a</sup>, C<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>, OC<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>, C<sub>1-4</sub>alkyleneOC<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>, C(=O)NR<sup>a</sup>SO<sub>2</sub>R<sup>a</sup>, C<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, C<sub>2-6</sub>alkenyleneN(R<sup>a</sup>)<sub>2</sub>, C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneOR<sup>a</sup>, C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneHet, OC<sub>2-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, OC<sub>1-4</sub>alkyleneCH(OR<sup>b</sup>)CH<sub>2</sub>N(R<sup>a</sup>)<sub>2</sub>, OC<sub>1-4</sub>alkyleneHet, OC<sub>2-4</sub>alkyleneOR<sup>a</sup>, OC<sub>2-4</sub>alkyleneNR<sup>a</sup>C(=O)OR<sup>a</sup>,



- 12 -

$\text{NR}^a\text{C}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{NR}^a\text{C}(=\text{O})\text{R}^a$ ,  $\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$ ,  
 $\text{N}(\text{SO}_2\text{C}_{1-4}\text{alkyl})_2$ ,  $\text{NR}^a(\text{SO}_2\text{C}_{1-4}\text{alkyl})$ ,  $\text{SO}_2\text{N}(\text{R}^a)_2$ ,  $\text{OSO}_2\text{CF}_3$ ,  
 $\text{C}_{1-3}\text{alkylenearyl}$ ,  $\text{C}_{1-4}\text{alkyleneHet}$ ,  $\text{C}_{1-6}\text{alkyleneOR}^b$ ,  
 $\text{C}_{1-3}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{C}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{NHC}(=\text{O})\text{C}_1\text{-C}_3\text{alkylene-}$   
 5  $\text{aryl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{3-8}\text{heterocycloalkyl}$ ,  $\text{arylOC}_{1-3}\text{-}$   
 $\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{arylOC}(=\text{O})\text{R}^b$ ,  $\text{NHC}(=\text{O})\text{C}_{1-3}\text{alkyleneC}_{3-8}\text{-}$   
 $\text{heterocycloalkyl}$ ,  $\text{NHC}(=\text{O})\text{C}_{1-3}\text{alkyleneHet}$ ,  $\text{OC}_{1-4}\text{al-}$   
 $\text{alkyleneOC}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^b$ ,  $\text{C}(=\text{O})\text{C}_{1-4}\text{alkyleneHet}$ , and  
 $\text{NHC}(=\text{O})\text{haloC}_{1-6}\text{alkyl}$ ;

10 or  $\text{R}^1$  and  $\text{R}^2$  are taken together to form a  
 3- or 4-membered alkylene or alkenylene chain  
 component of a 5- or 6-membered ring, optionally  
 containing at least one heteroatom;

$\text{R}^3$  is selected from the group consisting of  
 15 optionally substituted hydrogen,  $\text{C}_{1-6}\text{alkyl}$ ,  $\text{C}_{3-8}\text{cyclo-}$   
 $\text{alkyl}$ ,  $\text{C}_{3-8}\text{heterocycloalkyl}$ ,  $\text{C}_{1-4}\text{alkylenecycloalkyl}$ ,  
 $\text{C}_{2-6}\text{alkenyl}$ ,  $\text{C}_{1-3}\text{alkylenearyl}$ ,  $\text{arylC}_{1-3}\text{alkyl}$ ,  $\text{C}(=\text{O})\text{R}^a$ ,  
 $\text{aryl}$ , heteroaryl,  $\text{C}(=\text{O})\text{OR}^a$ ,  $\text{C}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{C}(=\text{S})\text{N}(\text{R}^a)_2$ ,  
 $\text{SO}_2\text{R}^a$ ,  $\text{SO}_2\text{N}(\text{R}^a)_2$ ,  $\text{S}(=\text{O})\text{R}^a$ ,  $\text{S}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{C}(=\text{O})\text{NR}^a\text{C}_{1-4}\text{-}$   
 20  $\text{alkyleneOR}^a$ ,  $\text{C}(=\text{O})\text{NR}^a\text{C}_{1-4}\text{alkyleneHet}$ ,  $\text{C}(=\text{O})\text{C}_{1-4}\text{alkyl-}$   
 $\text{enearyl}$ ,  $\text{C}(=\text{O})\text{C}_{1-4}\text{alkyleneheteroaryl}$ ,  $\text{C}_{1-4}\text{alkylenearyl}$   
 optionally substituted with one or more of halo,  
 $\text{SO}_2\text{N}(\text{R}^a)_2$ ,  $\text{N}(\text{R}^a)_2$ ,  $\text{C}(=\text{O})\text{OR}^a$ ,  $\text{NR}^a\text{SO}_2\text{CF}_3$ ,  $\text{CN}$ ,  $\text{NO}_2$ ,  $\text{C}(=\text{O})\text{R}^a$ ,  
 $\text{OR}^a$ ,  $\text{C}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ , and  $\text{OC}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{C}_{1-4}\text{-}$   
 25  $\text{alkyleneheteroaryl}$ ,  $\text{C}_{1-4}\text{alkyleneHet}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{-}$   
 $\text{C}_{1-4}\text{alkylenearyl}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{C}_{1-4}\text{alkylenehetero-}$   
 $\text{aryl}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{Het}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{N}(\text{R}^a)_2$ ,  
 $\text{C}_{1-4}\text{alkyleneOR}^a$ ,  $\text{C}_{1-4}\text{alkyleneNR}^a\text{C}(=\text{O})\text{R}^a$ ,  $\text{C}_{1-4}\text{alkyleneO-}$   
 $\text{C}_{1-4}\text{alkyleneOR}^a$ ,  $\text{C}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{-}$   
 30  $\text{OR}^a$ , and  $\text{C}_{1-4}\text{alkyleneOC}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^a$ ;

$\text{R}^a$  is selected from the group consisting of  
 hydrogen,  $\text{C}_{1-6}\text{alkyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{3-8}\text{heterocyclo-}$   
 $\text{alkyl}$ ,  $\text{C}_{1-3}\text{alkyleneN}(\text{R}^c)_2$ ,  $\text{aryl}$ ,  $\text{arylC}_{1-3}\text{alkyl}$ ,  $\text{C}_{1-3}\text{-}$

- 13 -

alkylenearyl, heteroaryl, heteroarylC<sub>1-3</sub>alkyl, and C<sub>1-3</sub>alkyleneheteroaryl;

or two R<sup>a</sup> groups are taken together to form a 5- or 6-membered ring, optionally containing at least one heteroatom;

R<sup>b</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, heteroC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkyleneheteroC<sub>1-3</sub>alkyl, arylheteroC<sub>1-3</sub>alkyl, aryl, heteroaryl, arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, and C<sub>1-3</sub>alkyleneheteroaryl;

R<sup>c</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, aryl, and heteroaryl;

Het is a 5- or 6-membered heterocyclic ring, saturated or partially or fully unsaturated, containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted with C<sub>1-4</sub>alkyl or C(=O)OR<sup>a</sup>;

and pharmaceutically acceptable salts and solvates (e.g., hydrates) thereof,

wherein the compound has at least about a 10-fold selective inhibition for PI3Kδ relative other Type-I PI3K isoforms in a cell-based assay.

In another embodiment, the invention is a method for treating a medical condition mediated by neutrophils, comprising administering to an animal in need thereof an effective amount of a compound that selectively inhibits phosphatidylinositol 3-kinase delta (PI3Kδ) activity in the neutrophils. Exemplary medical conditions that can be treated according to the method include those conditions characterized by an undesirable neutrophil function selected from the group consisting of stimulated

- 14 -

superoxide release, stimulated exocytosis, and chemotactic migration. Preferably, according to the method, phagocytic activity or bacterial killing by the neutrophils is substantially uninhibited.

5           In another embodiment, the invention is a method for disrupting a function of osteoclasts comprising contacting osteoclasts with a compound that selectively inhibits phosphatidylinositol 3-kinase delta (PI3K $\delta$ ) activity in the osteoclasts.  
10       According to the method, the compound can comprise a moiety that preferentially binds to bone.

          In another embodiment, the invention is a method of ameliorating a bone-resorption disorder in an animal in need thereof comprising administering  
15       to the animal an effective amount of a compound that inhibits phosphatidylinositol 3-kinase delta (PI3K $\delta$ ) activity in osteoclasts of the animal. A preferred bone-resorption disorder amenable to treatment according to the method is osteoporosis.


20           In another embodiment, the invention is a method for inhibiting the growth or proliferation of cancer cells of hematopoietic origin, comprising contacting the cancer cells with a compound that selectively inhibits phosphatidylinositol 3-kinase  
25       delta (PI3K $\delta$ ) activity in the cancer cells. The method can be advantageous in inhibiting the growth or proliferation of cancers selected from the group consisting of lymphomas, multiple myelomas, and leukemias.

30           In another embodiment, the invention is a method of inhibiting kinase activity of a phosphatidylinositol 3-kinase delta (PI3K $\delta$ ) polypeptide,

- 15 -

comprising contacting the PI3K $\delta$  polypeptide with a compound having the generic structure (I).

Preferred compounds useful according to the method include compounds selected from the group consisting of:



(II)

wherein Y is selected from the group consisting of null, S, and NH;

R<sup>4</sup> is selected from the group consisting of H, halo, NO<sub>2</sub>, OH, OCH<sub>3</sub>, CH<sub>3</sub>, and CF<sub>3</sub>;

R<sup>5</sup> is selected from the group consisting of H, OCH<sub>3</sub>, and halo;


or R<sup>4</sup> and R<sup>5</sup> together with C-6 and C-7 of the quinazoline ring system define a 5- or 6-membered aromatic ring optionally containing one or more O, N, or S atoms;

R<sup>6</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl, halophenyl, alkoxyphenyl, alkylphenyl, biphenyl, benzyl, pyridinyl, 4-methylpiperazinyl, C(=O)OC<sub>2</sub>H<sub>5</sub>, and morpholinyl;

R<sup>d</sup>, independently, is selected from the group consisting of NH<sub>2</sub>, halo, C<sub>1-3</sub>alkyl, S(C<sub>1-3</sub>alkyl),

- 16 -

OH,  $\text{NH}(\text{C}_{1-3}\text{alkyl})$ ,  $\text{N}(\text{C}_{1-3}\text{alkyl})_2$ ,  $\text{NH}(\text{C}_{1-3}\text{alkyleneph-en-yl})$ , and




; and

$q$  is 1 or 2,

provided that at least one of  $\text{R}^4$  and  $\text{R}^5$  is other than H when  $\text{R}^6$  is phenyl or 2-chlorophenyl.

More preferably, the compound is selected from the group consisting of:



(III)

wherein  $Y$  is selected from the group consisting of null, S, and NH;

$\text{R}^7$  is selected from the group consisting of H, halo, OH,  $\text{OCH}_3$ ,  $\text{CH}_3$ , and  $\text{CF}_3$ ;

- 17 -

R<sup>8</sup> is selected from the group consisting of is H, OCH<sub>3</sub>, and halo;

or R<sup>7</sup> and R<sup>8</sup> together with C-6 and C-7 of the quinazoline ring system define a 5- or 6-  
5 membered aromatic ring optionally containing one or more O, N, or S atoms;

R<sup>9</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl, halophenyl, alkylphenyl, biphenyl, benzyl, pyridinyl, 4-methylpiperazinyl, C(=O)-  
10 OC<sub>2</sub>H<sub>5</sub>, and morpholinyl;

R<sup>d</sup>, independently, is selected from the group consisting of NH<sub>2</sub>, halo, C<sub>1-3</sub>alkyl, S(C<sub>1-3</sub>alkyl), OH, NH(C<sub>1-3</sub>alkyl), N(C<sub>1-3</sub>alkyl)<sub>2</sub>, NH(C<sub>1-3</sub>alkylenephenyl); and

15 q is 1 or 2,  
provided that at least one of R<sup>7</sup> and R<sup>8</sup> is different from 6-halo or 6,7-dimethoxy groups, and that R<sup>9</sup> is different from 4-chlorophenyl.


In another embodiment, the invention is a  
20 method for disrupting leukocyte function, comprising contacting leukocytes with a compound having a general structure (I).

In another embodiment, the invention is a class of compounds that have been observed to  
25 inhibit PI3K $\delta$  activity in biochemical and cell-based assays, and are expected to exhibit therapeutic benefit in medical conditions in which PI3K $\delta$  activity is excessive or undesirable. Thus, the invention provides a class of compounds having the  
30 structure (II).

Preferably, the compounds have a general structure (IV)

- 18 -

5



10

(IV)

wherein Y is selected from the group consisting of null, S, and NH;

15

R<sup>10</sup> is selected from the group consisting of H, halo, OH, OCH<sub>3</sub>, CH<sub>3</sub>, and CF<sub>3</sub>;

R<sup>11</sup> is selected from the group consisting of H, OCH<sub>3</sub>, and halo;

20

or R<sup>10</sup> and R<sup>11</sup> together with C-6 and C-7 of the quinazoline ring system define a 5- or 6-membered aromatic ring optionally containing one or more O, N, or S atoms;

25

R<sup>12</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl, halophenyl, alkylphenyl, biphenyl, benzyl, pyridinyl, 4-methylpiperazinyl, C(=O)C<sub>2</sub>H<sub>5</sub>, and morpholinyl;

30

R<sup>d</sup>, independently, is selected from the group consisting of NH<sub>2</sub>, halo, C<sub>1-3</sub>alkyl, S(C<sub>1-3</sub>alkyl), OH, NH(C<sub>1-3</sub>alkyl), N(C<sub>1-3</sub>alkyl)<sub>2</sub>, NH(C<sub>1-3</sub>alkyleneph-en-yl), and

q is 1 or 2,

provided that:

- 19 -

(a) at least one of  $R^{10}$  and  $R^{11}$  is different from 6-halo or 6,7-dimethoxy groups;

(b)  $R^{12}$  is different from 4-chlorophenyl; and

5 (c) at least one of  $R^{10}$  and  $R^{11}$  is different from H when  $R^{12}$  is phenyl or 2-chlorophenyl and X is S.

These and other features and advantages of the present invention will be appreciated from the detailed description and examples that are set forth herein. The detailed description and examples are provided to enhance the understanding of the invention, but are not intended to limit the scope of the invention.

15

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the effect of a selective PI3K $\delta$  inhibitor of the invention on the activity of three PI3K isoforms.

20

Figure 2 shows the effect of a selective PI3K $\delta$  inhibitor on superoxide generation by human neutrophils in the presence of TNF or IgG.

Figure 3 shows the effect of a selective PI3K $\delta$  inhibitor on superoxide generation by human neutrophils in the presence of TNF or fMLP.

25

Figure 4 shows the effect of a selective PI3K $\delta$  inhibitor on elastase exocytosis in the presence of fMLP by human neutrophils.

30

Figure 5 shows the effect of a selective PI3K $\delta$  inhibitor on fMLP-induced chemotaxis by human neutrophils.



- 20 -

Figure 6 shows that a selective PI3K $\delta$  inhibitor does not affect phagocytosis and killing of *S. aureus* by neutrophils.

5 Figure 7 shows the effect of a selective PI3K $\delta$  inhibitor on proliferation and antibody production by human B lymphocytes.

Figure 8 shows the effect of a selective PI3K $\delta$  inhibitor on anti-IgM stimulated mouse splenic B lymphocyte proliferation.

10 Figure 9 shows the effect of a selective PI3K $\delta$  inhibitor on elastase exocytosis in an animal model.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

15

The invention provides compounds that selectively inhibit the activity of PI3K $\delta$ . The invention further provides methods of inhibiting PI3K $\delta$  activity, including methods of selectively  
20 modulating the activity of the PI3K $\delta$  isozyme in cells, especially leukocytes, osteoclasts, and cancer cells. The methods include *in vitro*, *in vivo*, and *ex vivo* applications.

Of particular benefit are methods of  
25 selectively modulating PI3K $\delta$  activity in the clinical setting in order to ameliorate disease or disorders mediated by PI3K $\delta$  activity. Thus, treatment of diseases or disorders characterized by excessive or inappropriate PI3K $\delta$  activity can be treated  
30 through use of selective modulators of PI3K $\delta$  according to the invention.

Other methods of the invention include enabling the further characterization of the physio-

- 21 -

logical role of the isozyme. Moreover, the invention provides pharmaceutical compositions comprising selective PI3K $\delta$  inhibitors. Also provided are articles of manufacture comprising a selective PI3K $\delta$  inhibitor compound (or a pharmaceutical composition comprising the compound) and instructions for using the compound. Details of these and other useful embodiments of the invention are now described.

The methods described herein benefit from the use of compounds that selectively inhibit, and preferably specifically inhibit, the activity of PI3K $\delta$  in cells, including cells *in vitro*, *in vivo*, or *ex vivo*. Cells useful in the methods include those that express endogenous PI3K $\delta$ , wherein endogenous indicates that the cells express PI3K $\delta$  absent recombinant introduction into the cells of one or more polynucleotides encoding a PI3K $\delta$  polypeptide or a biologically active fragment thereof. Methods also encompass use of cells that express exogenous PI3K $\delta$ , wherein one or more polynucleotides encoding PI3K $\delta$  or a biologically active fragment thereof have been introduced into the cell using recombinant procedures.

Of particular advantage, the cells can be *in vivo*, i.e., in a living subject, e.g., an animal or human, wherein a PI3K $\delta$  inhibitor can be used as a therapeutic to inhibit PI3K $\delta$  activity in the subject. Alternatively, the cells can be isolated as discrete cells or in a tissue, for *ex vivo* or *in vitro* methods. *In vitro* methods also encompassed by the invention can comprise the step of contacting a PI3K $\delta$  enzyme or a biologically active fragment thereof with an inhibitor compound of the invention.

- 22 -

The PI3K $\delta$  enzyme can include a purified and isolated enzyme, wherein the enzyme is isolated from a natural source (e.g., cells or tissues that normally express a PI3K $\delta$  polypeptide absent modification by recombinant technology) or isolated from cells modified by recombinant techniques to express exogenous enzyme.

The term "selective PI3K $\delta$  inhibitor" as used herein refers to a compound that inhibits the PI3K $\delta$  isozyme more effectively than other isozymes of the PI3K family. A "selective PI3K $\delta$  inhibitor" compound is understood to be more selective for PI3K $\delta$  than compounds conventionally and generically designated PI3K inhibitors, e.g., wortmannin or LY294002. Concomitantly, wortmannin and LY294002 are deemed "nonselective PI3K inhibitors." Compounds of any type that selectively negatively regulate PI3K $\delta$  expression or activity can be used as selective PI3K $\delta$  inhibitors in the methods of the invention. Moreover, compounds of any type that selectively negatively regulate PI3K $\delta$  expression or activity and that possess acceptable pharmacological properties can be used as selective PI3K $\delta$  inhibitors in the therapeutic methods of the invention.

The relative efficacies of compounds as inhibitors of an enzyme activity (or other biological activity) can be established by determining the concentrations at which each compound inhibits the activity to a predefined extent and then comparing the results. Typically, the preferred determination is the concentration that inhibits 50% of the activity in a biochemical assay, i.e., the 50% inhibitory concentration or "IC<sub>50</sub>." IC<sub>50</sub> determinations can be

- 23 -

accomplished using conventional techniques known in the art. In general, an  $IC_{50}$  can be determined by measuring the activity of a given enzyme in the presence of a range of concentrations of the inhibitor under study. The experimentally obtained values of enzyme activity then are plotted against the inhibitor concentrations used. The concentration of the inhibitor that shows 50% enzyme activity (as compared to the activity in the absence of any inhibitor) is taken as the  $IC_{50}$  value. Analogously, other inhibitory concentrations can be defined through appropriate determinations of activity. For example, in some settings it can be desirable to establish a 90% inhibitory concentration, i.e.,  $IC_{90}$ , etc.

Accordingly, a "selective PI3K $\delta$  inhibitor" alternatively can be understood to refer to a compound that exhibits a 50% inhibitory concentration ( $IC_{50}$ ) with respect to PI3K $\delta$  that is at least at least 10-fold, preferably at least 20-fold, and more preferably at least 30-fold, lower than the  $IC_{50}$  value with respect to any or all of the other Class I PI3K family members. The term "specific PI3K $\delta$  inhibitor" can be understood to refer to a selective PI3K $\delta$  inhibitor compound that exhibits an  $IC_{50}$  with respect to PI3K $\delta$  that is at least 50-fold, preferably at least 100-fold, more preferably at least 200-fold, and still more preferably at least 500-fold, lower than the  $IC_{50}$  with respect to any or all of the other PI3K Class I family members.

Among other things, the invention provides methods of inhibiting leukocyte function. More particularly, the invention provides methods of inhib-

- 24 -

iting or suppressing functions of neutrophils and T and B lymphocytes. With respect to neutrophils, it has unexpectedly been found that inhibition of PI3K $\delta$  activity inhibits functions of neutrophils. For  
5 example, it has been observed that the compounds of the invention elicit inhibition of classical neutrophil functions such as stimulated superoxide release, stimulated exocytosis, and chemotactic migration. However, it has been further observed that  
10 the method of the invention permits suppression of certain functions of neutrophils, while not substantially affecting other functions of these cells. For example, it has been observed that phagocytosis of bacteria by neutrophils is not substantially  
15 inhibited by the selective PI3K $\delta$  inhibitor compounds of the invention.

Thus, the invention includes methods for inhibiting neutrophil functions, without substantially inhibiting phagocytosis of bacteria. Neutrophil functions suitable for inhibition according to  
20 the method include any function that is mediated by PI3K $\delta$  activity or expression. Such functions include, without limitation, stimulated superoxide release, stimulated exocytosis or degranulation, chemotactic migration, adhesion to vascular endothelium (e.g., tethering/rolling of neutrophils, triggering of neutrophil activity, and/or latching of neutrophils to endothelium), transmural diapedesis or emigration through the endothelium to peripheral tissues. In general, these functions can be  
25 collectively termed "inflammatory functions," as they are typically related to neutrophil response to inflammation. The inflammatory functions of neutro-  
30

- 25 -

phils can be distinguished from the bacterial killing functions exhibited by these cells, e.g., phagocytosis and killing of bacteria. Accordingly, the invention further includes methods of treating disease states in which one or more of the inflammatory functions of neutrophils are abnormal or undesirable.

It has further been established through the invention that PI3K $\delta$  plays a role in the stimulated proliferation of lymphocytes, including B cells and T cells. Moreover, PI3K $\delta$  appears to play a role in stimulated secretion of antibodies by B cells. Selective PI3K $\delta$  inhibitor compounds of the invention have been employed to establish that these phenomena can be abrogated by inhibition of PI3K $\delta$ . Thus, the invention includes methods of inhibiting lymphocyte proliferation, and methods of inhibiting antibody production by B lymphocytes. Other methods enabled by the invention include methods of treating disease states in which one or more of these lymphocyte functions are abnormal or undesirable.

It has now been determined that PI3K $\delta$  activity can be inhibited selectively or specifically to facilitate treatment of a PI3K $\delta$ -mediated disease while reducing or eliminating complications that are typically associated with concomitant inhibition of the activity of other Class I PI 3-kinases. To illustrate this embodiment, methods of the invention can be practiced using members of a class of compounds that have been found to exhibit selective inhibition of PI3K $\delta$  relative to other PI3K isoforms.

- 26 -

The methods of this embodiment can be practiced using compounds having the general structure (III). Preferred methods employ compounds that have been empirically determined to exhibit at least

5 10-fold selective inhibition of PI3K $\delta$  relative to other PI3K isoforms. For example, the methods can be practiced using the following compounds:

3-(2-isopropylphenyl)-5-methyl-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one;

10 5-chloro-2-(9H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one;

5-chloro-3-(2-fluorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-(2-fluorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

15 3-(2-methoxyphenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-(2,6-dichlorophenyl)-5-methyl-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one;

20 3-(2-chlorophenyl)-6-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

5-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-(2-chlorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

25 3-(3-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-(2-chlorophenyl)-5-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

30 3-benzyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-butyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

- 27 -

- 3-(2-chlorophenyl)-7-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-morpholin-4-yl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one, acetate salt;
- 5 8-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-(2-chlorophenyl)-6,7-difluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-(2-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 10 6-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-(3-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 15 2-(9H-purin-6-ylsulfanylmethyl)-3-pyridin-4-yl-3H-quinazolin-4-one;  
3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-trifluoromethyl-3H-quinazolin-4-one;  
3-benzyl-5-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 20 3-(4-methylpiperazin-1-yl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one, acetate salt;  
3-(2-chlorophenyl)-6-hydroxy-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 25 [5-fluoro-4-oxo-2-(9H-purin-6-ylsulfanylmethyl)-4H-quinazolin-3-yl]acetic acid ethyl ester;  
3-(2,4-dimethoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-biphenyl-2-yl-5-chloro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 30 2-(6-aminopurin-9-ylmethyl)-3-(2-isopropylphenyl)-5-methyl-3H-quinazolin-4-one;




- 28 -

- 2-(6-aminopurin-9-ylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-3-biphenyl-2-yl-5-chloro-3H-quinazolin-4-one;
- 5 5-chloro-3-(2-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-3-(2-fluorophenyl)-5-methyl-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-fluorophenyl)-3H-quinazolin-4-one;
- 10 2-(6-aminopurin-9-ylmethyl)-8-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one;
- 15 2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-5-methyl-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-5-fluoro-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-3-benzyl-5-fluoro-3H-quinazolin-4-one;
- 20 2-(6-aminopurin-9-ylmethyl)-3-butyl-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-3-morpholin-4-yl-3H-quinazolin-4-one;
- 25 2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-7-fluoro-3H-quinazolin-4-one;
- 3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 3-phenyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 30 2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-isopropylphenyl)-3H-quinazolin-4-one; and

- 29 -

2-(6-aminopurin-9-ylmethyl)-5-chloro-3-o-tolyl-3H-quinazolin-4-one.

It has further been determined that the methods of the invention can be advantageously practiced using members of a class of compounds that exhibit PI3K $\delta$  inhibitory activity, thereby facilitating inhibitions of PI3K $\delta$  activity in diseases mediated thereby. For example, in this embodiment, the methods of the invention can be practiced using compounds having the general structure (I).



(I)

wherein A is an optionally substituted monocyclic or bicyclic ring system containing at least two nitrogen atoms, and at least one ring of the system is aromatic;

X is selected from the group consisting of C(R<sup>b</sup>)<sub>2</sub>, CH<sub>2</sub>CHR<sup>b</sup>, and CH=C(R<sup>b</sup>);

Y is selected from the group consisting of null, S, SO, SO<sub>2</sub>, NH, O, C(=O), OC(=O), C(=O)O, and NHC(=O)CH<sub>2</sub>S;

R<sup>1</sup> and R<sup>2</sup>, independently, are selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, aryl, heteroaryl, halo, NHC(=O)C<sub>1-3</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, NO<sub>2</sub>, OR<sup>a</sup>, CF<sub>3</sub>, OCF<sub>3</sub>, N(R<sup>a</sup>)<sub>2</sub>, CN, OC(=O)R<sup>a</sup>, C(=O)R<sup>a</sup>, C(=O)OR<sup>a</sup>, arylOR<sup>b</sup>, Het, NR<sup>a</sup>C(=O)C<sub>1-3</sub>alkyleneC(=O)OR<sup>a</sup>, arylOC<sub>1-3</sub>-

- 30 -

alkyleneN(R<sup>a</sup>)<sub>2</sub>, arylOC(=O)R<sup>a</sup>, C<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>,  
 OC<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>, C<sub>1-4</sub>alkyleneOC<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>,  
 C(=O)NR<sup>a</sup>SO<sub>2</sub>R<sup>a</sup>, C<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, C<sub>2-6</sub>alkenyleneN(R<sup>a</sup>)<sub>2</sub>,  
 C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneOR<sup>a</sup>, C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneHet, OC<sub>2-4</sub>-  
 5 alkyleneN(R<sup>a</sup>)<sub>2</sub>, OC<sub>1-4</sub>alkyleneCH(OR<sup>b</sup>)CH<sub>2</sub>N(R<sup>a</sup>)<sub>2</sub>, OC<sub>1-4</sub>-  
 alkyleneHet, OC<sub>2-4</sub>alkyleneOR<sup>a</sup>, OC<sub>2-4</sub>alkyleneNR<sup>a</sup>C(=O)OR<sup>a</sup>,  
 NR<sup>a</sup>C<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, NR<sup>a</sup>C(=O)R<sup>a</sup>, NR<sup>a</sup>C(=O)N(R<sup>a</sup>)<sub>2</sub>,  
 N(SO<sub>2</sub>C<sub>1-4</sub>alkyl)<sub>2</sub>, NR<sup>a</sup>(SO<sub>2</sub>C<sub>1-4</sub>alkyl), SO<sub>2</sub>N(R<sup>a</sup>)<sub>2</sub>, OSO<sub>2</sub>CF<sub>3</sub>,  
 C<sub>1-3</sub>alkylenearyl, C<sub>1-4</sub>alkyleneHet, C<sub>1-6</sub>alkyleneOR<sup>b</sup>,  
 10 C<sub>1-3</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, C(=O)N(R<sup>a</sup>)<sub>2</sub>, NHC(=O)C<sub>1-3</sub>alkylene-  
 aryl, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>heterocycloalkyl, arylOC<sub>1-3</sub>-  
 alkyleneN(R<sup>a</sup>)<sub>2</sub>, arylOC(=O)R<sup>b</sup>, NHC(=O)C<sub>1-3</sub>alkyleneC<sub>3-8</sub>-  
 heterocycloalkyl, NHC(=O)C<sub>1-3</sub>alkyleneHet, OC<sub>1-4</sub>alkyl-  
 eneOC<sub>1-4</sub>alkyleneC(=O)OR<sup>b</sup>, C(=O)C<sub>1-4</sub>alkyleneHet, and  
 15 NHC(=O)haloC<sub>1-6</sub>alkyl;

or R<sup>1</sup> and R<sup>2</sup> are taken together to form a  
 3- or 4-membered alkylene or alkenylene chain  
 component of a 5- or 6-membered ring, optionally  
 containing at least one heteroatom;

20 R<sup>3</sup> is selected from the group consisting of  
 optionally substituted hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-8</sub>cyclo-  
 alkyl, C<sub>3-8</sub>heterocycloalkyl, C<sub>1-4</sub>alkylenecycloalkyl,  
 C<sub>2-6</sub>alkenyl, C<sub>1-3</sub>alkylenearyl, arylC<sub>1-3</sub>alkyl, C(=O)R<sup>a</sup>,  
 aryl, heteroaryl, C(=O)OR<sup>a</sup>, C(=O)N(R<sup>a</sup>)<sub>2</sub>, C(=S)N(R<sup>a</sup>)<sub>2</sub>,  
 25 SO<sub>2</sub>R<sup>a</sup>, SO<sub>2</sub>N(R<sup>a</sup>)<sub>2</sub>, S(=O)R<sup>a</sup>, S(=O)N(R<sup>a</sup>)<sub>2</sub>, C(=O)NR<sup>a</sup>C<sub>1-4</sub>-  
 alkyleneOR<sup>a</sup>, C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneHet, C(=O)C<sub>1-4</sub>alkyl-  
 enearyl, C(=O)C<sub>1-4</sub>alkyleneheteroaryl, C<sub>1-4</sub>alkylenearyl  
 optionally substituted with one or more of halo  
 SO<sub>2</sub>N(R<sup>a</sup>)<sub>2</sub>, N(R<sup>a</sup>)<sub>2</sub>, C(=O)OR<sup>a</sup>, NR<sup>a</sup>SO<sub>2</sub>CF<sub>3</sub>, CN, NO<sub>2</sub>, C(=O)R<sup>a</sup>,  
 30 OR<sup>a</sup>, C<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, and OC<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>,  
 C<sub>1-4</sub>alkyleneheteroaryl, C<sub>1-4</sub>alkyleneHet, C<sub>1-4</sub>alkylene-  
 C(=O)C<sub>1-4</sub>alkylenearyl, C<sub>1-4</sub>alkyleneC(=O)C<sub>1-4</sub>alkylene-  
 heteroaryl, C<sub>1-4</sub>alkyleneC(=O)Het, C<sub>1-4</sub>alkyleneC(=O)-

- 31 -

$N(R^a)_2$ ,  $C_{1-4}$ alkylene $OR^a$ ,  $C_{1-4}$ alkylene $NR^aC(=O)R^a$ ,  
 $C_{1-4}$ alkylene $OC_{1-4}$ alkylene $OR^a$ ,  $C_{1-4}$ alkylene $N(R^a)_2$ ,  
 $C_{1-4}$ alkylene $C(=O)OR^a$ , and  $C_{1-4}$ alkylene $OC_{1-4}$ alkylene-  
 $C(=O)OR^a$ ;

5  $R^a$  is selected from the group consisting of  
hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocyclo-  
alkyl,  $C_{1-3}$ alkylene $N(R^c)_2$ , aryl, aryl $C_{1-3}$ alkyl,  
 $C_{1-3}$ alkylenearyl, heteroaryl, heteroaryl $C_{1-3}$ alkyl, and  
 $C_{1-3}$ alkyleneheteroaryl;

10 or two  $R^a$  groups are taken together to form  
a 5- or 6-membered ring, optionally containing at  
least one heteroatom;

$R^b$  is selected from the group consisting of  
hydrogen,  $C_{1-6}$ alkyl, hetero $C_{1-3}$ alkyl,  $C_{1-3}$ alkylenehet-  
15 ero $C_{1-3}$ alkyl, arylhetero $C_{1-3}$ alkyl, aryl, heteroaryl,  
aryl $C_{1-3}$ alkyl, heteroaryl $C_{1-3}$ alkyl,  $C_{1-3}$ alkylenearyl,  
and  $C_{1-3}$ alkyleneheteroaryl;

$R^c$  is selected from the group consisting of  
hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl, aryl, and hetero-  
20 aryl;

Het is a 5- or 6-membered heterocyclic  
ring, saturated or partially or fully unsaturated,  
containing at least one heteroatom selected from the  
group consisting of oxygen, nitrogen, and sulfur,  
25 and optionally substituted with  $C_{1-4}$ alkyl or  $C(=O)OR^a$ ;

and pharmaceutically acceptable salts and  
solvates (e.g., hydrates) thereof.

For example, methods of the invention can  
employ compounds that possess PI3K $\delta$  inhibitory  
30 activity, as follows:

3-(2-isopropylphenyl)-5-methyl-2-(9H-purin-6-yl-  
sulfanylmethyl)-3H-quinazolin-4-one;

- 32 -

- 5-chloro-2-(9H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one;  
5-chloro-3-(2-fluorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
5. 3-(2-fluorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-(2-methoxyphenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-(2,6-dichlorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
10 3-(2-chlorophenyl)-6-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
5-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
15 3-(2-chlorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-(2-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-(2-chlorophenyl)-5-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
20 3-benzyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-butyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
25 3-(2-chlorophenyl)-7-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-morpholin-4-yl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one, acetate salt;  
8-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
30 3-(2-chlorophenyl)-6,7-difluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

- 33 -

- 3-(3-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-quinazolin-4-one;  
6-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfan-  
ylmethyl)-3H-quinazolin-4-one;  
5 3-(3-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-quinazolin-4-one;  
2-(9H-purin-6-ylsulfanylmethyl)-3-pyridin-4-yl-3H-  
quinazolin-4-one;  
3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-  
10 trifluoromethyl-3H-quinazolin-4-one;  
3-benzyl-5-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-quinazolin-4-one;  
3-(4-methylpiperazin-1-yl)-2-(9H-purin-6-ylsulfan-  
ylmethyl)-3H-quinazolin-4-one, acetate salt;  
15 3-(2-chlorophenyl)-6-hydroxy-2-(9H-purin-6-ylsulfan-  
ylmethyl)-3H-quinazolin-4-one;  
[5-fluoro-4-oxo-2-(9H-purin-6-ylsulfanylmethyl)-4H-  
quinazolin-3-yl]acetic acid ethyl ester;  
3-biphenyl-2-yl-5-chloro-2-(9H-purin-6-ylsulfanyl-  
20 methyl)-3H-quinazolin-4-one;  
5-chloro-3-(2-methoxyphenyl)-2-(9H-purin-6-ylsulfan-  
ylmethyl)-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-3-(2-isopropylphenyl)-5-  
methyl-3H-quinazolin-4-one;  
25 2-(6-aminopurin-9-ylmethyl)-5-methyl-3-o-tolyl-3H-  
quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-3-biphenyl-2-yl-t-  
chloro-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-3-(2-fluorophenyl)-5-  
30 methyl-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-fluoro-  
phenyl)-3H-quinazolin-4-one;

- 34 -

- 2-(6-aminopurin-9-ylmethyl)-8-chloro-3-(2-chloro-phenyl)-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-chloro-phenyl)-3H-quinazolin-4-one;
- 5 2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-5-methyl-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-5-fluoro-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-3-benzyl-5-fluoro-3H-quinazolin-4-one;
- 10 2-(6-aminopurin-9-ylmethyl)-3-butyl-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-3-morpholin-4-yl-3H-quinazolin-4-one;
- 15 2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-7-fluoro-3H-quinazolin-4-one;
- 3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 3-phenyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 20 2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one;
- 3-(4-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 25 3-(2-chlorophenyl)-6,7-dimethoxy-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 3-(2-chlorophenyl)-7-nitro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 2-(6-aminopurin-9-ylmethyl)-6-bromo-3-(2-chlorophenyl)-3H-quinazolin-4-one;
- 30 2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-6,7-dimethoxy-3H-quinazolin-4-one;

- 35 -

6-bromo-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;


3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-benzo[g]quinazolin-4-one;

5 2-(6-aminopurin-9-ylmethyl)-5-chloro-3-o-tolyl-3H-quinazolin-4-one; and

2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-methoxyphenyl)-3H-quinazolin-4-one.

10 The invention further provides compounds that are selective inhibitors of PI3K $\delta$  activity. The compounds exhibit inhibition of PI3K $\delta$  in biochemical assays, and selectively disrupt function of PI3K $\delta$ -expressing cells in cell-based assays. As  
15 described elsewhere herein, the compounds of the invention have been demonstrated to inhibit certain functions in neutrophils and other leukocytes, as well as functions of osteoclasts.

In general, compounds provided by the  
20 invention have the general structure (I), a pharmaceutically acceptable salt thereof, or a prodrug thereof:



(I)

30

wherein A is an optionally substituted monocyclic or bicyclic ring system containing at



- 36 -

least two nitrogen atoms, and at least one ring of the system is aromatic;

X is selected from the group consisting of  $C(R^b)_2$ ,  $CH_2CHR^b$ , and  $CH=C(R^b)$ ;

5 Y is selected from the group consisting of null, S, SO,  $SO_2$ , NH, O,  $C(=O)$ ,  $OC(=O)$ ,  $C(=O)O$ , and  $NHC(=O)CH_2S$ ;

$R^1$  and  $R^2$ , independently, are selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl, aryl, heteroaryl, halo,  $NHC(=O)C_{1-3}$ alkylene $N(R^a)_2$ ,  $NO_2$ ,  $OR^a$ ,  $CF_3$ ,  $OCF_3$ ,  $N(R^a)_2$ , CN,  $OC(=O)R^a$ ,  $C(=O)R^a$ ,  $C(=O)OR^a$ ,  $arylOR^b$ , Het,  $NR^aC(=O)C_{1-3}$ alkylene $C(=O)OR^a$ ,  $arylOC_{1-3}$ alkylene $N(R^a)_2$ ,  $arylOC(=O)R^a$ ,  $C_{1-4}$ alkylene $C(=O)OR^a$ ,  $OC_{1-4}$ alkylene $C(=O)OR^a$ ,  $C_{1-4}$ alkylene $OC_{1-4}$ alkylene $C(=O)OR^a$ ,  $C(=O)NR^aSO_2R^a$ ,  $C_{1-4}$ alkylene $N(R^a)_2$ ,  $C_{2-6}$ alkenylene $N(R^a)_2$ ,  $C(=O)NR^aC_{1-4}$ alkylene $OR^a$ ,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $OC_{2-4}$ alkylene $N(R^a)_2$ ,  $OC_{1-4}$ alkylene $CH(OR^b)CH_2N(R^a)_2$ ,  $OC_{1-4}$ alkyleneHet,  $OC_{2-4}$ alkylene $OR^a$ ,  $OC_{2-4}$ alkylene $NR^aC(=O)OR^a$ ,  $NR^aC_{1-4}$ alkylene $N(R^a)_2$ ,  $NR^aC(=O)R^a$ ,  $NR^aC(=O)N(R^a)_2$ ,  $N(SO_2C_{1-4}alkyl)_2$ ,  $NR^a(SO_2C_{1-4}alkyl)$ ,  $SO_2N(R^a)_2$ ,  $OSO_2CF_3$ ,  $C_{1-3}$ alkylenearyl,  $C_{1-4}$ alkyleneHet,  $C_{1-6}$ alkylene $OR^b$ ,  $C_{1-3}$ alkylene $N(R^a)_2$ ,  $C(=O)N(R^a)_2$ ,  $NHC(=O)C_{1-3}$ alkylenearyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocycloalkyl,  $arylOC_{1-3}$ alkylene $N(R^a)_2$ ,  $arylOC(=O)R^b$ ,  $NHC(=O)C_{1-3}$ alkylene $C_{3-8}$ heterocycloalkyl,  $NHC(=O)C_{1-3}$ alkyleneHet,  $OC_{1-4}$ alkylene $OC_{1-4}$ alkylene $C(=O)OR^b$ ,  $C(=O)C_{1-4}$ alkyleneHet, and  $NHC(=O)haloC_{1-6}alkyl$ ;

or  $R^1$  and  $R^2$  are taken together to form a 3- or 4-membered alkylene or alkenylene chain component of a 5- or 6-membered ring, optionally containing at least one heteroatom;

$R^3$  is selected from the group consisting of optionally substituted hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cyclo-

- 37 -

alkyl, C<sub>3-8</sub>heterocycloalkyl, C<sub>1-4</sub>alkylenecycloalkyl, C<sub>2-6</sub>alkenyl, C<sub>1-3</sub>alkylenearyl, arylC<sub>1-3</sub>alkyl, C(=O)R<sup>a</sup>, aryl, heteroaryl, C(=O)OR<sup>a</sup>, C(=O)N(R<sup>a</sup>)<sub>2</sub>, C(=S)N(R<sup>a</sup>)<sub>2</sub>, SO<sub>2</sub>R<sup>a</sup>, SO<sub>2</sub>N(R<sup>a</sup>)<sub>2</sub>, S(=O)R<sup>a</sup>, S(=O)N(R<sup>a</sup>)<sub>2</sub>, C(=O)NR<sup>a</sup>C<sub>1-4</sub>-

5 alkyleneor<sup>a</sup>, C(=O)NR<sup>a</sup>C<sub>1-4</sub>alkyleneHet, C(=O)C<sub>1-4</sub>alkylenearyl, C(=O)C<sub>1-4</sub>alkyleneheteroaryl, C<sub>1-4</sub>alkylenearyl optionally substituted with one or more of halo SO<sub>2</sub>N(R<sup>a</sup>)<sub>2</sub>, N(R<sup>a</sup>)<sub>2</sub>, C(=O)OR<sup>a</sup>, NR<sup>a</sup>SO<sub>2</sub>CF<sub>3</sub>, CN, NO<sub>2</sub>, C(=O)R<sup>a</sup>, OR<sup>a</sup>, C<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, and OC<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, C<sub>1-4</sub>-

10 alkyleneheteroaryl, C<sub>1-4</sub>alkyleneHet, C<sub>1-4</sub>alkyleneC(=O)-C<sub>1-4</sub>alkylenearyl, C<sub>1-4</sub>alkyleneC(=O)C<sub>1-4</sub>alkyleneheteroaryl, C<sub>1-4</sub>alkyleneC(=O)Het, C<sub>1-4</sub>alkyleneC(=O)N(R<sup>a</sup>)<sub>2</sub>, C<sub>1-4</sub>alkyleneOR<sup>a</sup>, C<sub>1-4</sub>alkyleneNR<sup>a</sup>C(=O)R<sup>a</sup>, C<sub>1-4</sub>alkyleneO-C<sub>1-4</sub>alkyleneOR<sup>a</sup>, C<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, C<sub>1-4</sub>alkyleneC(=O)-

15 OR<sup>a</sup>, and C<sub>1-4</sub>alkyleneOC<sub>1-4</sub>alkyleneC(=O)OR<sup>a</sup>;

R<sup>a</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>heterocycloalkyl, C<sub>1-3</sub>alkyleneN(R<sup>c</sup>)<sub>2</sub>, aryl, arylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, heteroaryl, heteroarylC<sub>1-3</sub>alkyl, and

20 C<sub>1-3</sub>alkyleneheteroaryl;

or two R<sup>a</sup> groups are taken together to form a 5- or 6-membered ring, optionally containing at least one heteroatom;

R<sup>b</sup> is selected from the group consisting of

25 hydrogen, C<sub>1-6</sub>alkyl, heteroC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkyleneheteroC<sub>1-3</sub>alkyl, arylheteroC<sub>1-3</sub>alkyl, aryl, heteroaryl, arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, and C<sub>1-3</sub>alkyleneheteroaryl;

R<sup>c</sup> is selected from the group consisting of

30 hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, aryl, and heteroaryl;

Het is a 5- or 6-membered heterocyclic ring, saturated or partially or fully unsaturated,

- 38 -

containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted with  $C_{1-4}$ alkyl or  $C(=O)OR^a$ ; and pharmaceutically acceptable salts and solvates (e.g., hydrates) thereof.

As used herein, the term "alkyl" is defined as straight chained and branched hydrocarbon groups containing the indicated number of carbon atoms, typically methyl, ethyl, and straight chain and branched propyl and butyl groups. The hydrocarbon group can contain up to 16 carbon atoms, preferably one to eight carbon atoms. The term "alkyl" includes "bridged alkyl," i.e., a  $C_6$ - $C_{16}$  bicyclic or polycyclic hydrocarbon group, for example, norbornyl, adamantyl, bicyclo[2.2.2]octyl, bicyclo[2.2.1]heptyl, bicyclo[3.2.1]octyl, or decahydronaphthyl. The term "cycloalkyl" is defined as a cyclic  $C_3$ - $C_8$  hydrocarbon group, e.g., cyclopropyl, cyclobutyl, cyclohexyl, and cyclopentyl.

The term "alkenyl" is defined identically as "alkyl," except for containing a carbon-carbon double bond. "Cycloalkenyl" is defined similarly to cycloalkyl, except a carbon-carbon double bond is present in the ring.

The term "alkylene" is defined as an alkyl group having a substituent. For example, the term " $C_{1-3}$ alkylenearyl" refers to an alkyl group containing one to three carbon atoms, and substituted with an aryl group.

The term "hetero $C_{1-3}$ alkyl" is defined as a  $C_{1-3}$ alkyl group further containing a heteroatom selected from O, S, and  $NR^a$ . For example,  $-CH_2OCH_3$  or  $-CH_2CH_2SCH_3$ . The term "arylhetero $C_{1-3}$ alkyl" refers

- 39 -

to an aryl group having a heteroC<sub>1-3</sub>alkyl substituent.

5 The term "halo" or "halogen" is defined herein to include fluorine, bromine, chlorine, and iodine.

10 The term "haloalkyl" is defined herein as an alkyl group substituted with one or more halo substituents, either fluoro, chloro, bromo, iodo, or combinations thereof. Similarly, "halocycloalkyl" is defined as a cycloalkyl group having one or more halo substituents.

15 The term "aryl," alone or in combination, is defined herein as a monocyclic or polycyclic aromatic group, preferably a monocyclic or bicyclic aromatic group, e.g., phenyl or naphthyl. Unless otherwise indicated, an "aryl" group can be unsubstituted or substituted, for example, with one or more, and in particular one to three, halo, alkyl, phenyl, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkyl, nitro, amino, alkylamino, acylamino, alkylthio, alkylsulfinyl, and alkylsulfonyl. Exemplary aryl groups include phenyl, naphthyl, biphenyl, tetrahydronaphthyl, chlorophenyl, fluorophenyl, aminophenyl, methylphenyl, methoxyphenyl, trifluoromethylphenyl, nitrophenyl, carboxyphenyl, and the like. The terms "arylC<sub>1-3</sub>alkyl" and "heteroarylC<sub>1-3</sub>alkyl" are defined as an aryl or heteroaryl group having a C<sub>1-3</sub>alkyl substituent.

25 30 The term "heteroaryl" is defined herein as a monocyclic or bicyclic ring system containing one or two aromatic rings and containing at least one nitrogen, oxygen, or sulfur atom in an aromatic ring, and which can be unsubstituted or substituted,

- 40 -

for example, with one or more, and in particular one to three, substituents, like halo, alkyl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkyl, nitro, amino, alkylamino, acylamino, alkylthio, alkylsulfanyl, and alkylsulfonyl. Examples of heteroaryl groups include thienyl, furyl, pyridyl, oxazolyl, quinolyl, isoquinolyl, indolyl, triazolyl, isothiazolyl, isoxazolyl, imidazolyl, benzothiazolyl, pyrazinyl, pyrimidinyl, thiazolyl, and thiadiazolyl.

The term "Het" is defined as monocyclic, bicyclic, and tricyclic groups containing one or more heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur. A "Het" group also can contain an oxo group (=O) attached to the ring. Nonlimiting examples of Het groups include 1,3-dioxolane, 2-pyrazoline, pyrazolidine, pyrrolidine, piperazine, a pyrroline, 2H-pyran, 4H-pyran, morpholine, thiopholine, piperidine, 1,4-dithiane, and 1,4-dioxane.

The term "hydroxy" is defined as -OH.

The term "alkoxy" is defined as -OR, wherein R is alkyl.

The term "alkoxyalkyl" is defined as an alkyl group wherein a hydrogen has been replaced by an alkoxy group. The term "(alkylthio)alkyl" is defined similarly as alkoxyalkyl, except a sulfur atom, rather than an oxygen atom, is present.

The term "hydroxyalkyl" is defined as a hydroxy group appended to an alkyl group.

The term "amino" is defined as -NH<sub>2</sub>, and the term "alkylamino" is defined as -NR<sub>2</sub>, wherein at least one R is alkyl and the second R is alkyl or hydrogen.

- 41 -

The term "acylamino" is defined as  $RC(=O)N$ , wherein R is alkyl or aryl.

The term "alkylthio" is defined as -SR, wherein R is alkyl.

5       The term "alkylsulfinyl" is defined as  $R-SO_2$ , wherein R is alkyl.

      The term "amino" is defined as  $-NH_2$ , and the term "alkylamino" is defined as  $-NR_2$ , wherein at least one R is alkyl and the second R is alkyl or  
10       hydrogen.

      The term "acylamino" is defined as  $RC(=O)N$ , wherein R is alkyl or aryl.

      The term "alkylthio" is defined as -SR, wherein R is alkyl.

15       The term "alkylsulfinyl" is defined as  $R-SO_2$ , wherein R is alkyl.

      The term "alkylsulfonyl" is defined as  $R-SO_3$ , wherein R is alkyl.

      The term "nitro" is defined as  $-NO_2$ .

20       The term "trifluoromethyl" is defined as  $-CF_3$ .

      The term "trifluoromethoxy" is defined as  $-OCF_3$ .

      The term "cyano" is defined as  $-CN$ .


25       In preferred embodiments, X is selected from the group consisting of  $CH_2$ ,  $CH_2CH_2$ ,  $CH=CH$ ,  $CH(CH_3)$ ,  $CH(CH_2CH_3)$ ,  $CH_2CH(CH_3)$ , and  $C(CH_3)_2$ . In further preferred embodiments, Y is selected from the group consisting of null, S, and NH.

30       The A ring can be monocyclic or bicyclic. Monocyclic A ring systems are aromatic. Bicyclic A ring systems contain at least one aromatic ring, but both rings can be aromatic. Examples of A ring


- 42 -

systems include, but are not limited to, imidazolyl,  
pyrazolyl, 1,2,3-triazolyl, pyridizynyl, pyrimidin-  
yl, pyrazinyl, 1,3,5-triazinyl, purinyl, cinnolinyl,  
phthalazinyl, quinazolinyl, quinoxalinyl, 1,8-  
5 naphthyridinyl, pteridinyl, 1H-indazolyl, and benz-  
imidazolyl.


In a preferred group of compounds of  
formula (I), A is represented by an optionally  
substituted ring system selected from the group  
10 consisting of



15




20




25

- 43 -




5



10

, and


15



20

The A ring system optionally can be substituted with one to three, and preferably one to two, substituents selected from the group consisting of  $N(R^a)_2$ , halo,  $C_{1-3}$ alkyl,  $S(C_{1-3}$ alkyl),  $OR^a$ , and

25




30




- 44 -

Specific substituents include, but are not limited to,  $\text{NH}_2$ ,  $\text{NH}(\text{CH}_3)$ ,  $\text{N}(\text{CH}_3)_2$ ,  $\text{NHCH}_2\text{C}_6\text{H}_5$ ,  $\text{NH}(\text{C}_2\text{H}_5)$ ,  $\text{Cl}$ ,  $\text{F}$ ,  $\text{CH}_3$ ,  $\text{SCH}_3$ ,  $\text{OH}$ , and



In another preferred group of compounds of formula (I),  $\text{R}^1$  and  $\text{R}^2$ , independently, are represented by hydrogen,  $\text{OR}^a$ , halo,  $\text{C}_{1-6}\text{alkyl}$ ,  $\text{CF}_3$ ,  $\text{NO}_2$ ,  $\text{N}(\text{R}^a)_2$ ,  $\text{NR}^a\text{C}_{1-3}\text{alkyleneN}(\text{R}^a)_2$ , and  $\text{OC}_{1-3}\text{alkyleneOR}^a$ . Specific substituents include, but are not limited to,  $\text{H}$ ,  $\text{OCH}_3$ ,  $\text{Cl}$ ,  $\text{Br}$ ,  $\text{F}$ ,  $\text{CH}_3$ ,  $\text{CF}_3$ ,  $\text{NO}_2$ ,  $\text{OH}$ ,  $\text{N}(\text{CH}_3)_2$ ,



and  $\text{O}(\text{CH}_2)_2\text{OCH}_2\text{C}_6\text{H}_5$ .  $\text{R}^1$  and  $\text{R}^2$  also can be taken together to form a ring, for example, a phenyl ring.


In a preferred embodiment,  $\text{R}^3$  is selected from the group consisting of optionally substituted  $\text{C}_{1-6}\text{alkyl}$ , aryl, heteroaryl,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{3-8}\text{heterocycloalkyl}$ ,  $\text{C}(=\text{O})\text{OR}^a$ ,  $\text{C}_{1-4}\text{alkyleneHet}$ ,  $\text{C}_{1-4}\text{alkylene-cycloalkyl}$ ,  $\text{C}_{1-4}\text{alkylenearyl}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{C}_{1-4}\text{-alkylenearyl}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^a$ ,  $\text{C}_{1-4}\text{alkylene-}$

30


- 45 -

$C(=O)N(R^a)_2$ ,  $C_{1-4}alkyleneC(=O)Het$ ,  $C_{1-4}alkyleneN(R^a)_2$ ,  
and  $C_{1-4}alkyleneNR^aC(=O)R^a$ . Specific  $R^3$  groups  
include, but are not limited to


5




10



15




20




25


30




- 46 -



5



10




15



20

, and



25


The R<sup>3</sup> group can be substituted with one to three substituents, for example, halo, OR<sup>a</sup>, C<sub>1-6</sub>alkyl, aryl, heteroaryl, NO<sub>2</sub>, N(R<sup>a</sup>)<sub>2</sub>, NR<sup>a</sup>SO<sub>2</sub>CF<sub>3</sub>, NR<sup>a</sup>C(=O)R<sup>a</sup>, C(=O)OR<sup>a</sup>, N(R<sup>a</sup>)C<sub>1-4</sub>alkylene(R<sup>a</sup>)<sub>2</sub>, SO<sub>2</sub>N(R<sup>a</sup>)<sub>2</sub>, CN, C(=O)R<sup>a</sup>, C<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, OC<sub>1-4</sub>alkyleneC≡CR<sup>a</sup>, OC<sub>1-4</sub>alkylene-C(=O)N(R<sup>a</sup>)<sub>2</sub>, OC<sub>1-4</sub>alkylenearyl, OC<sub>1-4</sub>alkylenehetero-

30

- 47 -

aryl,  $\text{OC}_{1-4}\text{alkyleneHet}$ ,  $\text{OC}_{1-4}\text{alkyleneN(R}^a)_2$ , and  $\text{N(R}^a)_2\text{-C}_{1-4}\text{alkyleneN(R}^a)_2$ . Specific substituents for the  $\text{R}^3$  group include, but are not limited to, Cl, F,  $\text{CH}_3$ ,  $\text{CH(CH}_3)_2$ , OH,  $\text{OCH}_3$ ,  $\text{O(CH}_2)_3\text{N(CH}_3)_2$ ,  $\text{OCH}_2\text{C}\equiv\text{CH}$ ,  $\text{OCH}_2\text{C(=O)-NH}_2$ ,  $\text{C}_6\text{H}_5$ ,  $\text{NO}_2$ ,  $\text{NH}_2$ ,  $\text{NHC(=O)CH}_3$ ,  $\text{CO}_2\text{H}$ ,  $\text{N(CH}_3)_2\text{CH}_2\text{CH}_2\text{N(CH}_3)_2$ , and


10



15

As used herein, the quinazoline ring structure, and numbering of the ring structure, is


20



25

The purine ring structure, and numbering of the ring structure, is

- 48 -



5

The compounds provided by the invention

are exemplified as follows:

3-(2-isopropylphenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

5-chloro-2-(9H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one;

5-chloro-3-(2-fluorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-(2-fluorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-(2-methoxyphenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-(2,6-dichlorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-(2-chlorophenyl)-6-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

5-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-(2-chlorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-(2-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

3-(2-chlorophenyl)-5-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

- 49 -

- 3-benzyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 3-butyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 5 3-(2-chlorophenyl)-7-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 3-morpholin-4-yl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one, acetate salt;
- 8-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 10 3-(2-chlorophenyl)-6,7-difluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 3-(3-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 15 6-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 3-(3-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 2-(9H-purin-6-ylsulfanylmethyl)-3-pyridin-4-yl-3H-quinazolin-4-one;
- 20 3-(2-chlorophenyl)-8-trifluoromethyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 3-benzyl-5-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- 25 3-(4-methylpiperazin-1-yl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one, acetate salt;
- 3-(2-chlorophenyl)-6-hydroxy-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;
- [5-fluoro-4-oxo-2-(9H-purin-6-ylsulfanylmethyl)-4H-quinazolin-3-yl]acetic acid ethyl ester;
- 30 3-(2-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;

- 50 -

- 3-biphenyl-2-yl-5-chloro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
5-chloro-3-(2-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
5 2-(6-aminopurin-9-ylmethyl)-3-(2-isopropylphenyl)-5-methyl-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-3-biphenyl-2-yl-5-chloro-3H-quinazolin-4-one;  
10 2-(6-aminopurin-9-ylmethyl)-3-(2-fluorophenyl)-5-methyl-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-fluorophenyl)-3H-quinazolin-4-one;  
15 2-(6-aminopurin-9-ylmethyl)-8-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-5-methyl-3H-quinazolin-4-one;  
20 2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-5-fluoro-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-3-benzyl-5-fluoro-3H-quinazolin-4-one;  
25 2-(6-aminopurin-9-ylmethyl)-3-butyl-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-3-morpholin-4-yl-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-7-fluoro-3H-quinazolin-4-one;  
30 2-(6-aminopurin-9-ylmethyl)-6-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one;

- 51 -

3-(4-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-quinazolin-4-one;  
3-(2-chlorophenyl)-6,7-dimethoxy-2-(9H-purin-6-  
ylsulfanylmethyl)-3H-quinazoline-4-one;  
5 3-(2-chlorophenyl)-7-nitro-2-(9H-purin-6-ylsulfan-  
ylmethyl)-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-6-bromo-3-(2-chlorophen-  
yl)-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-6,7-  
10 dimethoxy-3H-quinazolin-4-one;  
6-bromo-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfan-  
ylmethyl)-3H-quinazolin-4-one;  
3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-benzo[g]quinazolin-4-one;  
15 2-(6-aminopurin-9-ylmethyl)-5-chloro-3-o-tolyl-3H-  
quinazolin-4-one; and  
2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-methoxy-  
phenyl)-3H-quinazolin-4-one.

20 The preferred compounds provided by the  
invention have the structure (IV), exemplified as  
follows:  
3-(2-isopropylphenyl)-5-methyl-2-(9H-purin-6-ylsul-  
fanylmethyl)-3H-quinazolin-4-one;  
25 5-chloro-2-(9H-purin-6-ylsulfanylmethyl)-3-o-tolyl-  
3H-quinazolin-4-one;  
5-chloro-3-(2-fluorophenyl)-2-(9H-purin-6-ylsulfan-  
ylmethyl)-3H-quinazolin-4-one;  
3-(2-fluorophenyl)-5-methyl-2-(9H-purin-6-ylsulfan-  
ylmethyl)-3H-quinazolin-4-one;  
30 3-(2,6-dichlorophenyl)-5-methyl-2-(9H-purin-6-ylsul-  
fanylmethyl)-3H-quinazolin-4-one;



- 52 -

- 3-(2-chlorophenyl)-6-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
5-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
5 3-(2-chlorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-(2-chlorophenyl)-5-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-benzyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
10 3-butyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-(2-chlorophenyl)-7-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
15 3-morpholin-4-yl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one, acetate salt;  
8-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-(2-chlorophenyl)-6,7-difluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
20 6-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
3-(3-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
25 2-(9H-purin-6-ylsulfanylmethyl)-3-pyridin-4-yl-3H-quinazolin-4-one;  
3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-trifluoromethyl-3H-quinazolin-4-one;  
3-benzyl-5-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one;  
30 3-(4-methylpiperazin-1-yl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one, acetate salt;

- 53 -

- 3- (2-chlorophenyl) -6-hydroxy-2- (9H-purin-6-ylsulfanylmethyl) -3H-quinazolin-4-one;  
[5-fluoro-4-oxo-2- (9H-purin-6-ylsulfanylmethyl) -4H-quinazolin-3-yl]acetic acid ethyl ester;
- 5 3-biphenyl-2-yl-5-chloro-2- (9H-purin-6-ylsulfanylmethyl) -3H-quinazolin-4-one;  
2- (6-aminopurin-9-ylmethyl) -3- (2-isopropylphenyl) -5-methyl-3H-quinazolin-4-one;  
2- (6-aminopurin-9-ylmethyl) -5-methyl-3-o-tolyl-3H-
- 10 quinazolin-4-one;  
2- (6-aminopurin-9-ylmethyl) -3-biphenyl-2-yl-5-chloro-3H-quinazolin-4-one;  
2- (6-aminopurin-9-ylmethyl) -3- (2-fluorophenyl) -5-methyl-3H-quinazolin-4-one;
- 15 2- (6-aminopurin-9-ylmethyl) -5-chloro-3- (2-fluorophenyl) -3H-quinazolin-4-one;  
2- (6-aminopurin-9-ylmethyl) -8-chloro-3- (2-chlorophenyl) -3H-quinazolin-4-one;  
2- (6-aminopurin-9-ylmethyl) -5-chloro-3- (2-chlorophenyl) -3H-quinazolin-4-one;
- 20 2- (6-aminopurin-9-ylmethyl) -3- (2-chlorophenyl) -5-methyl-3H-quinazolin-4-one;  
2- (6-aminopurin-9-ylmethyl) -3- (2-chlorophenyl) -5-fluoro-3H-quinazolin-4-one;
- 25 2- (6-aminopurin-9-ylmethyl) -3-benzyl-5-fluoro-3H-quinazolin-4-one;  
2- (6-aminopurin-9-ylmethyl) -3-butyl-3H-quinazolin-4-one;  
2- (6-aminopurin-9-ylmethyl) -3-morpholin-4-yl-3H-
- 30 quinazolin-4-one;  
2- (6-aminopurin-9-ylmethyl) -3- (2-chlorophenyl) -7-fluoro-3H-quinazolin-4-one; and

- 54 -

2-(6-aminopurin-9-ylmethyl)-5-chloro-3-o-tolyl-3H-quinazoline-4-one.

It is generally accepted that biological systems can exhibit very sensitive activities with respect to the absolute stereochemical nature of compounds. (See, E.J. Ariens, *Medicinal Research Reviews*, 6:451-466 (1986); E.J. Ariens, *Medicinal Research Reviews*, 7:367-387 (1987); K.W. Fowler, *Handbook of Stereoisomers: Therapeutic Drugs*, CRC Press, edited by Donald P. Smith, pp. 35-63 (1989); and S.C. Stinson, *Chemical and Engineering News*, 75:38-70 (1997.)

Therefore, the present invention includes all possible stereoisomers and geometric isomers of compounds of structural formulae (I)-(IV), and includes not only racemic compounds, but also the optically active isomers as well. When a compound of structural formulae (I)-(IV) is desired as a single enantiomer, it can be obtained either by resolution of the final product or by stereospecific synthesis from either isomerically pure starting material or use of a chiral auxiliary reagent, for example, see Z. Ma et al., *Tetrahedron: Asymmetry*, 8(6), pages 883-888 (1997). Resolution of the final product, an intermediate, or a starting material can be achieved by any suitable method known in the art. Additionally, in situations where tautomers of the compounds of structural formulae (I)-(IV) are possible, the present invention is intended to include all tautomeric forms of the compounds. Specific stereoisomers exhibit an excellent ability to inhibit kinase activity of PI3K $\delta$ .

- 55 -

The term "prodrug" as used herein refers to compounds that are rapidly transformed in vivo to a compound having structural formula (I) herein-above, for example, by hydrolysis. Prodrug design is discussed generally in Hardma et al. (Eds.),  
5 *Goodman and Gilman's The Pharmacological Basis of Therapeutics*, 9th ed., pp. 11-16 (1996). A thorough discussion is provided in Higuchi et al., *Prodrugs as Novel Delivery Systems*, Vol. 14, ASCD Symposium Series, and in Roche (ed.), *Bioreversible Carriers in Drug Design*, American Pharmaceutical Association and Pergamon Press (1987). Briefly, administration of a drug is followed by elimination from the body or some biotransformation whereby biological  
10 activity of the drug is reduced or eliminated. Alternatively, a biotransformation process can lead to a metabolic by-product, which is itself more active or equally active as compared to the drug initially administered. Increased understanding of these biotransformation processes permits the design of so-called "prodrugs," which, following a bio-  
15 transformation, become more physiologically active in their altered state. Prodrugs, therefore, encompass pharmacologically inactive compounds that are converted to biologically active metabolites.  
20

To illustrate, prodrugs can be converted into a pharmacologically active form through hydrolysis of, for example, an ester or amide linkage, thereby introducing or exposing a functional group  
25 on the resultant product. The prodrugs can be designed to react with an endogenous compound to form a water-soluble conjugate that further enhances the pharmacological properties of the compound, for  
30

- 56 -

example, increased circulatory half-life. Alternatively, prodrugs can be designed to undergo covalent modification on a functional group with, for example, glucuronic acid, sulfate, glutathione, amino acids, or acetate. The resulting conjugate can be inactivated and excreted in the urine, or rendered more potent than the parent compound. High molecular weight conjugates also can be excreted into the bile, subjected to enzymatic cleavage, and released back into the circulation, thereby effectively increasing the biological half-life of the originally administered compound.

#### 15 Methods for Identifying Negative Regulators of PI3K $\delta$ Activity

The PI3K $\delta$  protein, as well as fragments thereof possessing biological activity, can be used for screening putative negative regulator compounds in any of a variety of drug screening techniques. A negative regulator of PI3K $\delta$  is a compound that diminishes or abolishes the ability of PI3K $\delta$  to carry out any of its biological functions. An example of such compounds is an agent that decreases the ability of a PI3K $\delta$  polypeptide to phosphorylate phosphatidylinositol or to target appropriate structures within a cell. The selectivity of a compound that negatively regulates PI3K $\delta$  activity can be evaluated by comparing its activity on the PI3K $\delta$  to its activity on other proteins. Selective negative regulators include, for example, antibodies and other proteins or peptides that specifically bind to a PI3K $\delta$  polypeptide, oligonucleotides that specifi-

- 57 -

cally bind to PI3K $\delta$  polypeptides, and other nonpeptide compounds (e.g., isolated or synthetic organic molecules) that specifically interact with PI3K $\delta$  polypeptides. Negative regulators also include compounds as described above, but which interact with a specific binding partner of PI3K $\delta$  polypeptides.

Presently preferred targets for the development of selective negative regulators of PI3K $\delta$  include, for example:

(1) cytoplasmic regions of PI3K $\delta$  polypeptides that contact other proteins and/or localize PI3K $\delta$  within a cell;

(2) regions of PI3K $\delta$  polypeptides that bind specific binding partners;

(3) regions of the PI3K $\delta$  polypeptides that bind substrate;

(4) allosteric regulatory sites of the PI3K $\delta$  polypeptides that can or cannot interact directly with the active site upon regulatory signal;

(5) regions of the PI3K $\delta$  polypeptides that mediate multimerization.

For example, one target for development of modulators is the identified regulatory interaction of p85 with p110 $\delta$ , which can be involved in activation and/or subcellular localization of the p110 $\delta$  moiety. Still other selective modulators include those that recognize specific regulatory or PI3K $\delta$ -encoding nucleotide sequences. Modulators of PI3K $\delta$  activity can be therapeutically useful in treatment of a wide range of diseases and physiological conditions in which aberrant PI3K $\delta$  activity is involved.

- 58 -

Accordingly, the invention provides methods of characterizing the potency of a test compound as an inhibitor of PI3K $\delta$  polypeptide, said method comprising the steps of (a) measuring  
5 activity of a PI3K $\delta$  polypeptide in the presence of a test compound; (b) comparing the activity of the PI3K $\delta$  polypeptide in the presence of the test compound to the activity of the PI3K $\delta$  polypeptide in the presence of an equivalent amount of a reference  
10 compound (e.g., a PI3K $\delta$  inhibitor compound of the invention as described herein), wherein a lower activity of the PI3K $\delta$  polypeptide in the presence of the test compound than in the presence of the reference indicates that the test compound is a more  
15 potent inhibitor than the reference compound, and a higher activity of the PI3K $\delta$  polypeptide in the presence of the test compound than in the presence of the reference indicates that the test compound is a less potent inhibitor than the reference compound.

20 The invention further provides methods of characterizing the potency of a test compound as an inhibitor of PI3K $\delta$  polypeptide, comprising the steps of (a) determining an amount of a control compound (e.g., a PI3K $\delta$  inhibitor compound of the in-  
25 vention as described herein) that inhibits an activity of a PI3K $\delta$  polypeptide by a reference percentage of inhibition, thereby defining a reference inhibitory amount for the control compound; (b) determining an amount of a test compound that inhibits an  
30 activity of a PI3K $\delta$  polypeptide by a reference percentage of inhibition, thereby defining a reference inhibitory amount for the test compound; (c) comparing the reference inhibitory amount for the test

- 59 -

compound to the reference inhibitory amount for the control compound, wherein a lower reference inhibitory amount for the test compound than for the control compound indicates that the test compound is a more potent inhibitor than the control compound, and a higher reference inhibitory amount for the test compound than for the control compound indicates that the test compound is a less potent inhibitor than the control compound. In one aspect, the method uses a reference inhibitory amount which is the amount of the compound that inhibits the activity of the PI3K $\delta$  polypeptide by 50%, 60%, 70%, or 80%. In another aspect the method employs a reference inhibitory amount that is the amount of the compound that inhibits the activity of the PI3K $\delta$  polypeptide by 90%, 95%, or 99%. These methods comprise determining the reference inhibitory amount of the compounds in an *in vitro* biochemical assay, in an *in vitro* cell-based assay, or in an *in vivo* assay.

The invention further provides methods of identifying a negative regulator of PI3K $\delta$  activity, comprising the steps of (i) measuring activity of a PI3K $\delta$  polypeptide in the presence and absence of a test compound, and (ii) identifying as a negative regulator a test compound that decreases PI3K $\delta$  activity and that competes with a compound of the invention for binding to PI3K $\delta$ . Furthermore, the invention provides methods for identifying compounds that inhibit PI3K $\delta$  activity, comprising the steps of (i) contacting a PI3K $\delta$  polypeptide with a compound of the invention in the presence and absence of a test compound, and (ii) identifying a test compound



- 60 -

as a negative regulator of PI3K $\delta$  activity wherein the compound competes with a compound of the invention for binding to PI3K $\delta$ . The invention therefore provides a method for screening for candidate negative regulators of PI3K $\delta$  activity and/or to confirm the mode of action of candidate such negative regulators. Such methods can be employed against other PI3K isoforms in parallel to establish comparative activity of the test compound across the isoforms and/or relative to a compound of the invention.

In these methods, the PI3K $\delta$  polypeptide can be a fragment of p110 $\delta$  that exhibits kinase activity, i.e., a fragment comprising the catalytic site of p110 $\delta$ . Alternatively, the PI3K $\delta$  polypeptide can be a fragment from the p110 $\delta$ -binding domain of p85 and provides a method to identify allosteric modulators of PI3K $\delta$ . The methods can be employed in cells expressing cells expressing PI3K $\delta$  or its subunits, either endogenously or exogenously. Accordingly, the polypeptide employed in such methods can be free in solution, affixed to a solid support, modified to be displayed on a cell surface, or located intracellularly. The modulation of activity or the formation of binding complexes between the PI3K $\delta$  polypeptide and the agent being tested then can be measured.

Human PI3K polypeptides are amenable to biochemical or cell-based high throughput screening (HTS) assays according to methods known and practiced in the art, including melanophore assay systems to investigate receptor-ligand interactions, yeast-based assay systems, and mammalian cell expression systems. For a review, see Jayawickreme

- 61 -

and Kost, *Curr Opin Biotechnol*, 8:629-34 (1997).  
Automated and miniaturized HTS assays also are  
comprehended as described, for example, in Houston  
and Banks, *Curr Opin Biotechnol*, 8:734-40 (1997).

5               Such HTS assays are used to screen  
libraries of compounds to identify particular com-  
pounds that exhibit a desired property. Any library  
of compounds can be used, including chemical librar-  
ies, natural product libraries, and combinatorial  
10       libraries comprising random or designed oligopep-  
tides, oligonucleotides, or other organic compounds.

              Chemical libraries can contain known com-  
pounds, proprietary structural analogs of known  
compounds, or compounds that are identified from  
15       natural product screening.

              Natural product libraries are collections  
of materials isolated from natural sources, typi-  
cally, microorganisms, animals, plants, or marine  
organisms. Natural products are isolated from their  
20       sources by fermentation of microorganisms followed  
by isolation and extraction of the fermentation  
broths or by direct extraction from the microorgan-  
isms or tissues (plants or animal) themselves. Nat-  
ural product libraries include polyketides, nonribo-  
25       somal peptides, and variants (including nonnaturally  
occurring variants) thereof. For a review, see Cane  
et al., *Science*, 282:63-68 (1998).

              Combinatorial libraries are composed of  
large numbers of related compounds, such as pep-  
30       tides, oligonucleotides, or other organic compounds  
as a mixture. Such compounds are relatively  
straightforward to design and prepare by traditional  
automated synthesis protocols, PCR, cloning, or pro-

- 62 -

prietary synthetic methods. Of particular interest are peptide and oligonucleotide combinatorial libraries.

Still other libraries of interest include peptide, protein, peptidomimetic, multiparallel synthetic collection, recombinatorial, and polypeptide libraries. For a review of combinatorial chemistry and libraries created thereby, see Myers, *Curr Opin Biotechnol*, 8:701-07 (1997).

Once compounds have been identified that show activity as negative regulators of PI3K $\delta$  function, a program of optimization can be undertaken in an effort to improve the potency and or selectivity of the activity. This analysis of structure-activity relationships (SAR) typically involves of iterative series of selective modifications of compound structures and their correlation to biochemical or biological activity. Families of related compounds can be designed that all exhibit the desired activity, with certain members of the family, namely those possessing suitable pharmacological profiles, potentially qualifying as therapeutic candidates.

#### Therapeutic Uses of Inhibitors of PI3K $\delta$ Activity

The invention provides a method for selectively or specifically inhibiting PI3K $\delta$  activity therapeutically or prophylactically. The method comprises administering a selective or specific inhibitor of PI3K $\delta$  activity in an amount effective therefor. This method can be employed in treating

- 63 -

humans or animals who are or can be subject to any condition whose symptoms or pathology is mediated by PI3K $\delta$  expression or activity.

"Treating" as used herein refers to preventing a disorder from occurring in an animal that can be predisposed to the disorder, but has not yet been diagnosed as having it; inhibiting the disorder, i.e., arresting its development; relieving the disorder, i.e., causing its regression; or ameliorating the disorder, i.e., reducing the severity of symptoms associated with the disorder. "Disorder" is intended to encompass medical disorders, diseases, conditions, syndromes, and the like, without limitation.

The methods of the invention embrace various modes of treating an animal subject, preferably a mammal, more preferably a primate, and still more preferably a human. Among the mammalian animals that can be treated are, for example, companion animals (pets), including dogs and cats; farm animals, including cattle, horses, sheep, pigs, and goats; laboratory animals, including rats, mice, rabbits, guinea pigs, and nonhuman primates, and zoo specimens. Nonmammalian animals include, for example, birds, fish, reptiles, and amphibians.

In one aspect, the method of the invention can be employed to treat subjects therapeutically or prophylactically who have or can be subject to an inflammatory disorder. One aspect of the present invention derives from the involvement of PI3K $\delta$  in mediating aspects of the inflammatory process. Without intending to be bound by any theory, it is theorized that, because inflammation involves

- 64 -

processes are typically mediated by leukocyte (e.g., neutrophil, lymphocyte, etc.) activation and chemotactic transmigration, and because PI3K $\delta$  can mediate such phenomena, antagonists of PI3K $\delta$  can be used to suppress injury associated with inflammation.

"Inflammatory disorder" as used herein can refer to any disease, disorder, or syndrome in which an excessive or unregulated inflammatory response leads to excessive inflammatory symptoms, host tissue damage, or loss of tissue function. "Inflammatory disorder" also refers to a pathological state mediated by influx of leukocytes and/or neutrophil chemotaxis.

"Inflammation" as used herein refers to a localized, protective response elicited by injury or destruction of tissues, which serves to destroy, dilute, or wall off (sequester) both the injurious agent and the injured tissue. Inflammation is notably associated with influx of leukocytes and/or neutrophil chemotaxis. Inflammation can result from infection with pathogenic organisms and viruses and from noninfectious means such as trauma or reperfusion following myocardial infarction or stroke, immune response to foreign antigen, and autoimmune responses. Accordingly, inflammatory disorders amenable to the invention encompass disorders associated with reactions of the specific defense system as well as with reactions of the nonspecific defense system.

As used herein, the term "specific defense system" refers to the component of the immune system that reacts to the presence of specific antigens. Examples of inflammation resulting from a response

- 65 -

of the specific defense system include the classical response to foreign antigens, autoimmune diseases, and delayed type hypersensitivity response mediated by T-cells. Chronic inflammatory diseases, the rejection of solid transplanted tissue and organs, e.g., kidney and bone marrow transplants, and graft versus host disease (GVHD), are further examples of inflammatory reactions of the specific defense system.

The term "nonspecific defense system" as used herein refers to inflammatory disorders that are mediated by leukocytes that are incapable of immunological memory (e.g., granulocytes, and macrophages). Examples of inflammation that result, at least in part, from a reaction of the nonspecific defense system include inflammation associated with conditions such as adult (acute) respiratory distress syndrome (ARDS) or multiple organ injury syndromes; reperfusion injury; acute glomerulonephritis; reactive arthritis; dermatoses with acute inflammatory components; acute purulent meningitis or other central nervous system inflammatory disorders such as stroke; thermal injury; inflammatory bowel disease; granulocyte transfusion associated syndromes; and cytokine-induced toxicity.

"Autoimmune disease" as used herein refers to any group of disorders in which tissue injury is associated with humoral or cell-mediated responses to the body's own constituents. "Allergic disease" as used herein refers to any symptoms, tissue damage, or loss of tissue function resulting from allergy. "Arthritic disease" as used herein refers to any disease that is characterized by inflammatory

- 66 -

lesions of the joints attributable to a variety of etiologies. "Dermatitis" as used herein refers to any of a large family of diseases of the skin that are characterized by inflammation of the skin

5     attributable to a variety of etiologies. "Trans-plant rejection" as used herein refers to any immune reaction directed against grafted tissue, such as organs or cells (e.g., bone marrow), characterized by a loss of function of the grafted and surrounding

10    tissues, pain, swelling, leukocytosis, and thrombocytopenia.

          The therapeutic methods of the present invention include methods for the treatment of disorders associated with inflammatory cell activation.

15    "Inflammatory cell activation" refers to the induction by a stimulus (including, but not limited to, cytokines, antigens or auto-antibodies) of a proliferative cellular response, the production of soluble mediators (including but not limited to

20    cytokines, oxygen radicals, enzymes, prostanoids, or vasoactive amines), or cell surface expression of new or increased numbers of mediators (including, but not limited to, major histocompatibility anti-

25    gens or cell adhesion molecules) in inflammatory cells (including but not limited to monocytes, macrophages, T lymphocytes, B lymphocytes, granulocytes (i.e., polymorphonuclear leukocytes such as neutrophils, basophils, and eosinophils), mast

30    cells, dendritic cells, Langerhans cells, and endothelial cells). It will be appreciated by persons skilled in the art that the activation of one or a combination of these phenotypes in these cells can

- 67 -

contribute to the initiation, perpetuation, or exacerbation of an inflammatory disorder.

The compounds of the invention have been found to inhibit superoxide release by neutrophils. 5  
Superoxide is released by neutrophils in response to any of a variety of stimuli, including signals of infection, as a mechanism of cell killing. For example, superoxide release is known to be induced by tumor necrosis factor alpha (TNF $\alpha$ ), which is 10  
released by macrophages, mast cells, and lymphocytes upon contact with bacterial cell wall components such as lipopolysaccharide (LPS). TNF $\alpha$  is an extraordinarily potent and promiscuous activator of inflammatory processes, being involved in activation 15  
of neutrophils and various other cell types, induction of leukocyte/endothelial cell adhesion, pyrexia, enhanced MHC class I production, and stimulation of angiogenesis. Alternatively, superoxide release can be stimulated by formyl-Met-Leu-Phe 20  
(fMLP) or other peptides blocked at the N-terminus by formylated methionine. Such peptides are not normally found in eukaryotes, but are fundamentally characteristic of bacteria, and signal the presence of bacteria to the immune system. Leukocytes 25  
expressing the fMLP receptor, e.g., neutrophils and macrophages, are stimulated to migrate up gradients of these peptides (i.e., chemotaxis) toward loci of infection. As demonstrated herein, the compounds of the invention inhibit stimulated superoxide release 30  
by neutrophils in response to either TNF $\alpha$  or fMLP. Other functions of neutrophils, including stimulated exocytosis and directed chemotactic migration, also have been shown to be inhibited by the PI3K $\delta$  inhibi-



- 68 -

tors of the invention. Accordingly, the compounds of the invention can be expected to be useful in treating disorders, such as inflammatory disorders, that are mediated by any or all of these neutrophil functions.

The present invention enables methods of treating such diseases as arthritic diseases, such as rheumatoid arthritis, monoarticular arthritis, osteoarthritis, gouty arthritis, spondylitis; Behcet disease; sepsis, septic shock, endotoxic shock, gram negative sepsis, gram positive sepsis, and toxic shock syndrome; multiple organ injury syndrome secondary to septicemia, trauma, or hemorrhage; ophthalmic disorders such as allergic conjunctivitis, vernal conjunctivitis, uveitis, and thyroid-associated ophthalmopathy; eosinophilic granuloma; pulmonary or respiratory disorders such as asthma, chronic bronchitis, allergic rhinitis, ARDS, chronic pulmonary inflammatory disease (e.g., chronic obstructive pulmonary disease), silicosis, pulmonary sarcoidosis, pleurisy, alveolitis, vasculitis, emphysema, pneumonia, bronchiectasis, and pulmonary oxygen toxicity; reperfusion injury of the myocardium, brain, or extremities; fibrosis such as cystic fibrosis; keloid formation or scar tissue formation; atherosclerosis; autoimmune diseases, such as systemic lupus erythematosus (SLE), autoimmune thyroiditis, multiple sclerosis, some forms of diabetes, and Reynaud's syndrome; and transplant rejection disorders such as GVHD and allograft rejection; chronic glomerulonephritis; inflammatory bowel diseases such as chronic inflammatory bowel disease (CIBD), Crohn's disease, ulcerative colitis, and

- 69 -

necrotizing enterocolitis; inflammatory dermatoses such as contact dermatitis, atopic dermatitis, psoriasis, or urticaria; fever and myalgias due to infection; central or peripheral nervous system inflammatory disorders such as meningitis, encephalitis, and brain or spinal cord injury due to minor trauma; Sjögren's syndrome; diseases involving leukocyte diapedesis; alcoholic hepatitis; bacterial pneumonia; antigen-antibody complex mediated diseases; hypovolemic shock; Type I diabetes mellitus; acute and delayed hypersensitivity; disease states due to leukocyte dyscrasia and metastasis; thermal injury; granulocyte transfusion-associated syndromes; and cytokine-induced toxicity.

The method can have utility in treating subjects who are or can be subject to reperfusion injury, i.e., injury resulting from situations in which a tissue or organ experiences a period of ischemia followed by reperfusion. The term "ischemia" refers to localized tissue anemia due to obstruction of the inflow of arterial blood. Transient ischemia followed by reperfusion characteristically results in neutrophil activation and transmigration through the endothelium of the blood vessels in the affected area. Accumulation of activated neutrophils in turn results in generation of reactive oxygen metabolites, which damage components of the involved tissue or organ. This phenomenon of "reperfusion injury" is commonly associated with conditions such as vascular stroke (including global and focal ischemia), hemorrhagic shock, myocardial ischemia or infarction, organ transplantation, and cerebral vasospasm. To illustrate, reperfusion

- 70 -

injury occurs at the termination of cardiac bypass procedures or during cardiac arrest when the heart, once prevented from receiving blood, begins to reperfuse. It is expected that inhibition of PI3K $\delta$  activity will result in reduced amounts of reperfusion injury in such situations.

With respect to the nervous system, global ischemia occurs when blood flow to the entire brain ceases for a period. Global ischemia can result from cardiac arrest. Focal ischemia occurs when a portion of the brain is deprived of its normal blood supply. Focal ischemia can result from thromboembolytic occlusion of a cerebral vessel, traumatic head injury, edema, or brain tumor. Even if transient, both global and focal ischemia can cause widespread neuronal damage. Although nerve tissue damage occurs over hours or even days following the onset of ischemia, some permanent nerve tissue damage can develop in the initial minutes following the cessation of blood flow to the brain.

Ischemia also can occur in the heart in myocardial infarction and other cardiovascular disorders in which the coronary arteries have been obstructed as a result of atherosclerosis, thrombus, or spasm. Accordingly, the invention is believed to be useful for treating cardiac tissue damage, particularly damage resulting from cardiac ischemia or caused by reperfusion injury in mammals.

In another aspect, selective inhibitors of PI3K $\delta$  activity, such as the compounds of the invention, can be employed in methods of treating diseases of bone, especially diseases in which osteoclast function is abnormal or undesirable. As shown

- 71 -

in Example 6, below, compounds of the invention inhibit osteoclast function *in vitro*. Accordingly, the use of such compounds and other PI3K $\delta$  selective inhibitors can be of value in treating osteoporosis, Paget's disease, and related bone resorption disorders.

In a further aspect, the invention includes methods of using PI3K $\delta$  inhibitory compounds to inhibit the growth or proliferation of cancer cells of hematopoietic origin, preferably cancer cells of lymphoid origin, and more preferably cancer cells related to or derived from B lymphocytes or B lymphocyte progenitors. Cancers amenable to treatment using the method of the invention include, without limitation, lymphomas, e.g., malignant neoplasms of lymphoid and reticuloendothelial tissues, such as Burkitt's lymphoma, Hodgkins' lymphoma, non-Hodgkins lymphomas, lymphocytic lymphomas and the like; multiple myelomas; as well as leukemias such as lymphocytic leukemias, chronic myeloid (myelogenous) leukemias, and the like. In a preferred embodiment, PI3K $\delta$  inhibitory compounds can be used to inhibit or control the growth or proliferation of chronic myeloid (myelogenous) leukemia cells.

In another aspect, the invention includes a method for suppressing a function of basophils and/or mast cells, and thereby enabling treatment of diseases or disorders characterized by excessive or undesirable basophil and/or mast cell activity.

According to the method, a compound of the invention can be used that selectively inhibits the expression or activity of phosphatidylinositol 3-kinase delta (PI3K $\delta$ ) in the basophils and/or mast cells. Prefer-

- 72 -

ably, the method employs a PI3K $\delta$  inhibitor in an amount sufficient to inhibit stimulated histamine release by the basophils and/or mast cells. Accordingly, the use of such compounds and other PI3K $\delta$  selective inhibitors can be of value in treating diseases characterized by histamine release, i.e., allergic disorders, including disorders such as chronic obstructive pulmonary disease (COPD), asthma, ARDS, emphysema, and related disorders.

#### Pharmaceutical Compositions of Inhibitors of PI3K $\delta$ Activity

A compound of the present invention can be administered as the neat chemical, but it is typically preferable to administer the compound in the form of a pharmaceutical composition or formulation. Accordingly, the present invention also provides pharmaceutical compositions that comprise a chemical or biological compound ("agent") that is active as a modulator of PI3K $\delta$  activity and a biocompatible pharmaceutical carrier, adjuvant, or vehicle. The composition can include the agent as the only active moiety or in combination with other agents, such as oligo- or polynucleotides, oligo- or polypeptides, drugs, or hormones mixed with excipient(s) or other pharmaceutically acceptable carriers. Carriers and other ingredients can be deemed pharmaceutically acceptable insofar as they are compatible with other ingredients of the formulation and not deleterious to the recipient thereof.

Techniques for formulation and administration of pharmaceutical compositions can be found in

- 73 -

Remington's *Pharmaceutical Sciences*, 18th Ed., Mack Publishing Co, Easton, PA, 1990. The pharmaceutical compositions of the present invention can be manufactured using any conventional method, e.g.,  
5 mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, melt-spinning, spray-drying, or lyophilizing processes. However, the optimal pharmaceutical formulation will be determined by one of skill in the art  
10 depending on the route of administration and the desired dosage. Such formulations can influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the administered agent. Depending on the condition being treated,  
15 these pharmaceutical compositions can be formulated and administered systemically or locally.

The pharmaceutical compositions are formulated to contain suitable pharmaceutically acceptable carriers, and can optionally comprise excipients and auxiliaries that facilitate processing of  
20 the active compounds into preparations that can be used pharmaceutically. The administration modality will generally determine the nature of the carrier. For example, formulations for parenteral administration can comprise aqueous solutions of the active  
25 compounds in water-soluble form. Carriers suitable for parenteral administration can be selected from among saline, buffered saline, dextrose, water, and other physiologically compatible solutions. Preferred carriers for parenteral administration are  
30 physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiologically buffered saline. For tissue or cellular adminis-

- 74 -

tration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art. For preparations comprising proteins, the formulation can include stabilizing materials, such as polyols (e.g., sucrose) and/or surfactants (e.g., nonionic surfactants), and the like.

Alternatively, formulations for parenteral use can comprise dispersions or suspensions of the active compounds prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, and synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions can contain substances that increase the viscosity of the suspension, such as sodium carboxymethylcellulose, sorbitol, or dextran. Optionally, the suspension also can contain suitable stabilizers or agents that increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. Aqueous polymers that provide pH-sensitive solubilization and/or sustained release of the active agent also can be used as coatings or matrix structures, e.g., methacrylic polymers, such as the EUDRAGIT® series available from Röhm America Inc. (Piscataway, NJ). Emulsions, e.g., oil-in-water and water-in-oil dispersions, also can be used, optionally stabilized by an emulsifying agent or dispersant (surface active materials; surfactants). Suspensions can contain suspending agents such as ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide,

- 75 -

bentonite, agar-agar, gum tragacanth, and mixtures thereof.

Liposomes containing the active agent also can be employed for parenteral administration.

5 Liposomes generally are derived from phospholipids or other lipid substances. The compositions in liposome form also can contain other ingredients, such as stabilizers, preservatives, excipients, and the like. Preferred lipids include phospholipids and phosphatidyl cholines (lecithins), both natural  
10 and synthetic. Methods of forming liposomes are known in the art. See, e.g., Prescott (Ed.), *Methods in Cell Biology*, Vol. XIV, p. 33, Academic Press, New York (1976).

15 The pharmaceutical compositions comprising the agent in dosages suitable for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art. The preparations formulated for oral administration can  
20 be in the form of tablets, pills, capsules, cachets, dragees, lozenges, liquids, gels, syrups, slurries, elixirs, suspensions, or powders. To illustrate, pharmaceutical preparations for oral use can be obtained by combining the active compounds with a  
25 solid excipient, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries if desired, to obtain tablets or dragee cores. Oral formulations can employ liquid carriers similar in type to those  
30 described for parenteral use, e.g., buffered aqueous solutions, suspensions, and the like.

Preferred oral formulations include tablets, dragees, and gelatin capsules. These prep-



- 76 -

arations can contain one or excipients, which include, without limitation:

- a) diluents, such as sugars, including lactose, dextrose, sucrose, mannitol, or sorbitol;
- 5 b) binders, such as magnesium aluminum silicate, starch from corn, wheat, rice, potato, etc.;
- c) cellulose materials, such as methylcellulose, hydroxypropylmethyl cellulose, and sodium  
10 carboxymethylcellulose, polyvinylpyrrolidone, gums, such as gum arabic and gum tragacanth, and proteins, such as gelatin and collagen;
- d) disintegrating or solubilizing agents such as cross-linked polyvinyl pyrrolidone,  
15 starches, agar, alginic acid or a salt thereof, such as sodium alginate, or effervescent compositions;
- e) lubricants, such as silica, talc, stearic acid or its magnesium or calcium salt, and polyethylene glycol;
- 20 f) flavorants and sweeteners;
- g) colorants or pigments, e.g., to identify the product or to characterize the quantity (dosage) of active compound; and
- h) other ingredients, such as preserva-  
25 tives, stabilizers, swelling agents, emulsifying agents, solution promoters, salts for regulating osmotic pressure, and buffers.

Gelatin capsules include push-fit capsules made of gelatin, as well as soft, sealed capsules  
30 made of gelatin and a coating such as glycerol or sorbitol. Push-fit capsules can contain the active ingredient(s) mixed with fillers, binders, lubricants, and/or stabilizers, etc. In soft capsules,

- 77 -

the active compounds can be dissolved or suspended in suitable fluids, such as fatty oils, liquid paraffin, or liquid polyethylene glycol with or without stabilizers.

5                Dragee cores can be provided with suitable coatings such as concentrated sugar solutions, which also can contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable  
10                organic solvents or solvent mixtures.

                 The pharmaceutical composition can be provided as a salt of the active agent. Salts tend to be more soluble in aqueous or other protonic solvents than the corresponding free acid or base  
15                forms. Pharmaceutically acceptable salts are well known in the art. Compounds that contain acidic moieties can form pharmaceutically acceptable salts with suitable cations. Suitable pharmaceutically acceptable cations include, for example, alkali  
20                metal (e.g., sodium or potassium) and alkaline earth (e.g., calcium or magnesium) cations.

                 Compounds of structural formula (I) that contain basic moieties can form pharmaceutically acceptable acid addition salts with suitable acids.  
25                For example, Berge et al. describe pharmaceutically acceptable salts in detail in *J Pharm Sci*, 66:1 (1977). The salts can be prepared *in situ* during the final isolation and purification of the compounds of the invention or separately by reacting a  
30                free base function with a suitable acid.

                 Representative acid addition salts include, but are not limited to, acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesul-

- 78 -

fonate, bisulfate, butyrate, camphorate, camphorol-sulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethane-sulfonate (isothionate), lactate, maleate, methane-sulfonate or sulfate, nicotinate, 2-naphthalene-sulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, phosphate or hydrogen phosphate, glutamate, bicarbonate, p-toluenesulfonate, and undecanoate. Examples of acids that can be employed to form pharmaceutically acceptable acid addition salts include, without limitation, such inorganic acids as hydrochloric acid, hydrobromic acid, sulfuric acid, and phosphoric acid, and such organic acids as oxalic acid, maleic acid, succinic acid, and citric acid.

In light of the foregoing, any reference to compounds of the present invention appearing herein is intended to include compounds of structural formula (I)-(IV), as well as pharmaceutically acceptable salts and solvates, and prodrugs, thereof.

Basic addition salts can be prepared *in situ* during the final isolation and purification of the compounds of the invention or separately by reacting a carboxylic acid-containing moiety with a suitable base such as the hydroxide, carbonate, or bicarbonate of a pharmaceutically acceptable metal cation, or with ammonia or organic primary, secondary, or tertiary amine. Pharmaceutically acceptable basic addition salts include, but are not limited to, cations based on alkali metals or alkaline earth

- 79 -

metals such as lithium, sodium, potassium, calcium, magnesium, and aluminum salts and the like, and nontoxic quaternary ammonium and amine cations including ammonium, tetramethylammonium, tetraethyl-  
5 ammonium, methylammonium, dimethylammonium, trimethylammonium, ethylammonium, diethylammonium, triethylammonium, and the like. Other representative organic amines useful for the formation of base addition salts include ethylenediamine, ethanol-  
10 amine, diethanolamine, piperidine, piperazine, and the like.

Basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides such as methyl, ethyl, propyl, and butyl chlorides, bromides and iodides; dialkyl sulfates like di-  
15 methyl, diethyl, dibutyl, and diamyl sulfates; long chain alkyl halides such as decyl, lauryl, myristyl, and stearyl chlorides, bromides, and iodides; aryl-alkyl halides such as benzyl and phenethyl bromides; and others. Products having modified solubility or  
20 dispersibility are thereby obtained.

Compositions comprising a compound of the invention formulated in a pharmaceutical acceptable carrier can be prepared, placed in an appropriate  
25 container, and labeled for treatment of an indicated condition. Accordingly, there also is contemplated an article of manufacture, such as a container comprising a dosage form of a compound of the invention and a label containing instructions for use of the  
30 compound. Kits are also contemplated under the invention. For example, the kit can comprise a dosage form of a pharmaceutical composition and a package insert containing instructions for use of

- 80 -

the composition in treatment of a medical condition. In either case, conditions indicated on the label can include treatment of inflammatory disorders, cancer, etc.

5

**Methods of Administration of  
Inhibitors of PI3K $\delta$  Activity**

10                   Pharmaceutical compositions comprising an  
inhibitor of PI3K $\delta$  activity can be administered to  
the subject by any conventional method, including  
parenteral and enteral techniques. Parenteral ad-  
ministration modalities include those in which the  
15                   composition is administered by a route other than  
through the gastrointestinal tract, for example,  
intravenous, intraarterial, intraperitoneal, intra-  
medullary, intramuscular, intraarticular, intra-  
thecal, and intraventricular injections. Enteral  
20                   administration modalities include, for example, oral  
(including buccal and sublingual) and rectal admin-  
istration. Transepithelial administration modali-  
ties include, for example, transmucosal administra-  
tion and transdermal administration. Transmucosal  
25                   administration includes, for example, enteral  
administration as well as nasal, inhalation, and  
deep lung administration; vaginal administration;  
and rectal administration. Transdermal administra-  
tion includes passive or active transdermal or  
30                   transcutaneous modalities, including, for example,  
patches and iontophoresis devices, as well as  
topical application of pastes, salves, or ointments.  
Parenteral administration also can be accomplished  
using a high-pressure technique, e.g., POWDERJECT®.

- 81 -

Surgical techniques include implantation of depot (reservoir) compositions, osmotic pumps, and the like. A preferred route of administration for treatment of inflammation can be local or topical delivery for localized disorders such as arthritis, or systemic delivery for distributed disorders, e.g., intravenous delivery for reperfusion injury or for systemic conditions such as septicemia. For other diseases, including those involving the respiratory tract, e.g., chronic obstructive pulmonary disease, asthma, and emphysema, administration can be accomplished by inhalation or deep lung administration of sprays, aerosols, powders, and the like.

For the treatment of neoplastic diseases, especially leukemias and other distributed cancers, parenteral administration is typically preferred. Formulations of the compounds to optimize them for biodistribution following parenteral administration would be desirable. The PI3K $\delta$  inhibitor compounds can be administered before, during, or after administration of chemotherapy, radiotherapy, and/or surgery.

Moreover, the therapeutic index of the PI3K $\delta$  inhibitor compounds can be enhanced by modifying or derivatizing the compounds for targeted delivery to cancer cells expressing a marker that identifies the cells as such. For example, the compounds can be linked to an antibody that recognizes a marker that is selective or specific for cancer cells, so that the compounds are brought into the vicinity of the cells to exert their effects locally, as previously described (see for example,

- 82 -

Pietersz et al., *Immunol Rev*, 129:57 (1992); Trail et al., *Science*, 261:212 (1993); and Rowlinson-Busza et al., *Curr Opin Oncol*, 4:1142 (1992)). Tumor-directed delivery of these compounds enhances the therapeutic benefit by, inter alia, minimizing potential nonspecific toxicities that can result from radiation treatment or chemotherapy. In another aspect, PI3K $\delta$  inhibitor compounds and radio-isotopes or chemotherapeutic agents can be conjugated to the same anti-tumor antibody.

For the treatment of bone resorption disorders or osteoclast-mediated disorders, the PI3K $\delta$  inhibitors can be delivered by any suitable method. Focal administration can be desirable, such as by intraarticular injection. In some cases, it can be desirable to couple the compounds to a moiety that can target the compounds to bone. For example, a PI3K $\delta$  inhibitor can be coupled to compounds with high affinity for hydroxyapatite, which is a major constituent of bone. This can be accomplished, for example, by adapting a tetracycline-coupling method developed for targeted delivery of estrogen to bone (Orme et al., *Bioorg Med Chem Lett*, 4(11):1375-80 (1994)).

To be effective therapeutically in modulating central nervous system targets, the agents used in the methods of the invention should readily penetrate the blood brain barrier when peripherally administered. Compounds that cannot penetrate the blood brain barrier, however, can still be effectively administered by an intravenous route.

- 83 -

As noted above, the characteristics of the agent itself and the formulation of the agent can influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the administered agent. Such pharmacokinetic and pharmacodynamic information can be collected through preclinical *in vitro* and *in vivo* studies, later confirmed in humans during the course of clinical trials. Thus, for any compound used in the method of the invention, a therapeutically effective dose can be estimated initially from biochemical and/or cell-based assays. Then, dosage can be formulated in animal models to achieve a desirable circulating concentration range that modulates PI3K $\delta$  expression or activity. As human studies are conducted, further information will emerge regarding the appropriate dosage levels and duration of treatment for various diseases and conditions.

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD<sub>50</sub> (the dose lethal to 50% of the population) and the ED<sub>50</sub> (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the "therapeutic index," which typically is expressed as the ratio LD<sub>50</sub>/ED<sub>50</sub>. Compounds that exhibit large therapeutic indices, i.e., the toxic dose is substantially higher than the effective dose, are preferred. The data obtained from such cell culture assays and additional animal studies can be used in formulating a range of dosage for human use. The dosage of such



- 84 -

compounds lies preferably within a range of circulating concentrations that include the  $ED_{50}$  with little or no toxicity.

For the methods of the invention, any  
5 effective administration regimen regulating the timing and sequence of doses can be used. Doses of the agent preferably include pharmaceutical dosage units comprising an effective amount of the agent. As used herein, "effective amount" refers to an  
10 amount sufficient to modulate PI3K $\delta$  expression or activity and/or derive a measurable change in a physiological parameter of the subject through administration of one or more of the pharmaceutical dosage units.

15 Exemplary dosage levels for a human subject are of the order of from about 0.001 milligram of active agent per kilogram body weight (mg/kg) to about 100 mg/kg. Typically, dosage units of the active agent comprise from about 0.01 mg to about  
20 10,000 mg, preferably from about 0.1 mg to about 1,000 mg, depending upon the indication, route of administration, etc. Depending on the route of administration, a suitable dose can be calculated according to body weight, body surface area, or  
25 organ size. The final dosage regimen will be determined by the attending physician in view of good medical practice, considering various factors that modify the action of drugs, e.g., the agent's specific activity, the identity and severity of the  
30 disease state, the responsiveness of the patient, the age, condition, body weight, sex, and diet of the patient, and the severity of any infection. Additional factors that can be taken into account

- 85 -

include time and frequency of administration, drug combinations, reaction sensitivities, and tolerance/response to therapy. Further refinement of the dosage appropriate for treatment involving any of the formulations mentioned herein is done routinely by the skilled practitioner without undue experimentation, especially in light of the dosage information and assays disclosed, as well as the pharmacokinetic data observed in human clinical trials.

Appropriate dosages can be ascertained through use of established assays for determining concentration of the agent in a body fluid or other sample together with dose response data.

The frequency of dosing will depend on the pharmacokinetic parameters of the agent and the route of administration. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Accordingly, the pharmaceutical compositions can be administered in a single dose, multiple discrete doses, continuous infusion, sustained release depots, or combinations thereof, as required to maintain desired minimum level of the agent. Short-acting pharmaceutical compositions (i.e., short half-life) can be administered once a day or more than once a day (e.g., two, three, or four times a day). Long acting pharmaceutical compositions might be administered every 3 to 4 days, every week, or once every two weeks. Pumps, such as subcutaneous, intraperitoneal, or subdural pumps, can be preferred for continuous infusion.

- 86 -

The following Examples are provided to further aid in understanding the invention, and presuppose an understanding of conventional methods well-known to those persons having ordinary skill in the art to which the examples pertain, e.g., the construction of vectors and plasmids, the insertion of genes encoding polypeptides into such vectors and plasmids, or the introduction of vectors and plasmids into host cells. Such methods are described in detail in numerous publications including, for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press (1989), Ausubel et al. (Eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (1994); and Ausubel et al. (Eds.), *Short Protocols in Molecular Biology*, 4th ed., John Wiley & Sons, Inc. (1999). The particular materials and conditions described hereunder are intended to exemplify particular aspects of the invention and should not be construed to limit the reasonable scope thereof.

#### EXAMPLE 1

##### Preparation and Purification of Recombinant PI3K $\alpha$ , $\beta$ , and $\delta$

Recombinant PI3K heterodimeric complexes consisting of a p110 catalytic subunit and a p85 regulatory subunit were overexpressed using the BAC-TO-BAC® HT baculovirus expression system (GIBCO/BRL), and then purified for use in biochemical assays. The four Class I PI 3-kinases were cloned into baculovirus vectors as follows:

- 87 -

p110 $\delta$ : A FLAG<sup>®</sup>-tagged version of human p110 $\delta$  (SEQ ID NO:1) (see Chantry et al., *J Biol Chem*, 272:19236-41 (1997)) was subcloned using standard recombinant DNA techniques into the BamHI-XbaI site of the insect cell expression vector pFastbac HTb (Life Technologies, Gaithersburg, MD), such that the clone was in frame with the His tag of the vector. The FLAG<sup>®</sup> system is described in U.S. Patent Nos. 4,703,004; 4,782,137; 4,851,341; and 5,011,912, and reagents are available from Eastman Kodak Co.

p110 $\alpha$ : Similar to the method used for p110 $\delta$ , described above, a FLAG<sup>®</sup>-tagged version of p110 $\alpha$  (see Volinia et al., *Genomics*, 24(3):427-477 (1994)) was subcloned in BamHI-HindIII sites of pFastbac HTb (Life Technologies) such that the clone was in frame with the His tag of the vector.

p110 $\beta$ : A p110 $\beta$  (see Hu et al., *Mol Cell Biol*, 13:7677-88 (1993)) clone was amplified from the human MARATHON<sup>®</sup> Ready spleen cDNA library (Clontech, Palo Alto, CA) according to the manufacturer's protocol using the following primers:

5' Primer

5'-

GATCGAATTCGGCGCCACCATGGACTACAAGGACGACGATGACAAGTGCTTC  
AGTTTCATAATGCCTCC-3' (SEQ ID NO:3)

3' Primer

5'-GATCGCGGCCGCTTAAGATCTGTAGTCTTTCCGAAGTGTGTG-3'  
(SEQ ID NO:4)

The 5' primer was built to contain a FLAG<sup>®</sup> tag in frame with the p110 $\beta$  sequence. After amplification, the FLAG<sup>®</sup>-p110 $\beta$  sequence was subcloned using standard recombinant techniques into the EcoRI-NotI

- 88 -

sites of pFastbac HTa (Life Technologies), such that the clone was in frame with the His tag of the vector.

5 p110 $\gamma$ : The p110 $\gamma$  cDNA (see Stoyanov et al., *Science*, 269:690-93 (1995)) was amplified from a human Marathon Ready spleen cDNA library (Clontech) according to the manufacturer's protocol using the following primers:

5' Primer

10 5'-AGAATGCGGCCGCATGGAGCTGGAGAACTATAAACAGCCC-3' (SEQ ID NO:5)

3' Primer

5'-CGCGGATCCTTAGGCTGAATGTTTCTCTCCTTGTTG-3' (SEQ ID NO:6)

15 A FLAG<sup>®</sup> tag was subsequently attached to the 5' end of the p110 $\gamma$  sequence and was cloned in the *Bam*H1-*Spe*I sites of pFastbac HTb (Life Technologies) using standard recombinant DNA techniques, with the FLAG<sup>®</sup>-110 $\gamma$  sequence in-frame with the His tag of the  
20 vector.

p85 $\alpha$ : A *Bam*H1-*Eco*R1 fragment of FLAG<sup>®</sup>-tagged p85 cDNA (see Skolnik et al., *Cell*, 65:83-89 (1991)) was subcloned into the *Bam*H1-*Eco*R1 sites of the vector pFastbac dual (Life Technologies).

25 Recombinant baculoviruses containing the above clones were generated using manufacturer's recommended protocol (Life Technologies). Baculoviruses expressing His-tagged p110 $\alpha$ , p110 $\beta$ , or p110 $\delta$  catalytic subunit and p85 subunit were coinfectd  
30 into Sf21 insect cells. To enrich the heterodimeric enzyme complex, an excess amount of baculovirus expressing p85 subunit was infected, and the His-tagged p110 catalytic subunit complexed with p85 was

- 89 -

purified on nickel affinity column. Since p110 $\gamma$  does not associate with p85, Sf21 cells were infected with recombinant baculoviruses expressing His-tagged p110 $\gamma$  only. In an alternate approach, p101  
5 can be cloned into baculovirus, to permit coexpression with its preferred binding partner p110 $\gamma$ .

The 72-hour post-infected Sf21 cells (3 liters) were harvested and homogenized in a hypotonic buffer (20 mM HEPES-KOH, pH 7.8, 5 mM KCl, complete protease inhibitor cocktail (Roche Biochemicals, Indianapolis, IN), using a Dounce homogenizer. The homogenates were centrifuged at 1,000 x g for 15 min. The supernatants were further centrifuged at 10,000 x g for 20 min, followed by ultracentrifugation at 100,000 x g for 60 min. The  
15 soluble fraction was immediately loaded onto 10 mL of HITRAP<sup>®</sup> nickel affinity column (Pharmacia, Piscataway, NJ) equilibrated with 50 mL of Buffer A (50 mM HEPES-KOH, pH 7.8, 0.5 M NaCl, 10 mM imidazole). The column was washed extensively with  
20 Buffer A, and eluted with a linear gradient of 10-500 mM imidazole. Free p85 subunit was removed from the column during the washing step and only the heterodimeric enzyme complex eluted at 250 mM  
25 imidazole. Aliquots of nickel fractions were analyzed by 10% SDS-polyacrylamide gel electrophoresis (SDS-PAGE), stained with SYPRO<sup>®</sup> Red (Molecular Probes, Inc., Eugene, OR), and quantitated with STORM<sup>®</sup> PhosphoImager (Molecular Dynamics, Sunnyvale,  
30 CA). The active fractions were pooled and directly loaded onto a 5 mL Hi-trap heparin column preequilibrated with Buffer B containing 50 mM HEPES-KOH, pH 7.5, 50 mM NaCl, 2 mM dithiothreitol (DTT). The

- 90 -

column was washed with 50 mL of Buffer B and eluted with a linear gradient of 0.05-2 M NaCl. A single peak containing PI3K enzyme complex eluted at 0.8 M NaCl. SDS-polyacrylamide gel analysis showed that the purified PI3K enzyme fractions contained a 1:1 stoichiometric complex of p110 and p85 subunits. The protein profile of the enzyme complex during heparin chromatography corresponded to that of lipid kinase activity. The active fractions were pooled and frozen under liquid nitrogen.

## EXAMPLE 2

### PI3K $\delta$ High Throughput Screen (HTS) and Selectivity Assay

A high throughput screen of a proprietary chemical library was performed to identify candidate inhibitors of PI3K $\delta$  activity. PI3K $\delta$  catalyzes a phosphotransfer from  $\gamma$ -[ $^{32}$ P]ATP to PIP<sub>2</sub>/PS liposomes at the D3' position of the PIP<sub>2</sub> lipid inositol ring. This reaction is MgCl<sub>2</sub> dependent and is quenched in high molarity potassium phosphate buffer pH 8.0 containing 30 mM EDTA. In the screen, this reaction is performed in the presence or absence of library compounds. The reaction products (and all unlabeled products) are transferred to a 96-well, pre-wetted PVDF filter plate, filtered, and washed in high molarity potassium phosphate. Scintillant is added to the dried wells and the incorporated radioactivity is quantitated.

The majority of assay operations were performed using a BIOMEK<sup>®</sup> 1000 robotics workstations

- 91 -

(Beckman) and all plates were read using Wallac liquid scintillation plate counter protocols.

The 3X assay stocks of substrate and enzyme were made and stored in a trough (for robotics assays) or a 96-well, V-bottom, polypropylene plate (for manual assays). Reagents were stable for at least 3 hours at room temperature.

The 3X substrate for the HTS contained 0.6 mM Na<sub>2</sub>ATP, 0.10 mCi/mL  $\gamma$ -[<sup>32</sup>P]ATP (NEN, Pittsburgh, PA), 6  $\mu$ M PIP<sub>2</sub>/PS liposomes (Avanti Polar Lipids, Inc., Atlanta, GA), in 20 mM HEPES, pH 7.4.

The 3X enzyme stock for the HTS contained 1.8 nM PI3K $\delta$ , 150  $\mu$ g/mL horse IgG (used only as a stabilizer), 15 mM MgCl<sub>2</sub>, 3 mM DTT in 20 mM HEPES, pH 7.4.

The chemical high throughput screen (HTS) library samples (each containing a pool of 22 compounds) in dimethyl sulfoxide (DMSO) were diluted to 18.75  $\mu$ M or 37.8  $\mu$ M in double distilled water, and 20  $\mu$ L of the dilutions were placed in the wells of a 96-well polypropylene plate for assaying. The negative inhibitor control (or positive enzyme control) was DMSO diluted in water, and the positive inhibitor controls employed concentrations of LY294002 sufficient to provide 50% and 100% inhibition.

To the 20  $\mu$ L pooled chemical library dilutions, 20  $\mu$ L of 3X substrate was added. The reaction was initiated with 20  $\mu$ L of 3X enzyme, incubated at room temperature for 10 minutes. This dilution established a final concentration of 200  $\mu$ M ATP in the reaction volume. The reaction was stopped with 150  $\mu$ L quench buffer (1.0 M potassium phosphate pH 8.0, 30 mM EDTA). A portion of the



- 92 -

quenched solution (180  $\mu$ L) was then transferred to a PVDF filter plate (Millipore #MAIP NOB prewetted with sequential 200  $\mu$ L washes of 100% methanol, water, and finally 1.0 M potassium phosphate pH 8.0 wash buffer).


The PVDF filter plate was aspirated under moderate vacuum (2-5 mm Hg), washed with 5 x 200  $\mu$ L of wash buffer, and then dried by aspiration. The filter was subsequently blotted, allowed to air dry completely, and inserted into a Wallac counting cassette with 50  $\mu$ L of Ecoscint scintillation cocktail added per well. The incorporated radioactivity was quantitated, and data were analyzed, after normalizing to the enzyme positive control (set at 100%), to identify the curve intersection at the 50% inhibition value to estimate  $IC_{50}$  values for the inhibitors.

A total of 57 pooled master wells were selected for deconvolution, based on combined criteria of <42% residual activity at the tested concentration, and a total accepted hit rate of no more than 0.2%. At 22 compounds per well, a total of 1254 compounds were identified through this deconvolution and individually assayed at the 1X concentration of 27.7  $\mu$ M to identify which compounds exhibited the desired activity. From these assays, 73 compounds were selected and assayed further to develop  $IC_{50}$  curves. From the  $IC_{50}$  curve results, 34 compounds were selected for selectivity assays against PI3K $\alpha$  and PI3K $\beta$  (see assay protocol in Example 11).

From the selectivity assays, one compound, 3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-

- 93 -

3H-quinazolin-4-one (Compound D-000), was selected as being a relatively potent and selective compound. Catalog searches and selectivity assays of many analogous compounds of the potent and/or selective hits yielded only one compound that was both an active and selective analogue of D-000. This compound was purchased from Contract Services Corporation (Catalog #7232154) and differed from D-000 in substituting a phenyl group for the 2-chlorophenyl group of D-000.



As described above, the PI 3-kinase inhibitor LY294002 (Calbiochem, La Jolla, CA) does not have significant selectivity among the different PI 3-kinase isoforms tested. Under our assay conditions, LY294002 inhibited all three isoforms of PI 3-kinases with an  $IC_{50}$  of 0.3 to 1  $\mu$ M. However, when the compound D-000 was tested against the same PI 3-kinase isoforms distinct selectivity was observed. Specifically, as shown in Figure 1, D-000 inhibited the activity of the  $\delta$  isoform of PI3K with an  $IC_{50}$  of approximately 0.3  $\mu$ M, whereas under similar condi-

- 94 -

tions it did not inhibit activities of the  $\alpha$  and  $\beta$  isoforms at a limit of 100  $\mu$ M compound. These results show that D-000 selectively inhibits PI3K $\delta$  activity.

5

### EXAMPLES 3-7

Since PI3K $\delta$  is expressed at significant levels only in leukocytes, it is important to study the effects of the PI3K $\delta$ -selective inhibitor on leukocyte functions. Accordingly, the effects of PI3K $\delta$  inhibition in several types of leukocytes were examined. Neutrophils were examined to determine the effects that selective inhibition of PI3K $\delta$  might elicit (Example 3, below). It surprisingly was found that selective inhibition of PI3K $\delta$  activity appears to be significantly associated with inhibition of some but not all functions characteristic of activated neutrophils. In addition, the effects of PI3K $\delta$  inhibition on B cell and T cell function also were tested (Examples 4-5, below). Moreover, as PI3K $\delta$  also is expressed in osteoclasts, the effect of PI3K $\delta$  inhibition on the function of these specialized cells was studied (Example 6, below).

25

### EXAMPLE 3

#### Characterization of Role of PI3K $\delta$ in Neutrophil Function

30

The effects of a PI3K $\delta$  inhibitor of the invention, i.e., D-000, on neutrophil functions such

- 95 -

as superoxide generation, elastase exocytosis, chemotaxis, and bacterial killing were tested.

5                   A.    Preparation of neutrophils  
                    from human blood

                    Aliquots (8 mL) of heparinized blood from healthy volunteers were layered on 3 mL cushions of  
10                   7.3% FICOLL® (Sigma, St. Louis, MO) and 15.4%  
                    HYPAQUE® (Sigma) and centrifuged at 900 rpm for 30 min at room temperature in a table top centrifuge (Beckman). The neutrophil-rich band just above the FICOLL®-HYPAQUE® cushion was collected and washed  
15                   with Hanks' balanced salt solution (HBSS) containing 0.1% gelatin. Residual erythrocytes were removed by hypotonic lysis with 0.2% NaCl. The neutrophil preparation was washed twice with HBSS containing 0.1% gelatin and used immediately.

20

                    B.    Measurement of superoxide  
                    production from neutrophils

25                   Superoxide generation is one of the hallmarks of neutrophil activation. A variety of activators potentiate superoxide generation by neutrophils. The effect of the PI3Kδ inhibitor D-000 on superoxide generation by three different agonists:  
30                   TNF1α, IgG, and fMLP, each representing separate classes of activator, was measured. Superoxide generated by the neutrophils was measured by monitoring the change in absorbance upon reduction of cytochrome C by modification of the method described  
35                   by Green et al., (pp. 14.5.1-14.5.11 in Supp. 12,

- 96 -

*Curr Protocols Immunol* (Eds., Colligan et al.) (1994)), as follows. Individual wells of a 96-well plate were coated overnight at 4°C with 50 µL of 2 mg/mL solution of human fibrinogen or IgG. The wells were washed with PBS and the following reagents were added to each well: 50 µL of HBSS or superoxide dismutase (1 mg/mL), 50 µL of HBSS or TNF1α (50 ng/mL), 50 µL cytochrome C (2.7 mg/mL), and 100 µL of purified human neutrophil suspension (2 x 10<sup>6</sup> cells/mL). The plate was centrifuged for 2 min at 200 rpm and absorbance at 550 nm was monitored for 2 hr. To measure the relative amounts of superoxide generated, values obtained from the superoxide dismutase-containing wells were subtracted from all, and normalized to the values obtained from the wells without any inhibitor.

As shown in Figure 2, the PI3Kδ inhibitor D-000 inhibits TNF-induced superoxide generation by neutrophils in a concentration dependent manner. Superoxide generation induced by TNF was reduced to its half-maximal value at about 3 µM D-000. Figure 2 also reveals that superoxide generation induced by IgG was not significantly inhibited by D-000. In fact, even at 10 µM this PI3Kδ inhibitor did not have any effect on superoxide generation induced by IgG.

Next, the effect of D-000 on superoxide generation induced by another potent inducer, the bacterial peptide, formylated-Met-Leu-Phe (fMLP) was studied. Like the TNF-induced superoxide generation, fMLP-induced superoxide generation also was inhibited by D-000 (Figure 3). These results show that the PI3Kδ inhibitor D-000 can prevent stimulus

- 97 -

specific induction of superoxide generation by neutrophils, indicating that PI3K $\delta$  is involved in this process.

5                   C.    **Measurement of elastase exocytosis  
                          from neutrophils**

10                   In addition to superoxide generation,  
activated neutrophils also respond by releasing  
several proteases that are responsible for the de-  
struction of tissues and cartilage during inflamma-  
tion. As an indication of protease release, the  
effect of D-000 on elastase exocytosis was measured.  
15   Elastase exocytosis was quantitated by modification  
of the procedure described by Ossanna et al. (*J Clin  
Invest*, 77:1939-1951 (1986)), as follows. Purified  
human neutrophils ( $0.2 \times 10^6$ ) (treated with either  
DMSO or a serial dilution of D-000 in DMSO) were  
20   stimulated with fMLP in PBS containing 0.01 mg/mL  
cytochalasin B, 1.0  $\mu$ M sodium azide ( $\text{NaN}_3$ ), 5  $\mu$ g/mL  
L-methionine and 1  $\mu$ M fMLP for 90 min at 37°C in a  
96-well plate. At the end of the incubation period,  
the plate was centrifuged for 5 min at 1000 rpm, and  
25   90  $\mu$ L of the supernatant was transferred to 10  $\mu$ L of  
10 mM solution of an elastase substrate peptide,  
MeO-suc-Ala-Ala-Pro-Val-pNA, wherein MeO-suc =  
methoxy-succinyl; pNA = p-nitroanilide (Calbiochem,  
San Diego, CA). Absorbance at 410 nm was monitored  
30   for 2 hr in a 96-well plate reader. To measure the  
relative amounts of elastase exocytosed, all absorb-  
ance values were normalized to the values without  
any inhibitor. As shown in Figure 4, the PI3K $\delta$   
inhibitor D-000 inhibits fMLP-induced elastase

- 98 -

exocytosis significantly, and does so in a dose-dependent fashion. Inhibition was half-maximal at a concentration of about 2-3  $\mu$ M D-000.

5                   D.    Measurement of fMLP-induced human  
                    neutrophil migration

10                   Neutrophils have the intrinsic capacity to  
migrate through tissues, and are one of the first  
cell types to arrive at the sites of inflammation or  
tissue injury. The effect of D-000 on neutrophil  
migration towards a concentration gradient of fMLP  
was measured. The day before the migration assays  
15                   were performed, 6-well plates were coated with  
recombinant ICAM-1/Fc fusion protein (Van der Vieren  
et al., *Immunity*, 3:683-690 (1995)) (25  $\mu$ g/mL in  
bicarbonate buffer, pH 9.3) and left overnight at  
4°C. After washing, 1% agarose solution, in RPMI-  
20                   1640 with 0.5% bovine serum albumin (BSA), was added  
to wells with or without an inhibitor, and plates  
were placed into a refrigerator before punching  
holes in the gelled agarose to create plaques (1  
central hole surrounded by 6 peripheral ones per  
25                   well).

                    Human neutrophils were obtained as de-  
scribed above, and resuspended in RPMI medium  
supplemented with 0.5% BSA at  $5 \times 10^6$  cells/mL.  
After combining equal volumes of neutrophil suspen-  
30                   sion and medium (either with DMSO or a serial dilu-  
tion of the test compound in DMSO), neutrophils were  
aliquoted into the peripheral holes, while the  
central hole received fMLP (5  $\mu$ M). Plates were  
incubated at 37°C in the presence of 5% CO<sub>2</sub> for 4 hr,

- 99 -

followed by termination of migration by the addition of 1% glutaraldehyde solution in D-PBS. After removing the agarose layer, wells were washed with distilled water and dried.

5                   Analysis of neutrophil migration was conducted on a Nikon DIAPHOT® inverted microscope (1x objective) video workstation using the NIH 1.61 program. Using Microsoft Excel and Table Curve 4 (SSPS Inc., Chicago IL) programs, a migration index  
10                   was obtained for each of the studied conditions. Migration index was defined as the area under a curve representing number of migrated neutrophils versus the net distance of migration per cell.

                  As shown in Figure 5, the PI3Kδ inhibitor  
15                   D-000 had a profound effect on neutrophil migration, inhibiting this activity in a dose-dependent manner. The EC<sub>50</sub> of this compound for inhibition of neutrophil migration in this assay is about 1 μM. Based on a visual inspection of the recorded paths of the  
20                   cells in this assay, it appears that the total path length for the neutrophils was not significantly affected by the test compound. Rather, the compound affected neutrophil orientation or sense of direction, such that instead of migrating along the axis  
25                   of the chemoattractant gradient, the cells migrated in an undirected or less directed manner.

#### E.    Measurement of bactericidal capacity of neutrophils

30

                  Given that the PI3Kδ inhibitor D-000 affects certain neutrophil functions detailed above, it was of interest to see whether the compound



- 100 -

affects neutrophil-mediated bacterial killing. The effect of D-000 on neutrophil-mediated *Staphylococcus aureus* killing was studied according to the method described by Clark and Nauseef (pp. 7.23.4-7.23.6 in Vol. 2, Supp. 6, *Curr Protocols Immunol* (Eds., Colligan et al.) (1994)). Purified human neutrophils ( $5 \times 10^6$  cells/mL) (treated with either DMSO or a serial dilution of D-000 in DMSO) were mixed with autologous serum. Overnight-grown *S. aureus* cells were washed, resuspended in HBSS, and added to the serum-opsonized neutrophils at a 10:1 ratio. Neutrophils were allowed to internalize the bacteria by phagocytosis by incubation at 37°C for 20 min. The noninternalized bacteria were killed by 10 units/mL lysostaphin at 37°C for 5 min and the total mixture was rotated at 37°C. Samples were withdrawn at various times for up to 90 min and the neutrophils were lysed by dilution in water. Viable bacteria were counted by plating appropriate dilutions on trypticase-soy-agar plate and counting the *S. aureus* colonies after overnight growth.

As shown in Figure 6, neutrophil-mediated killing of *S. aureus* was similar in samples treated with DMSO (control) and with D-000. These results indicate that the PI3K $\delta$  inhibitor does not significantly affect the ability of neutrophils to kill *S. aureus*, suggesting that PI3K $\delta$  is not involved in this pathway of neutrophil function.

- 101 -

EXAMPLE 4Characterization of Role of  
PI3K $\delta$  in B Lymphocyte Function

5

The effects of the PI 3-kinase inhibitor on B cell functions including classical indices such as antibody production and specific stimulus-induced proliferation also were studied.

10

A. Preparation and stimulation of B  
cells from peripheral human blood

15

Heparinized blood (200 mL) from healthy volunteers was mixed with an equal volume of D-PBS, layered on 10 x 10 mL FICOLL-PAQUE® (Pharmacia), and centrifuged at 1600 rpm for 30 min at room temperature. Peripheral blood mononuclear cells (PBMC) were collected from the FICOLL®/serum interface, overlaid on 10 mL fetal bovine serum (FBS) and centrifuged at 800 rpm for 10 min to remove platelets. After washing, cells were incubated with DYNAL® Antibody Mix (B cell kit) (Dynal Corp., Lake Success, NY) for 20 min at 4-8°C. Following the removal of unbound antibody, PBL were mixed with anti-mouse IgG coated magnetic beads (Dynal) for 20 min at 4-8°C with gentle shaking followed by elimination of labeled non-B cells on the magnetic bead separator. This procedure was repeated once more. The B cells were resuspended in RPMI-1640 with 10% FBS, and kept on ice until further use.

25

30

- 102 -

B. Measurement of antibody production  
by human B cells

5                   To study antibody production, B cells were  
aliquoted at  $50-75 \times 10^3$  cells/well into 96-well  
plate with or without inhibitor, to which IL-2 (100  
U/mL) and PANSORBIN® (Calbiochem) *Staphylococcus*  
10 *aureus* cells (1:90,000) were added. Part of the  
media was removed after 24-36 hr, and fresh media  
(with or without inhibitor) and IL-2 were added.  
Cultures were incubated at 37°C, in the presence of  
a CO<sub>2</sub> incubator for additional 7 days. Samples from  
each condition (in triplicate) were removed, and  
15 analyzed for IgG and IgM, as measured by ELISA.  
Briefly, IMMULON® 4 96-well plates were coated (50  
µL/well) with either 150 ng/mL donkey antihuman IgG  
(H+L) (Jackson ImmunoResearch, West Grove PA), or 2  
µg/mL donkey antihuman IgG+IgM (H+L) (Jackson  
20 ImmunoResearch) in bicarbonate buffer, and left  
overnight at 4°C. After 3x washing with phosphate  
buffered saline containing 0.1% TWEEN®-80 (PBST)  
(350 µL/well), and blocking with 3% goat serum in  
PBST (100 µL/well) for 1 hr at room temperature,  
25 samples (100 µL/well) of B cell spent media diluted  
in PBST were added. For IgG plates the dilution  
range was 1:500 to 1:10000, and for IgM 1:50 to  
1:1000. After 1 hr, plates were exposed to biotin-  
conjugated antihuman IgG (100 ng/mL) or antihuman  
30 IgM (200 ng/mL) (Jackson ImmunoResearch) for 30 min,  
following by streptavidin-HRP (1:20000) for 30 min,  
and finally, to TMB solution (1:100) with H<sub>2</sub>O<sub>2</sub>  
(1:10000) for 5 min, with 3 x PBST washing between

- 103 -

steps. Color development was stopped by  $H_2SO_4$  solution, and plates were read on an ELISA plate reader.

As shown in Figure 7, D-000 significantly inhibited antibody production. IgM production was affected more than IgG production: half-maximal inhibition of IgM production was observed at about 1  $\mu M$ , versus about 7  $\mu M$  for comparable inhibition of IgG production.

10                   C.    Measurement of B Cell Proliferation  
                          in response to cell surface IgM  
                          stimulation

15                   In the above experiment, the B cells were  
stimulated using PANSORBIN®. The effect of D-000 on  
B cell proliferation response when they were stimulated  
through their cell surface IgM using anti-IgM antibody  
also was measured. Murine splenocytes  
20   (Balb/c) were plated into 96-well microtiter plates  
at  $2 \times 10^5$  cells per well in 10% FBS/RPMI. Appropriate  
dilutions of test inhibitor in complete medium were  
added to the cells and the plates were incubated for  
30-60 minutes prior to the addition of stimulus.  
25   Following the preincubation with test inhibitor an  
 $F(ab')_2$  preparation of goat antibody specific for the  
 $\mu$ -chain of mouse IgM was added to the wells at a  
final concentration of 25  $\mu g/mL$ . The plates were  
incubated at  $37^\circ C$  for 3 days and 1  $\mu Ci$  of [ $^3H$ ]-  
30   thymidine was added to each well for the final four  
hours of culture. The plates were harvested onto  
fiber filters washed and the incorporation of radio-  
label was determined using a beta counter (Matrix

- 104 -

96, Packard Instrument Co., Downers Grove, IL) and expressed as counts per minute (CPM).

Figure 8 shows the effect of D-000 on anti-IgM stimulated proliferation of B cells. The compound inhibited anti-IgM-stimulated B cell proliferation in a dose-dependent manner. At about 1  $\mu$ M, proliferation was reduced to its half-maximal value.

Because the compound D-000 inhibits B cell proliferation, it is envisioned that this compound and other PI3K $\delta$  inhibitors could be used to suppress undesirable proliferation of B cells in clinical settings. For example, in B cell malignancy, B cells of various stages of differentiation show unregulated proliferation. Based on the results shown above, one can infer that PI3K $\delta$  selective inhibitors could be used to control, limit, or inhibit growth of such cells.

#### 20 EXAMPLE 5

##### Characterization of Role of PI3K $\delta$ in T Lymphocyte Function

25 T cell proliferation in response to costimulation of CD3+CD28 was measured. T cells were purified from healthy human blood by negative selection using antibody coated magnetic beads according to the manufacturer's protocol (Dyna) and resuspended in RPMI. The cells were treated with either DMSO or a serial dilution of D-000 in DMSO and plated at  $1 \times 10^5$  cells/well on a 96-well plate precoated with goat antimouse IgG. Mouse monoclonal

- 105 -

anti-CD3 and anti-CD28 antibodies were then added to each well at 0.2 ng/mL and 0.2 µg/mL, respectively. The plate was incubated at 37°C for 24 hr and [<sup>3</sup>H]-thymidine (1 µCi/well) was added. After another 18 hr incubation the cells were harvested with an automatic cell harvester, washed and the incorporated radioactivity was quantified.

Although the PI3Kδ inhibitor D-000 inhibited anti-CD3- and anti-CD28-induced proliferation of T cells, its effect is not as strong as its effect on B cells or on some of the functions of neutrophils. Half-maximal inhibition of thymidine incorporation was not achieved at the highest tested concentration, i.e., 10 µM D-000.

#### EXAMPLE 6

##### Characterization of Role of PI3Kδ in Osteoclast Function

To analyze the effect of the PI3Kδ inhibitor D-000 on osteoclasts, mouse bone marrow cells were isolated and differentiated them to osteoclasts by treating the cells with Macrophage Colony Stimulating Factor<sup>-1</sup> (mCSF<sup>-1</sup>) and Osteoprotegerin Ligand (OPGL) in serum-containing medium (αMEM with 10% heat-inactivated FBS; Sigma) for 3 days. On day four, when the osteoclasts had developed, the medium was removed and cells were harvested. The osteoclasts were plated on dentine slices at 10<sup>5</sup> cells/-well in growth medium, i.e., αMEM containing 1% serum and 2% BSA with 55 µg/mL OPGL and 10 ng/mL mCSF-1. After 3 hr, the medium was changed to 1%

- 106 -

serum and 1% BSA, with or without osteopontin (25  $\mu\text{g/mL}$ ) and the PI3K inhibitors (100 nM). The medium was changed every 24 hours with fresh osteopontin and the inhibitors. At 72 hr, the medium was removed, and the dentine surfaces were washed with water to remove cell debris and stained with acid hematoxylin. Excess stain was washed and the pit depths were quantitated using confocal microscopy.

As shown in Table 1, in two experiments, the PI 3-kinase inhibitors had an inhibitory effect on osteoclast function. Both the nonspecific inhibitors LY294002 and wortmannin inhibited osteoclast activity. However, the PI3K $\delta$  inhibitor D-000 had the most profound effect, as at 100 nM this compound almost completely inhibited the osteoclast activity.

Table 1			
Osteopontin (OPN)	D-000 + OPN	LY294002 + OPN	Wortmannin + OPN
10 $\pm$ 0.5	1	4.6 $\pm$ 0.22	5.7 $\pm$ 0.6
9 $\pm$ 0.4	1	5.8 $\pm$ 0.5	5 $\pm$ 0.5

#### EXAMPLE 7


##### Characterization of Role of PI3K $\delta$ in Basophil Function

Assessment of the effect of a compound of the invention on basophil function was tested using a conventional histamine release assay, generally in accordance with the method described in Miura et al., *J Immunol*, 162:4198-206 (1999). Briefly, enriched basophils were preincubated with test com-

- 107 -

pounds at several concentrations from 0.1 nM to 1,000 nM, for 10 min at 37°C. Then, polyclonal goat antihuman IgE (0.1 µg/mL) or fMLP was added, and allowed to incubate for an additional 30 min.

5 Histamine released into the supernatant was measured using an automated fluorometric technique. Two compounds were tested, shown below.




20

D-026



- 108 -



D-999

A dose-dependent decrease in histamine release was observed for 3-(2-chlorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one (D-026) when the basophils were stimulated with anti-IgE. This suppression of histamine release was essentially 100% at 1,000 nM, with an  $EC_{50}$  of about 25 nM. Another compound, 3-(2-chlorophenyl)-2-(1H-pyrazolo[3,4-d]pyrimidin-4-ylsulfanylmethyl)-3H-quinazolin-4-one (D-999), in which the purine ring structure is rearranged, was less efficacious in the inhibition of histamine release. Neither compound elicited any effect when the basophils were stimulated with fMLP. For comparison, the nonselective PI3K inhibitor LY294002 was tested at 0.1 nM and 10,000 nM, showing close to 100% inhibition of histamine release at the highest concentration.

These data indicate that inhibitors of PI 3-kinase delta activity can be used to suppress release of histamine, which is one of the mediators of allergy. Since the activity of various PI 3-

- 109 -

kinases are required for protein trafficking, secretion, and exocytosis in many cell types, the above data suggest that histamine release by other cells, such as mast cells, also can be disrupted by PI 3-kinase delta-selective inhibitors.

#### CHEMICAL SYNTHESIS EXAMPLES

Specific nonlimiting examples of compounds of the invention are provided below. It is understood in the art that protecting groups can be employed where necessary in accordance with general principles of synthetic chemistry. These protecting groups are removed in the final steps of the synthesis under basic, acidic, or hydrogenolytic conditions readily apparent to those persons skilled in the art. By employing appropriate manipulation and protection of any chemical functionalities, synthesis of compounds of structural formula (I) not specifically set forth herein can be accomplished by methods analogous to the schemes set forth below.

Unless otherwise noted, all starting materials were obtained from commercial suppliers and used without further purification. All reactions and chromatography fractions were analyzed by thin-layer chromatography (TLC) on 250 mm silica gel plates, visualized with ultraviolet (UV) light or iodine (I<sub>2</sub>) stain. Products and intermediates were purified by flash chromatography or reverse-phase high performance liquid chromatography.

The following abbreviations are used in the synthetic examples: aq (aqueous), H<sub>2</sub>O (water), CHCl<sub>3</sub> (chloroform), HCl (hydrochloric acid), MeOH

- 110 -

(methanol), NaOH (sodium hydroxide), NaOMe (sodium methoxide), TFA (trifluoroacetic acid),  $K_2CO_3$  (potassium carbonate),  $SOCl_2$  (thionyl chloride),  $CH_2Cl_2$  (methylene chloride), EtOAc (ethyl acetate),  
5 DMF (dimethylformamide), EtOH (ethanol), DMSO (dimethyl sulfoxide),  $NaHCO_3$  (sodium bicarbonate), TLC (thin layer chromatography), HPLC (high performance liquid chromatography), HOBT (hydroxybenzotriazole), EDC (ethyldiethylaminopropylcarbodiimide), DIEA  
10 (diisopropylethylamine), and HOAc (acetic acid).

### I. General Procedures

#### Procedure A

15 Thionyl chloride was added to a rapidly stirring solution of anthranilic acid or benzoic acid in benzene, and the mixture was stirred at reflux for 5 to 18 hours. The reaction was  
20 concentrated in vacuo, and stripped down twice with benzene. The resulting oil was dissolved in  $CHCl_3$ , and to that solution was added the appropriate aniline. The reaction mixture was heated to reflux and stirred until complete, as determined by TLC, at  
25 which point the reaction mixture was cooled to ambient temperature. The precipitate was removed by filtration, and the filtrate concentrated in vacuo. The crude product was purified by chromatography and/or recrystallization from MeOH to provide amides  
30 1a-1r.

- 111 -

Procedure B

To a rapidly stirring suspension of an amide in glacial acetic acid was added chloroacetyl chloride. The reaction mixture was heated to 120°C, and allowed to stir at that temperature until complete, as determined by TLC. After brief cooling, the reaction mixture was concentrated in vacuo. The crude residue was purified by extraction, chromatography, and/or recrystallization to provide chlorides 2a-2r.

Procedure C

A mixture of a chloride, either a nitrogen or a sulfur nucleophile, for example, mercaptopurine monohydrate or adenine, and  $K_2CO_3$  in DMF was stirred at room temperature for 15-72 hours. The resulting suspension was poured into water, and kept at 4°C for several hours. The crude solid was filtered, washed with water, and purified by chromatography or recrystallization to provide the final product.

EXAMPLE 8Preparation of Intermediate Compounds: Amides2-Amino-N-(2-chlorophenyl)-4,5-dimethoxybenzamide (1a)

Prepared according to Procedure A using 4,5-dimethoxyanthranilic acid (5.0 g, 25.4 mmol) and  $SOCl_2$  (5.5 mL, 76.1 mmol) in benzene (100 mL),

- 112 -

followed by 2-chloroaniline (6.7 mL, 63.5 mmol) and  
CHCl<sub>3</sub> (75 mL). The product was washed with aqueous  
NaHCO<sub>3</sub> (2 x 25 mL) and HCl (0.5 M, 75 mL) and puri-  
fied by chromatography in CH<sub>2</sub>Cl<sub>2</sub> to provide 4.3 g of  
5 a brown foam (55%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.42 (dd,  
J=1.5, 8.3 Hz, 1H); 8.32 (br s, 1H); 7.40 (dd,  
J=1.4, 8.0 Hz, 1H); 7.31 (dt, J=1.4, 7.9 Hz, 1H);  
7.05 (dt, J=1.5, 7.7 Hz, 1H); 7.03 (s, 1H); 6.24 (s,  
1H); 3.88 (s, 3H); 3.87 (s, 3H). MS (ES): m/z  
10 307.0 (M+).

**2-Amino-5-bromo-N-(2-chlorophenyl)benzamide (1b)**

15 Prepared according to Procedure A using 2-  
amino-5-bromobenzoic acid (5.0 g, 23.1 mmol) and  
SOCl<sub>2</sub> (7.0 mL, 95.9 mmol) in benzene (50 mL),  
followed by 2-chloroaniline (7.3 mL, 69.3 mmol) and  
CHCl<sub>3</sub> (50 mL). The product was purified by two  
20 chromatographies in CH<sub>2</sub>Cl<sub>2</sub> to provide 1.48 g of a  
yellow orange solid (20%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.36  
(dd, J=1.2, 8.2 Hz, 1H); 8.20 (br s, 1H); 7.62 (d,  
J=2.1 Hz, 1H); 7.42 (dd, J=1.3, 8.0 Hz, 1H); 7.34  
(dd, J=2.2, 8.8 Hz, 1H); 7.28-7.33 (m, 1H); 7.09  
25 (dt, J=1.4, 7.7 Hz, 1H); 6.62 (d, J=8.7 Hz, 1H);  
5.57 (br s, 2H).

**2-Amino-N-(2-chlorophenyl)-4-fluorobenzamide (1c)**

30 Prepared according to Procedure A using 2-  
amino-4-fluorobenzoic acid (1.15 g, 7.41 mmol) and  
SOCl<sub>2</sub> (1.4 mL, 18.5 mmol) in benzene (25 mL),  
followed by 2-chloroaniline (1.6 mL, 14.8 mmol) and  
35 CHCl<sub>3</sub> (25 mL). The product was chromatographed in

- 113 -

CH<sub>2</sub>Cl<sub>2</sub>, then triturated from hexanes to provide 1.02 g of an off-white solid (52%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 12.91 (br s, 1H); 8.72 (dd, J=2.7, 12 Hz, 1H); 8.34 (dd, J=6.4, 9.2 Hz, 1H); 8.29 (dd, J=5.9, 8.8 Hz, 1H); 7.81 (dd, J=6.2, 8.8 Hz, 1H); 7.28 (dt, J=2.4, 8.4 Hz, 1H); 7.21 (dd, J=2.4, 9.0 Hz, 1H); 6.92 (ddd, J=2.4, 7.3, 9.1 Hz, 1H); 6.54 (ddd, J=2.4, 7.8, 8.8 Hz, 1H); 6.45 (dd, J=2.4, 11 Hz, 1H); 5.93 (br s, 2H). MS (ES): m/z 265.0 (M+).

**2-Amino-5-chloro-N-(2-chlorophenyl)benzamide (1d)**

Prepared according to Procedure A using 2-amino-5-chlorobenzoic acid (2.0 g, 11.7 mmol) and SOCl<sub>2</sub> (2.2 mL, 29.2 mmol) in benzene (50 mL), followed by 2-chloroaniline (2.5 mL, 23.3 mmol) and CHCl<sub>3</sub> (50 mL). The product was purified by recrystallization from MeOH to provide 1.72 g of a dark yellow solid (52%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.37 (dd, J=1.5, 8.3 Hz, 1H); 8.22 (br s, 1H); 7.48 (d, J=2.3 Hz, 1H); 7.42 (dd, J=1.5, 8.1 Hz, 1H); 7.31 (dt, J=1.4, 7.8 Hz, 1H); 7.22 (dd, J=2.4, 8.8 Hz, 1H); 7.09 (dt, J=1.5, 7.7 Hz, 1H); 6.67 (d, J=8.8 Hz, 1H); 5.56 (br s, 2H).

**2-Amino-N-(2-chlorophenyl)-6-fluorobenzamide (1e)**

Prepared according to Procedure A using 2-amino-6-fluorobenzoic acid (2.0 g, 12.9 mmol) and SOCl<sub>2</sub> (2.3 mL, 32.2 mmol) in benzene (50 mL), followed by 2-chloroaniline (2.7 mL, 25.8 mmol) and CHCl<sub>3</sub> (50 mL). The product was purified by chromatography in EtOAc/hexanes to provide 2.06 g of a

- 114 -

pale orange solid (60%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 9.00 (d, J=17 Hz, 1H); 8.47 (d, J=8.3 Hz, 1H); 7.41 (d, J=8.0 Hz, 1H); 7.30 (t, J=7.9 Hz, 1H); 7.10-7.20 (m, 1H); 7.07 (t, J=7.7 Hz, 1H); 6.49 (d, J=8.3 Hz, 1H); 6.03 (br s, 2H). MS (ES): m/z 265.0 (M+).

**2-Amino-6-chloro-N-(2-chlorophenyl)benzamide (1f)**

Prepared according to Procedure A using 2-amino-6-chlorobenzoic acid (2.5 g, 14.6 mmol) and SOCl<sub>2</sub> (2.7 mL, 36.4 mmol) in benzene (75 mL), followed by 2-chloroaniline (3.1 mL, 29.1 mmol) and CHCl<sub>3</sub> (75 mL). The product chromatographed in CH<sub>2</sub>Cl<sub>2</sub> to provide 1.05 g of a yellow orange solid (26%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.54 (d, J=8.1 Hz, 1H); 8.30 (br s, 1H); 7.41 (dd, J=1.5, 8.0 Hz, 1H); 7.33 (t, J=7.8 Hz, 1H); 7.10 (t, J=8.1 Hz, 1H); 7.09 (dt, J=1.6, 7.8 Hz, 1H); 6.78 (dd, J=0.4, 7.9 Hz, 1H); 6.63 (dd, J=0.9, 8.2 Hz, 1H); 4.69 (br s, 2H). MS (ES): m/z 303.0 (M+22), 281.0 (M+).

**2-Amino-N-(2-chlorophenyl)-6-methylbenzamide (1g)**

Prepared according to Procedure A using 2-amino-6-methylbenzoic acid (2.5 g, 16.5 mmol) and SOCl<sub>2</sub> (3.0 mL, 41.3 mmol) in benzene (75 mL), followed by 2-chloroaniline (3.5 mL, 33.0 mmol) and CHCl<sub>3</sub> (75 mL). The product was chromatographed in CH<sub>2</sub>Cl<sub>2</sub> to provide 2.19 g of a brown oil (51%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.58 (d, J=8.1 Hz, 1H); 7.99 (br s, 1H); 7.40 (dd, J=1.4, 8.0 Hz, 1H); 7.34 (t, J=7.7 Hz, 1H); 7.11 (t, J=7.8 Hz, 1H); 7.09 (dt, J=1.5, 7.7 Hz, 1H); 6.64 (d, J=7.5 Hz, 1H); 6.59 (d, J=8.1

- 115 -

Hz, 1H); 4.29 (br s, 2H); 2.45 (s, 3H). MS (ES):  
m/z 283.0 (M+22).

**2-Amino-3-chloro-N-(2-chlorophenyl)benzamide (1h)**

5

Prepared according to Procedure A using 2-amino-3-chlorobenzoic acid (1.0 g, 5.82 mmol) and SOCl<sub>2</sub> (1.1 mL, 14.6 mmol) in benzene (25 mL),  
10 followed by 2-chloroaniline (1.2 mL, 11.7 mmol) and CHCl<sub>3</sub> (25 mL). The product was recrystallized from MeOH to provide 1.29 g of a yellow solid (78%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.43 (dd, J=1.4, 8.3 Hz, 1H); 8.30 (br s, 1H); 7.47 (dd, J=1.1, 8.0 Hz, 1H); 7.42 (d, J=8.0 Hz, 2H); 7.33 (dt, J=1.4, 7.9 Hz, 1H); 7.09 (dt, J=1.5, 7.7 Hz, 1H); 6.68 (t, J=7.9 Hz, 1H);  
15 6.13 (br s, 2H). MS (ES): m/z 281.0 (M+).

**2-Amino-N-biphenyl-2-yl-6-chlorobenzamide (1i)**

20

Prepared according to Procedure A using 2-amino-6-chlorobenzoic acid (2.0 g, 11.7 mmol) and SOCl<sub>2</sub> (2.1 mL, 29.3 mmol) in benzene (60 mL),  
25 followed by 2-aminobiphenylamine (4.15 g, 24.5 mmol) and CHCl<sub>3</sub> (60 mL). The product was chromatographed in CH<sub>2</sub>Cl<sub>2</sub> to provide 2.16 g of a foamy dark-amber residue (57%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.48 (d, J=8.2 Hz, 1H); 7.79 (br s, 1H); 7.34-7.46 (m, 6H); 7.20-7.30 (m, 2H); 7.00 (t, J=8.1 Hz, 1H); 6.63 (dd, J=0.6, 7.9 Hz, 1H); 6.54 (d, J=8.3 Hz, 1H); 4.58 (br s, 2H). MS (ES): m/z 323.1 (M+).

30



- 116 -

**2-Amino-6-chloro-N-o-tolylbenzamide (1j)**

Prepared according to Procedure A using 2-amino-6-chlorobenzoic acid (1.0 g, 5.83 mmol) and SOCl<sub>2</sub> (1.1 mL, 14.6 mmol) in benzene (30 mL), followed by o-toluidine (1.4 mL, 12.8 mmol) and CHCl<sub>3</sub> (30 mL). The product was chromatographed in CH<sub>2</sub>Cl<sub>2</sub> to provide 840 mg of an oily yellow solid (55%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 7.96 (d, J=7.9 Hz, 1H); 7.60 (br s, 1H); 7.23-7.30 (m, 2H); 7.14 (t, J=7.5 Hz, 1H); 7.11 (t, J=8.3 Hz, 1H); 6.78 (d, J=7.9 Hz, 1H); 6.64 (d, J=8.2 Hz, 1H); 4.73 (br s, 2H); 2.35 (s, 3H). MS (ES): m/z 261.0 (M+).

**2-Amino-6-chloro-N-(2-fluorophenyl)benzamide (1k)**

Prepared according to Procedure A using 2-amino-6-chlorobenzoic acid (2.0 g, 11.7 mmol) and SOCl<sub>2</sub> (2.1 mL, 29.1 mmol) in benzene (60 mL), followed by 2-fluoroaniline (2.3 mL, 23.4 mmol) and CHCl<sub>3</sub> (60 mL). The product was chromatographed in CH<sub>2</sub>Cl<sub>2</sub> to provide 1.05 g of a yellow solid (34%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.45 (t, J=8.0 Hz, 1H); 8.01 (br s, 1H); 7.02-7.22 (m, 4H); 6.78 (dd, J=0.5, 7.9 Hz, 1H); 6.64 (dd, J=0.8, 8.2 Hz, 1H); 4.73 (br s, 2H). MS (ES): m/z 265.0 (M+).

- 117 -

**2-Amino-6-chloro-N-(2-methoxyphenyl)benzamide (11)**

Prepared according to Procedure A using 2-amino-6-chlorobenzoic acid (2.0 g, 11.7 mmol) and SOCl<sub>2</sub> (2.1 mL, 29.1 mmol) in benzene (60 mL), followed by o-anisidine (2.6 mL, 23.4 mmol) and CHCl<sub>3</sub> (60 mL). The product was chromatographed in CH<sub>2</sub>Cl<sub>2</sub> to provide 2.61 g of a dark yellow oil (81%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.53 (dd, J=1.7, 7.9 Hz, 1H); 8.39 (br s, 1H); 7.11 (dt, J=1.6, 7.8 Hz, 1H); 7.09 (t, J=8.1 Hz, 1H); 7.02 (dt, J=1.4, 7.8 Hz, 1H); 6.92 (dd, J=1.4, 8.0 Hz, 1H); 6.62 (dd, J=0.9, 8.2 Hz, 1H); 4.66 (br s, 2H); 3.87 (s, 3H). MS (ES): m/z 277.0 (M+).

**2-Amino-N-(2-chlorophenyl)-3-trifluoromethylbenzamide (1m)**

Prepared according to Procedure A using 3-trifluoromethylanthranilic acid (2.0 g, 9.75 mmol) and SOCl<sub>2</sub> (1.8 mL, 24.4 mmol) in benzene (50 mL), followed by 2-chloroaniline (2.1 mL, 19.5 mmol) and CHCl<sub>3</sub> (50 mL). The product was purified by recrystallization from MeOH to provide 2.38 g yellow crystals (78%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.40 (dd, J=1.4, 8.3 Hz, 1H); 8.25 (br s, 1H); 7.71 (d, J=7.8 Hz, 1H); 7.60 (d, J=7.8 Hz, 1H); 7.43 (dd, J=1.4, 8.0 Hz, 1H); 7.34 (dt, J=1.3, 7.9 Hz, 1H); 7.11 (dt, J=1.5, 7.7 Hz, 1H); 6.77 (t, J=7.8 Hz, 1H); 6.24 (br s, 2H). MS (ES): m/z 315.0 (M+).

- 118 -

**3-Aminonaphthalene-2-carboxylic acid (2-chlorophenyl)amide (1n)**

5                   Prepared according to Procedure A using 3-amino-2-napthoic acid (2.0 g, 10.7 mmol) and SOCl<sub>2</sub> (1.9 mL, 26.7 mmol) in benzene (50 mL), followed by 2-chloroaniline (2.3 mL, 21.4 mmol) and CHCl<sub>3</sub> (50 mL). The product was recrystallized from MeOH to  
10 provide 1.71 g of a brown solid (54%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 10.88 (br s, 1H); 9.21 (s, 1H); 8.91 (s, 1H); 8.70 (dd, J=1.0, 8.3 Hz, 1H); 7.95-8.01 (m, 1H); 7.87-7.94 (m, 1H); 7.60-7.68 (m, 2H); 7.41 (dd, J=1.3, 8.0 Hz, 1H); 7.34 (dt, J=1.2, 7.8 Hz, 1H);  
15 7.07 (dt, J=1.4, 7.7 Hz, 1H). MS (ES): m/z 297.1 (M+).

**2-Amino-N-(2-chlorophenyl)-4-nitrobenzamide (1o)**

20                   Prepared according to Procedure A using 4-nitroanthranilic acid (5.0 g, 27.5 mmol) and SOCl<sub>2</sub> (5.0 mL, 68.6 mmol) in benzene (150 mL), followed by 2-chloroaniline (5.8 mL, 55.0 mmol) and CHCl<sub>3</sub> (150  
25 mL). The product was purified by chromatography in CH<sub>2</sub>Cl<sub>2</sub> followed by recrystallization from MeOH to provide 2.20 g of an orange-brown solid (31%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.41 (dd, J=1.3, 8.3 Hz, 1H); 8.31 (br s, 1H); 7.67 (d, J=8.6 Hz, 1H); 7.57 (d, J=2.1  
30 Hz, 1H); 7.52 (dd, J=2.2, 8.5 Hz, 1H); 7.44 (dd, J=1.3, 8.1 Hz, 1H); 7.35 (dt, J=1.3, 7.9 Hz, 1H); 7.13 (dt, J=1.4, 7.8 Hz, 1H); 5.88 (br s, 2H). MS (ES): m/z 292.0 (M+).

- 119 -

**2-Amino-N-(2-chlorophenyl)-5-hydroxybenzamide (1p)**

Prepared according to Procedure A using 2-amino-5-hydroxybenzoic acid (5.0 g, 32.7 mmol) and SOCl<sub>2</sub> (6.0 mL, 81.6 mmol) in benzene (150 mL), followed by 2-chloroaniline (6.9 mL, 65.4 mmol) and CHCl<sub>3</sub> (150 mL). The product was purified by two chromatographies in MeOH/CH<sub>2</sub>Cl<sub>2</sub> to provide 990 mg of a brown solid (12%). <sup>1</sup>H NMR (MeOH-d<sub>4</sub>) δ: 7.92 (dd, J=1.6, 8.1 Hz, 1H); 7.48 (dd, J=1.5, 7.7 Hz, 1H); 7.34 (dt, J=1.5, 7.7 Hz, 1H); 7.20 (dt, J=1.7, 7.7 Hz, 1H); 7.16 (d, J=2.7 Hz, 1H); 6.83 (dd, J=2.7, 8.7 Hz, 1H); 6.76 (d, J=8.7 Hz, 1H); [6.24 (br s, 2H)]. MS (ES): m/z 263.0 (M+).

**2-Amino-N-(2-chlorophenyl)-4,5-difluorobenzamide (1q)**

Prepared according to Procedure A using 4,5-difluoroanthranilic acid (2.0 g, 11.6 mmol) and SOCl<sub>2</sub> (2.1 mL, 28.9 mmol) in benzene (60 mL), followed by 2-chloroaniline (2.4 mL, 23.2 mmol) and CHCl<sub>3</sub> (60 mL). The product was purified by two chromatographies in CH<sub>2</sub>Cl<sub>2</sub> and EtOAc/hexanes to provide 769 mg of a yellow solid (23%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.69-8.82 (m, 1H); 8.00 (dd, J=8.4, 9.0 Hz, 1H); 7.90 (dd, J=8.9, 12 Hz, 1H); 7.39 (dd, J=6.8, 10 Hz, 1H); 6.53 (dd, J=6.6, 12 Hz, 1H); 6.41 (br s, 2H); 5.79 (br s, 1H). MS (ES): m/z 283.1 (M+).

- 120 -

**2-Amino-N-(2-chlorophenyl)-5-fluorobenzamide (1r)**

Prepared according to Procedure A using 2-amino-5-fluorobenzoic acid (1.0 g, 6.45 mmol) and SOCl<sub>2</sub> (1.2 mL, 16.1 mmol) in benzene (30 mL), followed by 2-chloroaniline (1.4 mL, 12.9 mmol) and CHCl<sub>3</sub> (30 mL). The product was triturated from CH<sub>2</sub>Cl<sub>2</sub> to provide 985 mg of a mustard-yellow solid (58%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 7.66 (dd, J=2.9, 8.7 Hz, 1H); 7.52-7.55 (m, 1H); 7.32-7.37 (m, 3H); 7.09 (dt, J=3.0, 8.5 Hz, 1H); 6.71 (dd, J=4.3, 8.7 Hz, 1H). MS (ES): m/z 305.0 (M+40).

15

**EXAMPLE 9****Preparation of Intermediate Compounds: Chlorides**

20 **2-Chloromethyl-3-(2-chlorophenyl)-6,7-dimethoxy-3H-quinazolin-4-one (2a)**

Prepared according to Procedure B with 1a (2.95 g, 9.63 mmol) and chloroacetyl chloride (2.3 mL, 28.9 mmol) in acetic acid (30 mL). Purified by extraction from aq. K<sub>2</sub>CO<sub>3</sub> and recrystallization from isopropanol to afford 1.61 g of a brown crystalline solid (46%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 7.59-7.66 (m, 2H); 7.45-7.56 (m, 3H); 7.20 (s, 1H); 4.37 (d, J=12 Hz, 1H), 4.08 (d, J=12 Hz, 1H); 4.04 (s, 3H); 4.00 (s, 3H). MS (ES): m/z 365.0 (M+).

- 121 -

**6-Bromo-2-chloromethyl-3-(2-chlorophenyl)-3H-quinazolin-4-one (2b)**

5                   Prepared according to Procedure B with 1b  
(500 mg, 1.54 mmol) and chloroacetyl chloride (0.37  
mL, 4.61 mmol) in acetic acid (10 mL). Purified by  
recrystallization from isopropanol to afford 490 mg  
of an off-white solid (83%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.43  
10 (d, J=2.3 Hz, 1H); 7.91 (dd, J=2.3, 8.7 Hz, 1H);  
7.67 (d, J=8.7 Hz, 1H); 7.60-7.65 (m, 1H); 7.47-7.56  
(m, 2H); 7.52 (t, J=5.3 Hz, 1H); 7.47-7.56 (m, 1H);  
4.37 (d, J=12 Hz, 1H), 4.06 (d, J=12 Hz, 1H). MS  
(ES): m/z 385.0 (M+).

15

**2-Chloromethyl-3-(2-chlorophenyl)-7-fluoro-3H-quinazolin-4-one (2c)**

20                   Prepared according to Procedure B with 1c  
(500 mg, 1.89 mmol) and chloroacetyl chloride (0.45  
mL, 5.67 mmol) in acetic acid (10 mL). Purified by  
extraction from aqueous K<sub>2</sub>CO<sub>3</sub>, followed by recrystallization from isopropanol to afford 501 mg of a  
25 yellow crystalline solid (82%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ:  
8.32 (dd, J=6.0, 8.9 Hz, 1H); 7.59-7.66 (m, 1H);  
7.50-7.55 (m, 3H); 7.44 (dd, J=2.4, 9.4 Hz, 1H);  
7.27 (dt, J=2.5, 8.5 Hz, 1H); 4.37 (d, J=12 Hz, 1H),  
4.07 (d, J=12 Hz, 1H). MS (ES): m/z 323.0 (M+).

30

- 122 -

**6-Chloro-2-chloromethyl-3-(2-chlorophenyl)-3H-quinazolin-4-one (2d)**

5                   Prepared according to Procedure B with 1d  
(500 mg, 1.78 mmol) and chloroacetyl chloride (0.42  
mL, 5.33 mmol) in acetic acid (10 mL). Purified by  
recrystallization from isopropanol to afford 555 mg  
of a yellow solid (92%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.27 (d,  
10 J=1.9 Hz, 1H); 7.74-7.78 (m, 2H); 7.60-7.66 (m, 1H);  
7.48-7.57 (m, 3H); 4.37 (d, J=12 Hz, 1H), 4.07 (d,  
J=12 Hz, 1H). MS (ES): m/z 339.0 (M+).

**2-Chloromethyl-3-(2-chlorophenyl)-5-fluoro-3H-quinazolin-4-one (2e)**

15                   Prepared according to Procedure B with 1e  
(500 mg, 1.89 mmol) and chloroacetyl chloride (0.45  
20 mL, 5.67 mmol) in acetic acid (10 mL). Purified by  
extraction from aq. K<sub>2</sub>CO<sub>3</sub> and recrystallization from  
isopropanol to afford 430 mg of an off-white  
crystalline solid (70%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 7.76  
(dt, J=5.3, 8.2 Hz, 1H); 7.56-7.65 (m, 2H); 7.47-  
25 7.56 (m, 3H); 7.16-7.25 (m, 1H); 4.35 (d, J=12 Hz,  
1H), 4.07 (d, J=12 Hz, 1H). MS (ES): m/z 323.0  
(M+).

**5-Chloro-2-chloromethyl-3-(2-chlorophenyl)-3H-quinazolin-4-one (2f)**

30                   Prepared according to Procedure B with 1f  
(1.00 g, 3.56 mmol) and chloroacetyl chloride (0.85  
35 mL, 10.7 mmol) in acetic acid (15 mL). Purified by  
recrystallization from isopropanol to afford 791 mg  
of an off-white crystalline solid (65%). <sup>1</sup>H NMR

- 123 -

(CDCl<sub>3</sub>)  $\delta$ : 7.70 (s, 1H); 7.68 (d, J=3.8 Hz, 1H);  
7.61-7.65 (m, 1H); 7.55 (dd, J=2.7, 6.4 Hz, 1H);  
7.51 (d, J=3.1 Hz, 1H); 7.50 (s, 2H); 4.35 (d, J=12  
Hz, 1H), 4.05 (d, J=12 Hz, 1H). MS (ES): m/z 339.0  
(M+).

**2-Chloromethyl-3-(2-chlorophenyl)-5-methyl-3H-  
quinazolin-4-one (2g)**

Prepared according to Procedure B with 1g  
(2.18 g, 8.36 mmol) and chloroacetyl chloride (2.0  
mL, 25.1 mmol) in acetic acid (40 mL). Purified by  
two chromatographies in CH<sub>2</sub>Cl<sub>2</sub> and EtOAc/hexanes,  
followed by recrystallization from isopropanol to  
afford 638 mg of an off-white crystalline solid  
(24%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ : 7.73-7.80 (m, 3H); 7.58-  
7.64 (m, 3H); 7.41 (d, J=7.4 Hz, 1H); 4.40 (d, J=12  
Hz, 1H), 4.26 (d, J=12 Hz, 1H); 2.74 (s, 3H). MS  
(ES): m/z 319.0 (M+).

**8-Chloro-2-chloromethyl-3-(2-chlorophenyl)-3H-  
quinazolin-4-one (2h)**

Prepared according to Procedure B with 1h  
(500 mg, 1.78 mmol) and chloroacetyl chloride (0.49  
mL, 6.13 mmol) in acetic acid (10 mL). Purified by  
extraction from aqueous K<sub>2</sub>CO<sub>3</sub>, followed by recrystallization from isopropanol to afford 448 mg of a  
yellow solid (74%). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$ : 8.23 (dd,  
J=1.4, 8.0 Hz, 1H); 7.90 (dd, J=1.4, 7.8 Hz, 1H);  
7.61-7.66 (m, 1H); 7.51-7.55 (m, 3H); 7.47 (t, J=8.0  
Hz, 1H); 4.48 (d, J=12 Hz, 1H), 4.12 (d, J=12 Hz,  
1H). MS (ES): m/z 339.0 (M+).



- 124 -

**3-Biphenyl-2-yl-5-chloro-2-chloromethyl-3H-quinazolin-4-one (2i)**

5                   Prepared according to Procedure B with 1i  
(2.0 g, 6.20 mmol) and chloroacetyl chloride (1.5  
mL, 18.6 mmol) in acetic acid (30 mL). Purified by  
chromatography in CH<sub>2</sub>Cl<sub>2</sub>, followed by recrystalli-  
zation from isopropanol to afford 1.44 g of an off-  
10 white solid (61%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 7.61-7.64 (m,  
1H); 7.58-7.59 (m, 1H); 7.54-7.57 (m, 2H); 7.52-7.53  
(m, 1H); 7.45-7.52 (m, 2H); 7.24 (s, 5H); 3.92-4.03  
(m, 2H). MS (ES): m/z 381.0 (M+).

15                   **5-Chloro-2-chloromethyl-3-o-tolyl-3H-quinazolin-4-**  
**one (2j)**

                  Prepared according to Procedure B with 1j  
20 (750 mg, 2.88 mmol) and chloroacetyl chloride (0.69  
mL, 8.63 mmol) in acetic acid (15 mL). Purified by  
chromatography in CH<sub>2</sub>Cl<sub>2</sub>, followed by recrystalli-  
zation from isopropanol to afford 340 mg of a white  
solid (37%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 7.69 (d, J=2.1 Hz,  
25 1H); 7.68 (q, J=7.4 Hz, 1H); 7.54 (dd, J=2.2, 7.0  
Hz, 1H); 7.35-7.47 (m, 3H); 7.21-7.25 (m, 1H); 4.27  
(d, J=12 Hz, 1H); 4.11 (d, J=12 Hz, 1H); 2.18 (s,  
3H). MS (ES): m/z 319.0 (M+).

30                   **5-Chloro-2-chloromethyl-3-(2-fluorophenyl)-3H-**  
**quinazolin-4-one (2k)**

                  Prepared according to Procedure B with 1k  
35 (1.0 g, 3.78 mmol) and chloroacetyl chloride (0.90  
mL, 11.3 mmol) in acetic acid (20 mL). Purified by  
chromatography in CH<sub>2</sub>Cl<sub>2</sub> to afford 484 mg of a pale

- 125 -

pink solid (40%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 7.69 (s, 1H); 7.68 (d, J=3.2 Hz, 1H); 7.56 (d, J=3.0 Hz, 1H); 7.54 (d, J=3.0 Hz, 1H); 7.40-7.47 (m, 1H); 7.35-7.38 (m, 1H); 7.27-7.32 (m, 1H); 4.35 (d, J=12 Hz, 1H); 4.18 (d, J=12 Hz, 1H). MS (ES): m/z 323.0 (M+).

**5-Chloro-2-chloromethyl-3-(2-methoxyphenyl)-3H-quinazolin-4-one (21)**

10

Prepared according to Procedure B with 11 (2.6 g, 9.41 mmol) and chloroacetyl chloride (2.2 mL, 28.2 mmol) in acetic acid (40 mL). Purified by chromatography in CH<sub>2</sub>Cl<sub>2</sub>, followed by recrystallization from isopropanol to afford 874 mg of a pale yellow solid (28%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 7.55-7.74 (m, 2H); 7.47-7.54 (m, 2H); 7.34 (dd, J=1.7, 7.8 Hz, 1H); 7.13 (dt, J=1.2, 7.7 Hz, 1H); 7.08 (dd, J=1.0, 8.4 Hz, 1H); 4.29 (d, J=12 Hz, 1H); 4.11 (d, J=12 Hz, 1H); 3.80 (s, 3H). MS (ES): m/z 335.0 (M+).

20

**2-Chloromethyl-3-(2-chlorophenyl)-8-trifluoromethyl-3H-quinazolin-4-one (2m)**

25

Prepared according to Procedure B with 1m (500 mg, 1.59 mmol) and chloroacetyl chloride (0.38 mL, 4.77 mmol) in acetic acid (10 mL). Purified by recrystallization from isopropanol to afford 359 mg of a white crystalline solid (61%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.51 (dd, J=1.0, 8.0 Hz, 1H); 8.14 (d, J=7.3 Hz, 1H); 7.65 (dd, J=2.5, 5.6 Hz, 1H); 7.62 (d, J=3.9 Hz, 1H); 7.48-7.60 (m, 3H); 4.44 (d, J=12 Hz, 1H), 4.12 (d, J=12 Hz, 1H). MS (ES): m/z 373.0 (M+).

35

- 126 -

**2-Chloromethyl-3-(2-chlorophenyl)-3H-benzo[g]quinazolin-4-one (2n)**

5                   Prepared according to Procedure B with 1n  
(500 mg, 1.68 mmol) and chloroacetyl chloride (0.40  
mL, 5.05 mmol) in acetic acid (10 mL). Purified by  
chromatography in CH<sub>2</sub>Cl<sub>2</sub> followed by recrystalli-  
zation from isopropanol to afford 232 mg of a light-  
10 brown solid (39%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.92 (s, 1H);  
8.29 (s, 1H); 8.81 (d, J=8.3, 1H); 8.32 (d, J=8.3  
Hz, 1H); 7.51-7.69 (m, 4H); 7.55 (d, J=5.2 Hz, 1H);  
7.53 (d, J=3.8 Hz, 1H); 4.43 (d, J=12 Hz, 1H), 4.12  
(d, J=12 Hz, 1H). MS (ES): m/z 355.0 (M+).

15

**2-Chloromethyl-3-(2-chlorophenyl)-7-nitro-3H-quinazolin-4-one (2o)**

20                   Prepared according to Procedure B with 1o  
(500 mg, 1.71 mmol) and chloroacetyl chloride (0.41  
mL, 5.14 mmol) in acetic acid (10 mL). Purified by  
extraction from aqueous K<sub>2</sub>CO<sub>3</sub>, followed by two  
chromatographies in CH<sub>2</sub>Cl<sub>2</sub> to afford 338 mg of a  
25 yellow oil (56%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.64 (d, J=2.2  
Hz, 1H); 8.48 (d, J=8.8 Hz, 1H); 8.32 (dd, J=2.2,  
8.7 Hz, 1H); 7.66 (dd, J=2.5, 6.0 Hz, 1H); 7.52-7.59  
(m, 3H); 4.41 (d, J=12 Hz, 1H), 4.10 (d, J=12 Hz,  
1H). MS (ES): m/z 350.0 (M+).

30

**Acetic acid 2-chloromethyl-3-(2-chlorophenyl)-4-oxo-3,4-dihydro-quinazolin-6-yl ester (2p)**

35                   Prepared according to Procedure B with 1p  
(670 mg, 2.55 mmol) and chloroacetyl chloride (0.61  
mL, 7.65 mmol) in acetic acid (10 mL). Purified by

- 127 -

chromatography in 0-3% MeOH/CH<sub>2</sub>Cl<sub>2</sub>, followed by  
recrystallization from isopropanol to afford 523 mg  
of the acetate as pale-peach crystals (57%). <sup>1</sup>H NMR  
(CDCl<sub>3</sub>) δ: 8.00 (d, J=2.7 Hz, 1H); 7.82 (d, J=8.8  
5 Hz, 1H); 7.60-7.66 (m, 1H); 7.56 (dd, J=2.7, 8.8 Hz,  
1H); 7.51 (t, J=4.7 Hz, 2H); 7.50 (s, 1H); 4.38 (d,  
J=12 Hz, 1H), 4.08 (d, J=12 Hz, 1H); 2.36 (s, 3H).  
MS (ES): m/z 363.0 (M+).

10 **2-Chloromethyl-3-(2-chlorophenyl)-6,7-difluoro-3H-  
quinazolin-4-one (2q)**

Prepared according to Procedure B with 1q  
15 (700 mg, 2.48 mmol) and chloroacetyl chloride (0.60  
mL, 7.43 mmol) in acetic acid (12 mL). Purified by  
chromatography in CH<sub>2</sub>Cl<sub>2</sub>, followed by recrystalli-  
zation from isopropanol to afford 219 mg of a yellow  
crystalline solid (26%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.07  
20 (dd, J=8.5, 9.7 Hz, 1H); 7.64 (dd, J=2.5, 5.6 Hz,  
1H); 7.60 (dd, J=3.5, 11 Hz, 1H); 7.55 (q, J=2.9 Hz,  
3H); 7.52 (d, J=1.9 Hz, 1H); 7.49-7.51 (m, 1H); 4.36  
(d, J=12 Hz, 1H), 4.06 (d, J=12 Hz, 1H). MS (ES):  
m/z 341.0 (M+).

25

**2-Chloromethyl-3-(2-chlorophenyl)-6-fluoro-3H-  
quinazolin-4-one (2r)**

30 Prepared according to Procedure B with 1r  
(850 mg, 3.21 mmol) and chloroacetyl chloride (0.77  
mL, 9.63 mmol) in acetic acid (15 mL). Purified by  
extraction from aqueous K<sub>2</sub>CO<sub>3</sub>, followed by chromatog-  
raphy in EtOAc/hexanes. A second chromatography in  
35 acetone/hexanes afforded 125 mg of a white solid

- 128 -

(12%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 7.95 (dd, J=2.9, 8.2 Hz, 1H); 7.81 (dd, J=4.8, 9.0 Hz, 1H); 7.61-7.66 (m, 1H); 7.57 (dd, J=2.7, 8.6 Hz, 1H); 7.57 (dd, J=2.7, 8.6 Hz, 1H); 7.52 (dd, J=3.2, 6.9 Hz, 1H); 7.52 (br s, 2H); 4.38 (d, J=12 Hz, 1H), 4.08 (d, J=12 Hz, 1H). MS (ES): m/z 323.0 (M+).

### EXAMPLE 10

#### Preparation of PI3Kδ Inhibitor Compounds

##### Compound D-001

2-(6-Aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-6,7-dimethoxy-3H-quinazolin-4-one

15

Prepared according to Procedure C using Intermediate 2a (200 mg, 0.546 mmol), adenine (81 mg, 0.601 mmol), K<sub>2</sub>CO<sub>3</sub> (83 mg, 0.601 mmol), and DMF (4 mL). The crude product was recrystallized from ethanol (EtOH) to provide 164 mg of a beige solid (65%), mp 281.5-282.7°C (decomposes). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.06 (s, 1H); 8.04 (s, 1H); 7.76-7.81 (m, 1H); 7.70-7.76 (m, 1H); 7.60-7.67 (m, 2H); 7.45 (s, 1H); 7.22 (s, 2H); 6.90 (s, 1H); 5.08 (d, J=17 Hz, 1H); 4.91 (d, J=17 Hz, 1H); 3.87 (s, 3H); 3.87 (s, 3H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 159.9, 156.2, 155.4, 152.9, 150.0, 149.7, 149.4, 143.0, 141.9, 133.7, 132.1, 131.9, 131.2, 130.8, 129.3, 118.4, 113.6, 108.4, 105.8, 56.5, 56.1, 44.7. MS (ES): m/z 464.1 (M+). Anal. calcd. for C<sub>22</sub>H<sub>18</sub>ClN<sub>7</sub>O<sub>3</sub>•0.1C<sub>2</sub>H<sub>6</sub>O•0.05KCl: C, 56.47; H, 3.97; Cl, 7.88; N, 20.76. Found: C, 56.54; H, 4.05; Cl, 7.77; N, 20.55.

- 129 -

## Compound D-002

2-(6-Aminopurin-9-ylmethyl)-6-bromo-3-(2-chlorophenyl)-3H-quinazolin-4-one

5

Prepared according to Procedure C using Intermediate 2b (100 mg, 0.260 mmol), adenine (39 mg, 0.286 mmol), K<sub>2</sub>CO<sub>3</sub> (40 mg, 0.286 mmol), and DMF (2 mL). The crude product was recrystallized from EtOH to provide 52 mg of an off-white solid (41%), mp 284.2-284.7°C (decomposes). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.24 (d, J=2.0 Hz, 1H); 8.05 (s, 1H); 8.03 (s, 1H); 7.98 (dd, J=1.9, 8.6 Hz, 1H); 7.74-7.83 (m, 2H); 7.59-7.68 (m, 2H); 7.46 (d, J=8.7 Hz, 1H); 7.22 (s, 2H); 5.12 (d, J=17 Hz, 1H); 4.94 (d, J=17 Hz, 1H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 159.5, 156.2, 152.9, 152.0, 150.1, 145.8, 141.8, 138.4, 133.1, 132.2, 131.9, 131.1, 130.9, 130.1, 129.4, 128.9, 122.4, 120.4, 118.4, 45.0. MS (ES): m/z 482.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>13</sub>ClBrN<sub>7</sub>O•0.1KCl: C, 49.01; H, 2.67; Cl, 7.96; N, 20.00. Found: C, 48.82; H, 2.82; Cl, 8.00; N, 19.79.

## Compound D-003

2-(6-Aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-7-fluoro-3H-quinazolin-4-one

Prepared according to Procedure C using Intermediate 2c (100 mg, 0.310 mmol), adenine (46 mg, 0.340 mmol), K<sub>2</sub>CO<sub>3</sub> (47 mg, 0.340 mmol), and DMF (1 mL). The crude product was recrystallized from EtOH to provide 57 mg of a beige solid (44%), mp 216.8-217.2°C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.22 (dd, J=6.3, 8.7 Hz, 1H); 8.05 (s, 1H); 8.03 (s, 1H); 7.78-7.80

- 130 -

(m, 2H); 7.61-7.64 (m, 2H); 7.46 (dt, J=2.1, 8.6 Hz, 1H); 7.32 (d, J=9.8 Hz, 1H); 7.22 (s, 2H); 5.13 (d, J=17 Hz, 1H); 4.95 (d, J=17 Hz, 1H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 166.1 (d, J=253 Hz), 159.6, 155.8, 152.5, 149.7, 148.6 (d, J=14 Hz), 141.4, 132.8, 131.8, 131.6, 130.8, 130.5, 129.8 (d, J=11 Hz), 129.0, 118.1, 117.4, 116.2 (d, J=24 Hz), 112.7 (d, J=22 Hz), 44.6. MS (ES): m/z 422.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>13</sub>ClFN<sub>7</sub>O•0.1H<sub>2</sub>O(0.15KCl): C, 55.25; H, 3.06; Cl, 9.38; N, 22.55. Found: C, 55.13; H, 2.92; Cl, 9.12; N, 22.30.

**Compound D-004**

2-(6-Aminopurin-9-ylmethyl)-6-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one

Prepared according to Procedure C using Intermediate 2d (100 mg, 0.294 mmol), adenine (44 mg, 0.323 mmol), K<sub>2</sub>CO<sub>3</sub> (45 mg, 0.323 mmol), and DMF (1 mL). The crude product was recrystallized from EtOH to provide 50 mg of a yellow solid (39%), mp 294.5-294.8°C (decomposes). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.10 (d, J=2.2 Hz, 1H); 8.05 (s, 1H); 8.03 (s, 1H); 7.86 (dd, J=2.4, 8.8 Hz, 1H); 7.75-7.82 (m, 2H); 7.59-7.67 (m, 2H); 7.53 (d, J=8.7 Hz, 1H); 7.22 (br s, 2H); 5.13 (d, J=17 Hz, 1H); 4.95 (d, J=17 Hz, 1H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 159.7, 156.2, 152.9, 151.9, 150.1, 145.5, 141.8, 135.7, 133.1, 132.3, 132.2, 131.9, 131.1, 130.9, 130.0, 129.4, 125.9, 122.0, 118.4, 44.9. MS (ES): m/z 438.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>13</sub>Cl<sub>2</sub>N<sub>7</sub>O: C, 54.81; H, 2.99; N, 22.37. Found: C, 54.72; H, 2.87; N, 22.18.

- 131 -

## Compound D-005

2-(6-Aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-5-fluoro-3H-quinazolin-4-one

5

Prepared according to Procedure C using Intermediate 2e (200 mg, 0.619 mmol), adenine (92 mg, 0.681 mmol), K<sub>2</sub>CO<sub>3</sub> (94 mg, 0.680 mmol), and DMF (4 mL). The crude product was chromatographed in MeOH/CH<sub>2</sub>Cl<sub>2</sub> to provide 168 mg of an off-white solid (64%), mp 159-172°C (gradually decomposes). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.10 (s, 1H); 8.08 (s, 1H); 7.73-7.89 (m, 3H); 7.57-7.71 (m, 2H); 7.37-7.48 (m, 2H); 7.34 (d, J=11 Hz, 1H); 7.30 (d, J=8.3 Hz, 1H); 5.14 (d, J=17 Hz, 1H); 4.94 (d, J=17 Hz, 1H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 160.8 (d, J=264 Hz), 157.5 (d, J=4.2 Hz), 155.8, 152.4, 152.4, 150.0, 148.7, 142.1, 136.4 (d, J=11 Hz), 133.0, 132.2, 132.1, 131.2, 130.9, 129.4, 123.8 (d, J=3.6 Hz), 118.4, 114.5 (d, J=20 Hz), 110.2 (d, J=6.0 Hz), 44.9. MS (ES): m/z 422.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>13</sub>ClFN<sub>7</sub>O: C, 56.95; H, 3.11; Cl, 8.40; N, 23.24. Found: C, 54.62; H, 3.32; Cl, 9.40; N, 21.29.

25

## Compound D-006

2-(6-Aminopurin-9-ylmethyl)-5-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one

30

Prepared according to Procedure C using Intermediate 2f (300 mg, 0.883 mmol), adenine (131 mg, 0.972 mmol), K<sub>2</sub>CO<sub>3</sub> (134 mg, 0.972 mmol), and DMF (4 mL). The crude product was chromatographed in MeOH/CH<sub>2</sub>Cl<sub>2</sub> and recrystallized from EtOH to provide 188 mg of a pale orange crystalline solid (49%), mp

35



- 132 -

245.7-246.0<sup>°</sup> (starts to sweat at 220°C). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.06 (s, 1H); 8.04 (s, 1H); 7.76-7.81 (m, 2H); 7.72 (d, J=8.0 Hz, 1H); 7.59-7.66 (m, 3H); 7.41 (d, J=8.1 Hz, 1H); 7.26 (br s, 2H); 5.11 (d, J=17 Hz, 1H); 4.93 (d, J=17 Hz, 1H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 158.5, 156.2, 152.9, 152.2, 150.1, 149.2, 141.8, 135.4, 133.3, 133.2, 132.1, 132.0, 131.2, 130.9, 130.4, 129.4, 127.3, 118.4, 117.7, 44.9. MS (ES): m/z 438.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>13</sub>Cl<sub>2</sub>N<sub>7</sub>O•0.1C<sub>2</sub>H<sub>6</sub>O•0.05H<sub>2</sub>O: C, 54.67; H, 3.11; Cl, 15.98; N, 22.09. Found: C, 54.35; H, 3.00; Cl, 15.82; N, 22.31.

**Compound D-007**

2-(6-Aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-5-methyl-3H-quinazolin-4-one

Prepared according to Procedure C using Intermediate 2g (250 mg, 0.783 mmol), adenine (116 mg, 0.862 mmol), K<sub>2</sub>CO<sub>3</sub> (119 mg, 0.862 mmol), and DMF (4 mL). The crude product was recrystallized from EtOH to provide 93 mg of a pale yellow solid (28%), mp 190.7-190.9°C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.05 (s, 1H); 8.03 (s, 1H); 7.76-7.79 (m, 1H); 7.71-7.74 (m, 1H); 7.59-7.67 (m, 1H); 7.34 (d, J=7.4 Hz, 1H); 7.28 (d, J=8.2 Hz, 1H); 7.24 (br s, 2H); 5.07 (d, J=17 Hz, 1H); 4.92 (d, J=17 Hz, 1H); 2.73 (s, 3H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 161.1, 156.2, 152.8, 150.9, 150.1, 148.3, 141.9, 141.0, 134.6, 133.6, 132.2, 131.9, 131.3, 130.8, 130.3, 129.3, 125.9, 119.1, 118.4, 44.8, 22.8. MS (ES): m/z 418.1 (M<sup>+</sup>). Anal. calcd. for C<sub>21</sub>H<sub>16</sub>ClN<sub>7</sub>O•H<sub>2</sub>O: C, 57.87; H, 4.16; Cl, 8.13; N,

- 133 -

22.49. Found: C, 57.78; H, 3.99; Cl, 8.38; N,  
22.32.

**Compound D-008**

5     **2-(6-Aminopurin-9-ylmethyl)-8-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one**

Prepared according to Procedure C using  
10     Intermediate 2h (100 mg, 0.294 mmol), adenine (44 mg, 0.324 mmol), K<sub>2</sub>CO<sub>3</sub> (45 mg, 0.324 mmol), and DMF (1 mL). The crude product was chromatographed in MeOH/CH<sub>2</sub>Cl<sub>2</sub> to provide 50 mg of a pale yellow solid (39%), mp 273.3-273.5°C (discolors). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.11 (dd, J=1.3, 8.0 Hz, 1H); 8.08 (s, 1H);  
15     8.05 (s, 1H); 8.00 (dd, J=1.3, 7.8 Hz, 1H); 7.79-7.83 (m, 2H); 7.63-7.66 (m, 2H); 7.56 (t, J=7.9 Hz, 1H); 7.21 (br s, 2H); 5.17 (d, J=17 Hz, 1H); 4.97 (d, J=17 Hz, 1H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 160.2,  
20     156.1, 152.8, 152.2, 150.2, 143.3, 142.0, 135.6, 133.1, 132.3, 131.9, 131.1, 131.0, 130.9, 129.4, 128.4, 126.0, 122.5, 118.4, 45.0. MS (ES): m/z 438.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>13</sub>Cl<sub>2</sub>N<sub>7</sub>O•0.1CH<sub>4</sub>O•0.6H<sub>2</sub>O(0.15KCl): C, 52.09; H,  
25     3.18; N, 21.15. Found: C, 51.85; H, 2.93; N, 21.01.

**Compound D-009**

30     **2-(6-Aminopurin-9-ylmethyl)-3-biphenyl-2-yl-5-chloro-3H-quinazolin-4-one**

Prepared according to Procedure C using  
Intermediate 2i (400 mg, 1.05 mmol), adenine (155 mg, 1.15 mmol), K<sub>2</sub>CO<sub>3</sub> (159 mg, 1.15 mmol), and DMF (5  
35     mL). The crude product was recrystallized from EtOH

- 134 -

to provide 344 mg of a white solid (68%), mp 299.9-300.1°C (discolors). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.08 (s, 1H); 7.89 (s, 1H); 7.58-7.73 (m, 5H); 7.51 (d, J=7.9 Hz, 1H); 7.46 (d, J=7.5 Hz, 2H); 7.27-7.41 (m, 3H);  
5 7.14-7.27 (m, 3H); 5.14 (d, J=17 Hz, 1H); 4.82 (d, J=17 Hz, 1H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 159.6, 156.2, 152.8, 152.5, 150.0, 149.0, 141.7, 140.2, 137.7, 135.0, 133.3, 133.2, 131.8, 130.7, 130.1, 129.8, 129.5, 128.8, 128.6, 128.4, 127.1, 118.4, 117.6,  
10 45.3. MS (ES): m/z 480.1 (M<sup>+</sup>). Anal. calcd. for C<sub>26</sub>H<sub>18</sub>ClN<sub>7</sub>O: C, 65.07; H, 3.78; Cl, 7.39; N, 20.43. Found: C, 64.77; H, 3.75; Cl, 7.43; N, 20.35.

**Compound D-010**

15 **5-Chloro-2-(9H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one**

Prepared according to Procedure C using  
20 Intermediate 2j (200 mg, 0.626 mmol), 6-mercaptopurine monohydrate (93 mg, 0.546 mmol), K<sub>2</sub>CO<sub>3</sub> (95 mg, 0.689 mmol), and DMF (4 mL). The crude product was recrystallized from EtOH to provide 125 mg of an off-white solid (46%), mp 213.9°C. <sup>1</sup>H NMR  
25 (DMSO-d<sub>6</sub>) δ: 13.53 (br s, 1H); 8.49 (s, 1H); 8.44 (s, 1H); 7.78 (t, J=7.9 Hz, 1H); 7.63 (d, J=8.2 Hz, 1H); 7.59 (d, J=7.7 Hz, 1H); 7.49 (d, J=6.9 Hz, 1H); 7.24-7.41 (m, 3H); 4.32-4.45 (m, 2H); 2.14 (s, 3H).  
<sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 158.9, 157.2, 154.2, 151.5,  
30 149.7, 149.6, 143.5, 136.1, 135.9, 135.1, 133.2, 131.3, 130.3, 130.0, 129.9, 129.1, 127.6, 127.1, 117.8, 32.4, 17.5. MS (ES): m/z 438.0 (M<sup>+</sup>). Anal. calcd. for C<sub>21</sub>H<sub>15</sub>ClN<sub>6</sub>OS: C, 58.00; H, 3.48; Cl, 8.15;

- 135 -

N, 19.32; S, 7.37. Found: C, 58.05; H, 3.38; Cl, 8.89; N, 18.38; S, 7.00.

**Compound D-011**

5 **5-Chloro-3-(2-fluorophenyl)-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one**

Prepared according to Procedure C using  
10 Intermediate 2k (210 mg, 0.650 mmol), 6-mercapto-purine monohydrate (122 mg, 0.715 mmol), K<sub>2</sub>CO<sub>3</sub> (99 mg, 0.715 mmol), and DMF (4 mL). The crude product was recrystallized from EtOH to provide 240 mg of an off-white solid (84%), mp 244.0°C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  
15 δ: 13.56 (br s, 1H); 8.50 (s, 1H); 8.45 (s, 1H); 7.81 (t, J=8.0 Hz, 1H); 7.74 (t, J=7.7 Hz, 1H); 7.67 (d, J=8.1 Hz, 1H); 7.62 (d, J=7.7 Hz, 1H); 7.46-7.55 (m, 1H); 7.29-7.42 (m, 2H); 4.47-4.59 (m, 2H). <sup>13</sup>C  
NMR (DMSO-d<sub>6</sub>) ppm: 158.4, 157.3 (d, J=249 Hz),  
20 156.4, 153.8, 151.0, 149.1, 143.2, 135.0, 132.9, 131.8 (d, J=8.0 Hz), 130.8, 129.9, 126.7, 125.3 (d, J=3.5 Hz), 123.6 (d, J=13 Hz), 117.0, 116.2 (d, J=19 Hz), 31.7. MS (ES): m/z 439.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>12</sub>ClFN<sub>6</sub>OS: C, 54.74; H, 2.76; Cl, 8.08; N,  
25 19.15; S, 7.31. Found: C, 54.42; H, 2.88; Cl, 8.08; N, 18.87; S, 7.08.

- 136 -

## Compound D-012

2-(6-Aminopurin-9-ylmethyl)-5-chloro-3-(2-fluorophenyl)-3H-quinazolin-4-one

5

Prepared according to Procedure C using Intermediate 2k (210 mg, 0.650 mmol), adenine (97 mg, 0.715 mmol), K<sub>2</sub>CO<sub>3</sub> (99 mg, 0.715 mmol), and DMF (4 mL). The crude product was recrystallized from EtOH to provide 137 mg of a tan solid (50%), mp 295.6-295.8°C (decomposes). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.05 (s, 1H); 8.04 (s, 1H); 7.75 (t, J=7.6 Hz, 1H); 7.74 (t, J=7.9 Hz, 1H); 7.62-7.69 (m, 1H); 7.61 (d, J=7.6 Hz, 1H); 7.47-7.55 (m, 1H); 7.48 (d, J=7.8 Hz, 1H); 7.41 (d, J=8.0 Hz, 1H); 7.24 (br s, 2H); 5.19 (d, J=17 Hz, 1H); 5.03 (d, J=17 Hz, 1H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 158.7, 157.6 (d, J=250 Hz), 156.2, 152.8, 152.4, 150.0, 149.2, 141.8, 135.4, 133.3, 132.5 (d, J=8.0 Hz), 131.0, 130.4, 127.3, 126.2 (d, J=3.5 Hz), 123.1 (d, J=14 Hz), 118.4, 117.6, 117.2 (d, J=19 Hz), 45.1. MS (ES): m/z 422.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>13</sub>ClFN<sub>7</sub>O•0.05C<sub>2</sub>H<sub>6</sub>O: C, 56.92; H, 3.16; Cl, 8.36; N, 23.12. Found: C, 56.79; H, 3.20; Cl, 8.46; N, 22.79.

25

## Compound D-013

3-Biphenyl-2-yl-5-chloro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one

30

Prepared according to Procedure C using Intermediate 2i (400 mg, 1.05 mmol), 6-mercaptopurine monohydrate (196 mg, 1.15 mmol), K<sub>2</sub>CO<sub>3</sub> (159 mg, 1.15 mmol), and DMF (5 mL). The crude product was chromatographed in MeOH/CH<sub>2</sub>Cl<sub>2</sub> and subsequently

35

- 137 -

recrystallized from EtOH to provide 439 mg of a pale yellow crystalline solid (84%), mp 222.0-222.5°C (dec). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.56 (br s, 1H); 8.55 (s, 1H); 8.45 (s, 1H); 7.73 (t, J=8.0 Hz, 1H); 7.64 (d, J=7.7 Hz, 1H); 7.50-7.59 (m, 4H); 7.41-7.48 (m, 1H); 7.25-7.38 (m, 5H); 4.41 (d, J=16 Hz, 1H); 4.16 (d, J=16 Hz, 1H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 160.2, 157.0, 153.7, 151.5, 149.7, 149.3, 143.5, 139.9, 137.8, 135.1, 134.1, 133.3, 131.5, 130.5, 130.3, 130.1, 129.1, 128.9, 128.4, 128.4, 126.9, 117.5, 32.3. MS (ES): m/z 497.0 (M<sup>+</sup>). Anal. calcd. for C<sub>26</sub>H<sub>17</sub>ClN<sub>6</sub>OS: C, 62.84; H, 3.45; Cl, 7.13; N, 16.91; S, 6.45. Found: C, 62.60; H, 3.47; Cl, 7.15; N, 16.65; S, 6.41.

#### Compound D-014

5-Chloro-3-(2-methoxyphenyl)-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one

Prepared according to Procedure C using Intermediate 21 (250 mg, 0.746 mmol), 6-mercaptopurine monohydrate (140 mg, 0.821 mmol), K<sub>2</sub>CO<sub>3</sub> (113 mg, 0.821 mmol), and DMF (4 mL). The crude product was recrystallized from EtOH to provide 254 mg of an off-white solid (76%), mp 237.0°C (dec; discolors at 154.6°C). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.53 (br s, 1H); 8.52 (s, 1H); 8.45 (s, 1H); 7.78 (t, J=7.9 Hz, 1H); 7.64 (d, J=8.0 Hz, 1H); 7.59 (d, J=7.7 Hz, 1H); 7.48 (d, J=7.3 Hz, 1H); 7.42 (t, J=7.7 Hz, 1H); 7.15 (d, J=8.2 Hz, 1H); 7.03 (t, J=7.5 Hz, 1H); 4.45 (s, 2H); 3.76 (s, 3H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 158.9, 157.1, 154.8, 154.7, 151.5, 149.6, 143.6, 135.1, 133.2, 131.3, 130.4, 130.0, 127.0, 124.8, 121.2, 117.8,

- 138 -

112.7, 56.1, 32.0. MS (ES): m/z 451.0 (M+). Anal. calcd. for  $C_{21}H_{15}ClN_6O_2S \cdot 0.15C_2H_6O \cdot 0.05KCl$ : C, 55.43; H, 3.47; Cl, 8.07; N, 18.21; S, 6.95. Found: C, 55.49; H, 3.68; Cl, 7.95; N, 17.82; S, 6.82.

5

**Compound D-015**

**3-(2-Chlorophenyl)-5-fluoro-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one**

10

Prepared according to Procedure C using Intermediate 2e (200 mg, 0.619 mmol), 6-mercaptopurine monohydrate (116 mg, 0.681 mmol),  $K_2CO_3$  (94 mg, 0.681 mmol), and DMF (5 mL). The crude product was recrystallized from EtOH to provide 152 mg of a white solid (56%), mp 222.7-223.8°C (discolors).  $^1H$  NMR (DMSO- $d_6$ )  $\delta$ : 13.56 (br s, 1H); 8.48 (s, 1H); 8.44 (s, 1H); 7.89 (dt, J=5.6, 8.1 Hz, 1H); 7.76 (dd, J=1.6, 7.3 Hz, 1H); 7.67 (d, J=7.4 Hz, 1H); 7.56 (d, J=8.1 Hz, 1H); 7.47 (t, J=7.1 Hz, 1H), 7.41-7.53 (m, 2H); 7.37 (dd, J=8.7, 11 Hz, 1H); 4.38-4.52 (m, 2H).  $^{13}C$  NMR (DMSO- $d_6$ ) ppm: 160.9 (d, J=264 Hz), 157.6, 156.8, 154.1, 151.5, 149.6, 149.0, 143.6, 136.4 (d, J=11 Hz), 133.9, 132.2, 131.7, 131.6, 130.5, 130.2, 128.8, 123.6, 114.4 (d, J=20 Hz), 110.2, 32.0. MS (ES): m/z 439.0 (M+). Anal. calcd. for  $C_{20}H_{12}ClFN_6OS \cdot 0.5C_2H_6O$ : C, 54.61; H, 3.27; Cl, 7.68; N, 18.19; S, 6.94. Found: C, 54.37; H, 3.26; Cl, 7.89; N, 18.26; S, 6.55.

30

- 139 -

## Compound D-016

3-(2-Chlorophenyl)-6,7-dimethoxy-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one

5 Prepared according to Procedure C using Intermediate 2a (200 mg, 0.546 mmol), 6-mercapto-purine monohydrate (102 mg, 0.601 mmol), K<sub>2</sub>CO<sub>3</sub> (83 mg, 0.601 mmol), and DMF (5 mL). The crude product was recrystallized from EtOH to provide 172 mg of an off-white solid (65%), mp 160-180°C (gradually decomposes). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.55 (br s, 1H); 8.49 (s, 1H); 8.44 (s, 1H); 7.72 (d, J=6.9 Hz, 1H); 7.66 (d, J=6.9 Hz, 1H) 7.38-7.54 (m, 3H); 7.22 (s, 1H); 4.36-4.52 (m, 2H); 3.94 (s, 3H); 3.89 (s, 3H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 160.1, 155.4, 151.5, 151.1, 149.4, 143.2, 134.6, 132.3, 131.6, 131.5, 130.4, 128.7, 113.6, 108.4, 105.8, 56.5, 56.1, 32.0. MS (ES): m/z 481.1 (M<sup>+</sup>). Anal. calcd. for C<sub>22</sub>H<sub>17</sub>ClN<sub>6</sub>O<sub>3</sub>S•0.5C<sub>2</sub>H<sub>6</sub>O•0.05KCl: C, 54.41; H, 3.97; Cl, 7.33; N, 16.55; S, 6.32. Found: C, 54.43; H, 3.94; Cl, 7.69; N, 16.69; S, 6.52.

## Compound D-017

25 6-Bromo-3-(2-chlorophenyl)-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one

30 Prepared according to Procedure C using Intermediate 2b (200 mg, 0.519 mmol), 6-mercapto-purine monohydrate (97 mg, 0.570 mmol), K<sub>2</sub>CO<sub>3</sub> (79 mg, 0.572 mmol), and DMF (5 mL). The crude product was recrystallized from EtOH to provide 123 mg of an off-white solid (47%), mp 212-242°C (gradually decomposes). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.07 (br s, 1H);

35



- 140 -

8.48 (s, 1H); 8.44 (s, 1H); 8.24 (d, J=2.3 Hz, 1H);  
8.06 (dd, J=2.3, 8.7 Hz, 1H); 7.76 (dd, J=1.9, 7.4  
Hz, 1H); 7.70 (d, J=8.7 Hz, 1H); 7.66 (d, J=8.1 Hz,  
1H); 7.51 (dd, J=2.1, 7.9 Hz, 1H); 7.46 (dd, J=1.9,  
5 7.9 Hz, 1H); 4.47 (s, 2H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm:  
159.7, 156.8, 153.6, 151.5, 146.1, 143.6, 138.5,  
134.0, 132.1, 131.8, 131.5, 130.5, 130.2, 129.9,  
128.9, 128.8, 122.2, 120.3, 32.0. MS (ES): m/z  
499.0 (M<sup>+</sup>). Anal. calcd. for  
10 C<sub>20</sub>H<sub>12</sub>ClBrN<sub>6</sub>OS•0.2C<sub>2</sub>H<sub>6</sub>O•0.05KCl: C, 47.79; H, 2.59; N,  
16.39; S, 6.25. Found: C, 47.56; H, 2.54; N, 16.25;  
S, 6.58.

## Compound D-018

15 3-(2-Chlorophenyl)-(9H-purin-6-ylsulfanylmethyl)-  
trifluoromethyl-3H-quinazolin-4-one

Prepared according to Procedure C using  
20 Intermediate 2m (200 mg, 0.536 mmol), 6-mercapto-  
purine monohydrate (100 mg, 0.588 mmol), K<sub>2</sub>CO<sub>3</sub> (82  
mg, 0.593 mmol), and DMF (4 mL). The crude product  
was recrystallized from EtOH to provide 148 mg of a  
white solid (56%), mp 218.5-219.4°C. <sup>1</sup>H NMR (DMSO-  
25 d<sub>6</sub>) δ: 13.52 (br s, 1H); 8.48 (s, 1H); 8.44 (s,  
1H); 8.43 (d, J=6.0 Hz, 1H); 8.26 (d, J=7.5 Hz, 1H);  
7.84 (dd, J=2.5, 6.7 Hz, 1H); 7.70-7.75 (m, 2H);  
7.51-7.59 (m, 2H); 4.40-4.55 (m, 2H). <sup>13</sup>C NMR (DMSO-  
d<sub>6</sub>) ppm: 160.0, 157.2, 154.2, 151.4, 149.6, 144.4,  
30 143.4, 133.8, 133.0 (q, J=5.1 Hz), 132.0, 131.9,  
131.6, 131.4, 130.6, 129.0, 127.3, 125.2 (q, J=30  
Hz), 123.6 (q, J=273 Hz), 121.8, 32.6. MS (ES): m/z  
489.0 (M<sup>+</sup>). Anal. calcd. for C<sub>21</sub>H<sub>12</sub>ClF<sub>3</sub>N<sub>6</sub>OS: C,

- 141 -

51.59; H, 2.47; Cl, 7.25; N, 17.19; S, 6.56. Found: C, 51.51; H, 2.55; Cl, 7.37; N, 17.05; S, 6.38.

**Compound D-019**

5     **3-(2-Chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-benzo[g]quinazolin-4-one**

Prepared according to Procedure C using  
10     Intermediate 2n (200 mg, 0.563 mmol), 6-mercapto-  
purine monohydrate (105 mg, 0.619 mmol), K<sub>2</sub>CO<sub>3</sub> (86  
mg, 0.619 mmol), and DMF (4 mL). The crude product  
was recrystallized from EtOH to provide 128 mg of a  
dark yellow solid (48%), mp 247.8-254.4°C (decom-  
15     poses). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.56 (br s, 1H); 8.90  
(s, 1H); 8.50 (s, 1H); 8.46 (s, 1H); 8.34 (s, 1H);  
8.27 (d, J=8.2 Hz, 1H); 8.16 (d, J=8.2 Hz, 1H); 7.81  
(dd, J=1.6, 7.3 Hz, 1H); 7.70 (t, J=7.5 Hz, 1H);  
7.61-7.74 (m, 2H); 7.49 (t, J=7.5 Hz, 1H); 7.44-7.53  
20     (m, 1H); 4.44-4.56 (m, 2H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm:  
161.3, 151.6, 151.5, 143.9, 142.2, 136.7, 134.4,  
132.5, 131.8, 131.6, 130.5, 129.7, 129.3, 128.8,  
128.6, 128.3, 128.3, 127.1, 125.2, 119.5, 32.4. MS  
(ES): m/z 471.0 (M<sup>+</sup>). Anal. calcd. for  
25     C<sub>24</sub>H<sub>15</sub>ClN<sub>6</sub>OS•0.2C<sub>2</sub>H<sub>6</sub>O•0.05KCl: C, 60.57; H, 3.37; Cl,  
7.69; N, 17.37; S, 6.63. Found: C, 60.24; H, 3.46;  
Cl, 7.50; N, 17.34; S, 6.69.

- 142 -

## Compound D-020

6-Chloro-3-(2-chlorophenyl)-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one

5

Prepared according to Procedure C using Intermediate 2d (200 mg, 0.587 mmol), 6-mercaptopurine monohydrate (110 mg, 0.646 mmol),  $K_2CO_3$  (90 mg, 0.651 mmol), and DMF (5 mL). The crude product was recrystallized from EtOH to provide 113 mg of a yellow crystalline solid (42%), mp 237.1-238.2°C (decomposes).  $^1H$  NMR (DMSO- $d_6$ )  $\delta$ : 13.55 (br s, 1H); 8.48 (s, 1H); 8.44 (s, 1H); 8.11 (s, 1H); 7.94 (d, J=8.3 Hz, 1H); 7.78 (d, J=8.1 Hz, 2H); 7.66 (d, J=6.7 Hz, 1H); 7.48-7.56 (m, 2H); 4.48 (s, 2H).  $^{13}C$  NMR (DMSO- $d_6$ ) ppm: 159.8, 156.8, 153.5, 151.5, 149.6, 145.8, 143.6, 135.7, 134.0, 132.2, 132.1, 131.7, 131.5, 130.5, 130.2, 129.8, 128.8, 125.8, 121.9, 32.0. MS (ES): m/z 455.0 (M+). Anal. calcd. for  $C_{20}H_{12}Cl_2N_6OS \cdot 0.1C_2H_6O \cdot 0.6H_2O (0.15KCl)$ : C, 50.34; H, 2.89; Cl, 15.82; N, 17.44; S, 6.65. Found: C, 50.02; H, 2.63; Cl, 15.51; N, 17.39; S, 6.81.

## Compound D-021

8-Chloro-3-(2-chlorophenyl)-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one

Prepared according to Procedure C using Intermediate 2h (200 mg, 0.589 mmol), 6-mercaptopurine monohydrate (124 mg, 0.726 mmol),  $K_2CO_3$  (100 mg, 0.726 mmol), and DMF (4 mL). The crude product was recrystallized from EtOH to provide 202 mg of a white solid (75%), mp 211.9-212.7°C (decomposes).  $^1H$  NMR (DMSO- $d_6$ )  $\delta$ : 13.54 (br s, 1H); 8.47 (s, 1H);

- 143 -

8.44 (s, 1H); 8.12 (d, J=7.9 Hz, 1H); 8.07 (d, J=7.6 Hz, 1H); 7.78 (d, J=7.5 Hz, 1H); 7.67 (d, J=7.1 Hz, 1H); 7.58 (t, J=7.9 Hz, 1H); 7.42-7.54 (m, 2H); 4.52 (s, 2H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 160.3, 156.9, 5 153.9, 151.5, 149.7, 143.5, 135.7, 134.0, 132.1, 131.8, 131.4, 131.1, 130.5, 130.3, 128.9, 128.3, 126.1, 122.4, 32.5. MS (ES): m/z 455.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>12</sub>Cl<sub>2</sub>N<sub>6</sub>OS: C, 52.76; H, 2.66; Cl, 15.57; N, 18.46; S, 7.04. Found: C, 52.65; H, 2.79; 10 Cl, 15.32; N, 18.47; S, 7.18.

**Compound D-022**

3-(2-Chlorophenyl)-7-fluoro-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one 15

Prepared according to Procedure C using Intermediate 2c (200 mg, 0.619 mmol), 6-mercapto-purine monohydrate (116 mg, 0.681 mmol), K<sub>2</sub>CO<sub>3</sub> (95 20 mg, 0.687 mmol), and DMF (4 mL). The crude product was recrystallized from EtOH to provide 143 mg of a white crystalline solid (53%), mp 151.4-154.2°C (discolors). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.55 (br s, 1H); 8.48 (s, 1H); 8.44 (s, 1H); 8.23 (dd, J=6.3, 8.7 Hz, 1H); 7.77 (dd, J=1.7, 7.4 Hz, 1H); 7.64 (d, J=7.4 25 Hz, 1H); 7.57 (d, J=9.8 Hz, 1H); 7.45-7.52 (m, 3H); 4.48 (s, 2H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 169.0 (d, J=253 Hz), 162.6, 159.3, 157.0, 154.0, 152.2, 151.7 (d, J=13 Hz), 146.1, 136.5, 134.7, 134.2, 134.0, 30 133.0, 132.6 (d, J=11 Hz), 131.3, 120.2, 118.9 (d, J=24 Hz), 115.3 (d, J=22 Hz), 34.6. MS (ES): m/z 439.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>12</sub>ClFN<sub>6</sub>OS•0.4-C<sub>2</sub>H<sub>6</sub>O•0.4H<sub>2</sub>O(0.15KCl: C, 52.52; H, 3.22; Cl, 8.57;

- 144 -

N, 17.67. Found: C, 52.25; H, 3.11; Cl, 8.20; N, 17.69.

**Compound D-023**

5     **3-(2-Chlorophenyl)-7-nitro-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one**

Prepared according to Procedure C using  
10     Intermediate 2o (216 mg, 0.617 mmol), 6-mercapto-purine monohydrate (116 mg, 0.681 mmol), K<sub>2</sub>CO<sub>3</sub> (94 mg, 0.680 mmol), and DMF (4 mL). The crude product was recrystallized from EtOH to provide 212 mg of a yellow crystalline solid (74%), mp 218.0-218.3°C  
15     (decomposes). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.56 (br s, 1H); 8.49 (s, 1H); 8.42 (s, 1H); 8.38-8.45 (m, 2H); 8.31 (d, J=8.4 Hz, 1H); 7.81 (d, J=6.5 Hz, 1H); 7.68 (d, J=6.7 Hz, 1H); 7.43-7.58 (m, 2H); 4.53 (s, 2H). <sup>13</sup>C  
NMR (DMSO-d<sub>6</sub>) ppm: 157.7, 154.4, 153.3, 149.8,  
20     149.3, 147.6, 145.2, 141.4, 131.5, 129.8, 129.7, 129.2, 128.4, 127.1, 126.7, 122.7, 120.3, 119.4, 29.9. MS (ES): m/z 466.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>12</sub>ClN<sub>7</sub>O<sub>3</sub>S•0.4C<sub>2</sub>H<sub>6</sub>O•0.05KCl: C, 51.19; H, 2.97; Cl, 7.63; N, 20.09; S, 6.57. Found: C, 51.27; H, 2.88;  
25     Cl, 7.40; N, 20.04; S, 6.52.

**Compound D-024**

30     **3-(2-Chlorophenyl)-6-hydroxy-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one**

Prepared according to Procedure C using  
Intermediate 2p (200 mg, 0.552 mmol), 6-mercapto-purine monohydrate (117 mg, 0.685 mmol), K<sub>2</sub>CO<sub>3</sub> (95  
35     mg, 0.687 mmol), and DMF (4 mL). The crude product

- 145 -

was recrystallized from EtOH to provide 182 mg of a white solid, a mixture of the desired product and the acetyl derivative. A portion of this material (120 mg) was suspended in a mixture of MeOH (2 mL) and aqueous NaHCO<sub>3</sub> (satd., 1 mL) and stirred rapidly for 4 hours. The mixture was concentrated *in vacuo*, suspended in H<sub>2</sub>O (10 mL), and stored at 4°C overnight. The white solid was collected and dried to 103 mg (66%), mp 186-214°C (gradually decomposes).  
<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.48 (s, 1H); 8.45 (s, 1H); 7.71 (d, J=6.8 Hz, 1H); 7.62-7.64 (m, 2H); 7.43-7.51 (m, 2H); 7.40-7.43 (m, 1H); 7.35 (d, J=8.8 Hz, 1H); 4.39-4.52 (m, 2H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 160.6, 157.1, 156.2, 151.4, 150.8, 149.3, 144.1, 140.2, 134.5, 132.2, 131.6, 131.4, 130.4, 129.3, 128.7, 124.8, 121.7, 109.8, 32.0. MS (ES): m/z 437.0 (M<sup>+</sup>).  
Anal. calcd. for (2 C<sub>20</sub>H<sub>13</sub>ClN<sub>6</sub>O<sub>2</sub>S•0.1C<sub>2</sub>H<sub>6</sub>O•0.6H<sub>2</sub>O: C, 49.68; H, 3.88; Cl, 7.26; N, 17.21; S, 6.57. Found: C, 49.43; H, 3.62; Cl, 7.32; N, 17.07; S, 6.58.

## Compound D-025

5-Chloro-3-(2-chlorophenyl)-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one

Prepared according to Procedure C using Intermediate 2f (300 mg, 0.883 mmol), 6-mercaptopurine monohydrate (165 mg, 0.972 mmol), K<sub>2</sub>CO<sub>3</sub> (134 mg, 0.972 mmol), and DMF (4 mL). The crude product was recrystallized from EtOH to provide 341 mg of a pale orange crystalline solid (85%), mp 233.7-234.4°C (decomposes). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.58 (br s, 1H); 8.50 (s, 1H); 8.47 (s, 1H); 7.77-7.85 (m, 2H); 7.68 (d, J=8.1 Hz, 2H); 7.65 (d, J=7.7 Hz, 1H);

- 146 -

7.41-7.56 (m, 2H); 4.45 (d, J=1.2 Hz, 2H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 158.7, 156.8, 153.8, 151.5, 149.6, 149.5, 143.5, 135.4, 134.1, 133.3, 132.2, 131.6, 131.6, 130.5, 130.2, 128.8, 127.1, 117.6, 32.0. MS (ES): m/z 455.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>12</sub>Cl<sub>2</sub>N<sub>6</sub>OS•C<sub>2</sub>H<sub>6</sub>O•0.3H<sub>2</sub>: C, 52.14; H, 3.70; Cl, 13.99; N, 16.58; S, 6.33. Found: C, 52.07; H, 3.37; Cl, 13.40; N, 16.65; S, 6.42.

10      **Compound D-026**

**3-(2-Chlorophenyl)-5-methyl-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one**

15                      Prepared according to Procedure C using Intermediate 2g (300 mg, 0.940 mmol), 6-mercapto-purine monohydrate (176 mg, 1.03 mmol), K<sub>2</sub>CO<sub>3</sub> (142 mg, 1.03 mmol), and DMF (5 mL). The crude product was recrystallized from EtOH to provide 324 mg of a  
20      white crystalline solid (79%), mp 227.8-230.1°C (decomposes). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.57 (br s, 1H); 8.49 (s, 1H); 8.47 (s, 1H); 7.69-7.78 (m, 2H); 7.66 (d, J=7.3 Hz, 1H); 7.55 (d, J=7.9 Hz, 1H); 7.39-7.52 (m, 2H); 7.36 (d, J=6.9 Hz, 1H); 4.38-4.50 (m, 2H);  
25      2.74 (s, 3H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 161.2, 156.3, 152.4, 151.5, 148.6, 143.9, 141.0, 134.6, 134.5, 132.3, 131.7, 131.4, 130.4, 130.2, 128.7, 125.7, 119.0, 32.0, 22.8. MS (ES): m/z 435.0 (M<sup>+</sup>). Anal. calcd. for C<sub>21</sub>H<sub>15</sub>ClN<sub>6</sub>OS•0.65C<sub>2</sub>H<sub>6</sub>O•0.1H<sub>2</sub>O: C, 57.40; H, 4.13; Cl, 7.60; N, 18.01; S, 6.87. Found: C, 57.11;  
30      H, 3.96; Cl, 7.45; N, 17.79; S, 6.90.

- 147 -

**Compound D-027****3-(2-Chlorophenyl)-6,7-difluoro-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one**

5

Prepared according to Procedure C using Intermediate 2q (200 mg, 0.586 mmol), 6-mercapto-purine monohydrate (110 mg, 0.645 mmol), K<sub>2</sub>CO<sub>3</sub> (89 mg, 0.645 mmol), and DMF (4 mL). The crude product was recrystallized from EtOH to provide 143 mg of a pale yellow crystalline solid (53%), mp 207.8°C (discolors; sweats at 136°C). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.57 (br s, 1H); 8.49 (s, 1H); 8.46 (s, 1H); 8.11 (t, J=9.4 Hz, 1H); 7.88 (dd, J=7.3, 11 Hz, 1H); 7.77 (dd, J=1.7, 7.3 Hz, 1H); 7.67 (d, J=7.4 Hz, 1H); 7.42-7.55 (m, 2H); 4.48 (s, 2H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 159.5 (d, J=2.5 Hz), 154.6 (dd, J=14, 255 Hz), 154.0 (d, J=1.5 Hz), 151.5, 149.3 (dd, J=14, 250 Hz), 145.1 (d, J=12 Hz), 143.9, 133.9, 132.1, 131.8, 131.4, 130.5, 128.9, 118.0 (d, J=4.9 Hz), 115.8 (d, J=18 Hz), 114.6 (d, J=20 Hz), 32.0. MS (ES): m/z 457.0 (M<sup>+</sup>). Anal. calcd. for C<sub>20</sub>H<sub>11</sub>ClF<sub>2</sub>N<sub>6</sub>OS: C, 52.58; H, 2.43; Cl, 7.76; N, 18.40; S, 7.02. Found: C, 51.81; H, 2.37; Cl, 7.49; N, 18.04; S, 7.55.

25

**Compound D-028****3-(2-Chlorophenyl)-6-fluoro-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one**

30

Prepared according to Procedure C using Intermediate 2r (118 mg, 0.365 mmol), 6-mercapto-purine monohydrate (68 mg, 0.402 mmol), K<sub>2</sub>CO<sub>3</sub> (56 mg, 0.402 mmol), and DMF (2 mL). The crude product was recrystallized from EtOH to provide 103 mg of an

35



- 148 -

off-white crystalline solid (64%), mp 232.8-233.0°C (discolors). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.56 (br s, 1H); 8.48 (s, 1H); 8.44 (s, 1H); 7.81-7.86 (m, 3H); 7.76 (d, J=7.5 Hz, 1H); 7.67 (d, J=7.5 Hz, 1H); 7.40-7.54 (m, 2H); 4.48 (br s, 2H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 160.8 (d, J=247 Hz), 160.2 (d, J=3.3 Hz), 156.9, 152.3 (d, J=1.9 Hz), 151.5, 149.7, 144.0, 143.6, 134.1, 132.1, 131.7, 131.5, 130.5, 130.4, 130.2, 128.8, 124.0 (d, J=24 Hz), 122.0 (d, J=8.7 Hz), 111.7 (d, J=24 Hz), 32.0. MS (ES): m/z 439.0 (M<sup>+</sup>).  
Anal. calcd. for C<sub>20</sub>H<sub>12</sub>ClFN<sub>6</sub>OS•0.2C<sub>2</sub>H<sub>6</sub>O•0.1H<sub>2</sub>O: C, 54.46; H, 3.00; Cl, 7.88; N, 18.68. Found: C, 54.09; H, 2.73; Cl, 7.80; N, 18.77.

Compound D-029

2-(6-Aminopurin-9-ylmethyl)-3-(2-isopropylphenyl)-5-methyl-3H-quinazolin-4-one

Thionyl chloride (2.2 mL, 30 mmol) was added to a stirred solution of 2-amino-6-methylbenzoic acid (1.51 g, 10 mmol) in benzene (50 mL) and the mixture was heated at reflux for 18 h. Once cooled, the solvent was removed *in vacuo* and stripped down twice with benzene (25 mL). The residue was dissolved in CHCl<sub>3</sub> (50 mL) and treated with 2-isopropylaniline (2.83 mL, 20 mmol). The slurry was then heated at reflux for 3 h. At that time TLC (50% EtOAc/hexane) indicated that the reaction was complete. After cooling to room temperature, the reaction mixture was poured atop a 4 cm plug of silica gel and flushed through with 20% EtOAc/hexane. The product containing fractions were combined and concentrated *in vacuo*. The residue was

- 149 -

dissolved in HOAc (50 mL) and treated with chloro-  
actyl chloride (1.6 mL, 20 mmol) and the mixture was  
heated at reflux for 18 h. The reaction was cooled  
and concentrated in vacuo. The remaining HOAc was  
5 removed by azeotroping with toluene (25 mL) three  
times. The residue was dissolved in toluene (10 mL)  
and poured through a 4 cm plug of silica gel, flush-  
ing through with 20 % EtOAc/hexane. The product  
containing fractions were identified by LCMS (MS  
10 (ES): m/z 327 (M+)), and concentrated in vacuo to  
afford 975 mg (30%) as a white foam. The white foam  
chloride (450 mg, 1.36 mmol) was dissolved in DMF  
(10 mL) and treated with adenine (275 mg, 2.04 mmol)  
and K<sub>2</sub>CO<sub>3</sub> (281 mg, 2.04 mmol) and the mixture was  
15 stirred overnight at room temperature. The suspen-  
sion was then poured into 200 mL of water, stirred  
at room temperature for 30 min then chilled in the  
refrigerator for 30 min. The resultant solid was  
collected by vacuum filtration and recrystallized  
20 from EtOH to afford 285 mg (49%) of an off white  
solid. mp 258.0-258.2°C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.19  
(s, 1H), 8.09 (s, 1H), 7.60 (m, 3H), 7.45 (m, 2H),  
7.23 (m, 3H), 5.11 (d, J=17.5 Hz, 1H), 4.71 (d,  
J=17.5 Hz, 1H), 2.68 (s, 3H), 2.73 (q, J=6.9 Hz,  
25 1H), 1.34 (d, J=6.8 Hz, 3H), 1.13 (d, J=6.8 Hz, 3H).  
<sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 161.9, 156.2, 152.8, 151.6,  
150.1, 148.4, 146.1, 142.2, 140.8, 134.3, 133.7,  
130.6, 130.0, 129.0, 127.7, 127.6, 125.8, 119.2,  
118.4, 44.8, 28.3, 24.4, 23.3, 22.9. MS (ES): m/z  
30 426.4 (M+). Anal. calcd. for C<sub>24</sub>H<sub>23</sub>N<sub>7</sub>O: C, 67.75; H,  
5.45; N, 23.04. Found: C, 67.60; H, 5.45; N, 22.82.

- 150 -

## Compound D-030

2-(6-Aminopurin-9-ylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one

5                   Thionyl chloride (2.2 mL, 30 mmol) was added to a stirred solution of 2-amino-6-methylbenzoic acid (1.51 g, 10 mmol) in benzene (50 mL) and the mixture was heated at reflux for 18 h. Once  
10 cooled, the solvent was removed *in vacuo* and stripped down twice with benzene (25 mL). The residue was dissolved in CHCl<sub>3</sub> (50 mL) and treated with o-toluidine (2.13 mL, 20 mmol). The slurry was then heated at reflux for 3 h. At that time TLC  
15 (50% EtOAc/hexane) indicated that the reaction was complete. After cooling to room temperature, the reaction mixture was poured atop a 4 cm plug of silica gel and flushed through with 20% EtOAc/hexane. The product containing fractions were combined  
20 and concentrated *in vacuo*. The residue was dissolved in HOAc (50 mL) and treated with chloroacetyl chloride (1.6 mL, 20 mmol) and the mixture was heated at reflux for 18 h. The reaction was cooled and concentrated *in vacuo*. The remaining HOAc was  
25 removed by azeotroping with toluene (25 mL) three times. The residue was dissolved in toluene (10 mL) and poured through a 4 cm plug of silica gel, flushing through with 20 % EtOAc/hexane. The product containing fractions were identified by LCMS  
30 [MS (ES): m/z 299 (M<sup>+</sup>)], and concentrated *in vacuo* to afford 476 mg (16%) as a white foam. The white foam chloride (470 mg, 1.57 mmol) was dissolved in DMF (10 mL) and treated with adenine (423 mg, 3.14 mmol) and K<sub>2</sub>CO<sub>3</sub> (433 mg, 3.14 mmol) and the mixture

- 151 -

was stirred overnight at room temperature. The suspension was then poured into 200 mL of H<sub>2</sub>O, stirred at room temperature for 30 min then chilled in the refrigerator for 30 min. The resultant solid was collected by vacuum filtration and recrystallized from EtOH to afford 123 mg (20%) of an off white solid. mp 281.5-282.7°C (decomposes). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.07 (s, 1H); 8.05 (s, 1H); 7.61 (t, J=7.8 Hz, 1H), 7.48 (m, 4H), 7.25 (m, 3H), 5.09 (d, J=17.4 Hz, 1H), 4.76 (d, J=17.4 Hz, 1H), 2.73 (s, 3H), 2.18 (s, 3H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 161.3, 156.2, 152.8, 151.4, 150.0, 148.5, 142.2, 140.9, 136.1, 135.4, 134.3, 131.7, 130.1, 130.0, 129.0, 128.0, 125.8, 119.2, 118.5, 44.8, 22.9, 17.4. MS (ES): m/z 398.2 (M<sup>+</sup>). Anal. calcd. for C<sub>22</sub>H<sub>19</sub>N<sub>7</sub>O: C, 66.49; H, 4.82; N, 24.67. Found: C, 66.29; H, 4.78; N, 24.72.

**Compound D-031**

**3-(2-Fluorophenyl)-5-methyl-2-(9H-purin-6-yl-sulfanylmethyl)-3H-quinazolin-4-one**

Thionyl chloride (2.2 mL, 30 mmol) was added to a stirred solution of 2-amino-6-methylbenzoic acid (1.51 g, 10 mmol) in benzene (50 mL) and the mixture was heated at reflux for 18 h. Once cooled, the solvent was removed in vacuo and stripped down twice with benzene (25 mL). The residue was dissolved in CHCl<sub>3</sub> (50 mL) and treated with 2-fluoroaniline (1.93 mL, 20 mmol). The slurry was then heated at reflux for 3 h. At that time TLC (50% EtOAc/hexane) indicated that the reaction was complete. After cooling to room temperature, the

- 152 -

reaction mixture was poured atop a 4 cm plug of silica gel and flushed through with 20% EtOAc/hexane. The product containing fractions were combined and concentrated in vacuo. The residue was dissolved in HOAc (50 mL) and treated with chloroacetyl chloride (1.6 mL, 20 mmol) and the mixture was heated at reflux for 18 h. The reaction was cooled and concentrated in vacuo. The remaining HOAc was removed by azeotroping with toluene (25 mL) three times. The residue was dissolved in toluene (10 mL) and poured through a 4 cm plug of silica gel, flushing through with 20 % EtOAc/hexane. The product containing fractions were identified by LCMS [MS (ES): m/z 303 (M+)], and concentrated in vacuo to afford 1.12 g (37%) as a white foam. The white foam chloride (455 mg, 1.50 mmol) was dissolved in DMF (10 mL) and treated with 6-mercaptopurine monohydrate (510 mg, 3.0 mmol) and K<sub>2</sub>CO<sub>3</sub> (414 mg, 3.0 mmol) and the mixture was stirred overnight at room temperature. The suspension was then poured into 200 mL of water, stirred at room temperature for 30 min then chilled in the refrigerator for 30 min. The resultant solid was collected by vacuum filtration and recrystallized from EtOH to afford 487 mg (77%) of an off white solid. mp 151.9-152.2°C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.48 (s, 1H), 8.44 (s, 1H), 7.70 (m, 2H), 7.48 (m, 2H), 7.33 (m, 3H), 4.55 (d, J=15.1 Hz, 1H), 4.48 (d, J=15.1 Hz, 1H), 2.73 (s, 3H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) ppm: 161.3, 157.8 (d, J=249.1 Hz), 156.9, 152.8, 151.5, 149.6, 148.6, 143.6, 140.9, 134.7, 131.9 (d, J=8.0 Hz), 131.4, 130.2, , 125.6 (d, J=3.6 Hz), 125.5, 124.4 (d, J=13.5 Hz), 118.8, 116.6 (d, J=19.6 Hz), 56.4, 22.9. MS (ES): m/z

- 153 -

419.5 (M+). Anal. calcd. for  $C_{21}H_{15}FN_6OS \cdot 0.15 C_2H_6O$ :  
C, 60.14; H, 3.77; F, 4.47; N, 19.76; S, 7.54.  
Found: C, 59.89; H, 3.88; F, 4.42; N, 19.42; S, 7.23.

5      **Compound D-032**

**2-(6-Aminopurin-9-ylmethyl)-5-chloro-3-o-tolyl-3H-quinazolin-4-one**

10                      Prepared according to Procedure C using 2j  
(200 mg, 0.626 mmol), adenine (93 mg, 0.689 mmol),  
 $K_2CO_3$  (95 mg, 0.689 mmol), and DMF (3 mL). The crude  
product was chromatographed in MeOH/ $CH_2Cl_2$  to provide  
101 mg of an off-white solid (39%), mp 262.0-  
15      266.5°C.  $^1H$  NMR (DMSO- $d_6$ )  $\delta$ : 8.08 (s, 1H); 8.07 (s,  
1H); 7.70 (t,  $J=8.0$  Hz, 1H); 7.58 (dd,  $J=0.6$ , 7.9  
Hz, 1H); 7.43-7.57 (m, 4H); 7.36 (dd,  $J=0.7$ , 8.0 Hz,  
1H); 7.26 (br s, 2H); 5.12 (d,  $J=18$  Hz, 1H); 4.78  
(d,  $J=18$  Hz, 1H); 2.20 (s, 3H).  $^{13}C$  NMR (DMSO- $d_6$ )  
20      ppm: 158.7, 156.2, 152.9, 152.7, 150.0, 149.4,  
142.1, 136.1, 135.1, 135.0, 133.2, 131.8, 130.3,  
130.1, 128.9, 128.1, 127.2, 118.5, 117.9, 44.9,  
17.4. MS (ES):  $m/z$  418.1 (M+). Anal. calcd. for  
 $C_{21}H_{16}ClN_7O \cdot 0.1H_2O \cdot 0.05KCl$ : C, 59.57; H, 3.86; Cl,  
25      8.79; N, 23.16. Found: C, 59.65; H, 3.80; Cl, 8.70;  
N, 22.80.

- 154 -

## Compound D-033

2-(6-Aminopurin-9-ylmethyl)-5-chloro-3-(2-methoxy-phenyl)-3H-quinazolin-4-one

5

Prepared according to Procedure C using 21  
(250 mg, 0.746 mmol), adenine (111 mg, 0.821 mmol),  
K<sub>2</sub>CO<sub>3</sub> (113 mg, 0.821 mmol), and DMF (4 mL). The  
crude product was chromatographed in MeOH/CH<sub>2</sub>Cl<sub>2</sub> and  
10 recrystallized from EtOH to provide 124 mg of a  
brown solid (38%), mp 257.0-257.1°C. <sup>1</sup>H NMR (DMSO-  
d<sub>6</sub>) δ: 8.06 (s, 1H); 8.01 (s, 1H); 7.71 (t, J=8.0  
Hz, 1H); 7.57 (dd, J=0.9, 7.9 Hz, 1H); 7.52-7.59 (m,  
1H); 7.50 (dd, J=1.6, 7.8 Hz, 1H); 7.38 (dd, J=1.1,  
15 8.2 Hz, 1H); 7.27 (dd, J=0.6, 8.3 Hz, 1H); 7.24 (br  
s, 2H); 7.17 (dt, J=0.9, 7.6 Hz, 1H); 5.07 (d, J=17  
Hz, 1H); 4.97 (d, J=17 Hz, 1H); 3.79 (s, 3H). <sup>13</sup>C  
NMR (DMSO-d<sub>6</sub>) ppm: 158.8, 156.2, 154.7, 153.2,  
152.8, 150.1, 149.3, 142.0, 135.1, 133.2, 131.8,  
20 130.1, 130.1, 127.2, 123.8, 121.6, 118.4, 117.9,  
113.1, 56.2, 44.8. MS (ES): m/z 434.0 (M<sup>+</sup>). Anal.  
calcd. for C<sub>21</sub>H<sub>16</sub>ClN<sub>7</sub>O<sub>2</sub>•0.5H<sub>2</sub>O•0.04KCl: C, 56.57; H,  
3.84; Cl, 8.27; N, 21.99. Found: C, 56.29; H, 3.75;  
Cl, 8.21; N, 21.61.

25

The following compounds were made gener-  
ally in accordance with the above-described methods  
and serve to further illustrate specific embodiments  
of the compounds of the invention:

30

3-(2,6-dichlorophenyl)-5-methyl-2-(9H-purin-6-ylsul-  
fanylmethyl)-3H-quinazolin-4-one (D-034)  
3-(2-isopropylphenyl)-5-methyl-2-(9H-purin-6-ylsul-  
fanylmethyl)-3H-quinazolin-4-one (D-035)

- 155 -

- 3-(2-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-quinazolin-4-one (D-036)
- 3-benzyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quin-  
azolin-4-one (D-037)
- 5 3-butyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quin-  
azolin-4-one (D-038)
- 3-morpholin-4-yl-2-(9H-purin-6-ylsulfanylmethyl)-3H-  
quinazolin-4-one, acetate salt (D-039)
- 3-(3-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-  
10 3H-quinazolin-4-one (D-040)
- 3-(3-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-quinazolin-4-one (D-041)
- 2-(9H-purin-6-ylsulfanylmethyl)-3-pyridin-4-yl-3H-  
quinazolin-4-one (D-042)
- 15 3-benzyl-5-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-quinazolin-4-one (D-043)
- 3-(4-methylpiperazin-1-yl)-2-(9H-purin-6-ylsulfan-  
ylmethyl)-3H-quinazolin-4-one, acetate salt (D-044)
- [5-fluoro-4-oxo-2-(9H-purin-6-ylsulfanylmethyl)-4H-  
20 quinazolin-3-yl]acetic acid ethyl ester (D-045)
- 3-(2-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-quinazolin-4-one (D-046)
- 3-(2-methoxyphenyl)-5-methyl-2-(9H-purin-6-ylsulfan-  
ylmethyl)-3H-quinazolin-4-one (D-047)
- 25 2-(6-aminopurin-9-ylmethyl)-3-(2-fluorophenyl)-5-  
methyl-3H-quinazolin-4-one (D-048)
- 2-(6-aminopurin-9-ylmethyl)-3-benzyl-5-fluoro-3H-  
quinazolin-4-one (D-049)
- 2-(6-aminopurin-9-ylmethyl)-3-butyl-3H-quinazolin-4-  
30 one (D-050)
- 2-(6-aminopurin-9-ylmethyl)-3-morpholin-4-yl-3H-  
quinazolin-4-one, acetate salt (D-051)




- 156 -

3-(4-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-quinazolin-4-one (D-052).

Additional compounds of the present inven-  
tion were prepared by the following synthetic pro-  
cedures.

The following intermediates were prepared  
by the above-described Procedure A.




- 3a R=cyclopropyl
- 3b R=cyclopropylmethyl
- 3c R=phenethyl
- 3d R=cyclopentyl
- 3e R=3-(2-chloro)pyridyl
- 3f R=4-(2-methyl)benzoic acid
- 3g R=4-nitrobenzyl
- 3h R=cyclohexyl
- 3i R=E-(2-phenyl)cyclopropyl

Additional compounds of the present inven-  
tion (D-053 through D-070) having the following core  
structure are discussed in the following Experimen-  
tal Section. All were prepared following Procedure  
C.

- 157 -


Core structure:



Compound No.	R	R'
D-053	cyclopropyl	C
D-054	cyclopropylmethyl	B
D-055	cyclopropylmethyl	A
D-056	cyclopropylmethyl	C
D-057	phenethyl	B
D-058	phenethyl	C
D-059	cyclopentyl	B
D-060	cyclopentyl	A
D-061	3-(2-chloro)pyridyl	B
D-062	3-(2-chloro)pyridyl	A
D-063	4-(2-methyl)benzoic acid	B
D-064	cyclopropyl	B
D-065	cyclopropyl	A
D-066	4-nitrobenzyl	B
D-067	cyclohexyl	B
D-068	cyclohexyl	A
D-069	cyclohexyl	C
D-070	E-(2-phenyl)cyclopropyl	B


- 158 -

5




A

10



B

20



25

C

- 159 -

2-(2-Amino-9H-purin-6-ylsulfanylmethyl)-3-cyclopropyl-5-methyl-3H-quinazolin-4-one (D-053)

5                   Prepared according to procedure C using 3a  
(100mg, 0.4 mmol), 2-amino-6-mercaptopurine (80 mg,  
0.48 mmol), and K<sub>2</sub>CO<sub>3</sub> (77 mg, 0.56 mmol). The  
product was purified by trituration from H<sub>2</sub>O. <sup>1</sup>H NMR  
(DMSO-d<sub>6</sub>) δ: 7.89 (d, J=0.9 Hz, 1H); 7.54 (t, J=7.4  
10 Hz, 1H); 7.34 (d, J=8.1 Hz, 1H); 7.19 (d, J=7.2 Hz,  
1H); 6.28 (s, 2H); 4.94 (s, 2H); 2.70 (s, 3H); 1.24  
(d, J=6.5 Hz, 2H); 0.91 (s, 2H). MS (ES): m/z 380  
(M+H), 190.

15                   3-Cyclopropylmethyl-5-methyl-2-(9H-purin-6-  
ylsulfanylmethyl)-3H-quinazolin-4-one (D-054)

                  Prepared according to procedure C using 3b  
20 (300mg, 1.14 mmol), 6-mercaptopurine monohydrate  
(214 mg, 1.26 mmol), and K<sub>2</sub>CO<sub>3</sub> (189 mg, 1.37 mmol).  
The product was purified by trituration from H<sub>2</sub>O,  
followed by recrystallization from MeOH. <sup>1</sup>H NMR  
(DMSO-d<sub>6</sub>) δ: 13.60 (br s, 1H); 8.72 (s, 1H); 8.48  
25 (s, 1H); 7.63 (t, J=7.8 Hz, 1H); 7.42 (d, J=8.0 Hz,  
1H); 7.28 (d, J=7.3 Hz, 1H); 5.01 (s, 2H); 4.11 (d,  
J=6.8 Hz, 2H); 2.78 (s, 3H); 1.35 (quint, J=6.2 Hz,  
1H); 0.44-0.59 (m, 4H). MS (ES): m/z 379 (M+H),  
325.

30

- 160 -

2-(6-Aminopurin-9-ylmethyl)-3-cyclopropylmethyl-5-methyl-3H-quinazolin-4-one (D-055)

5                   Prepared according to procedure C using 3b  
(300mg, 1.14 mmol), adenine (170 mg, 1.26 mmol), and  
K<sub>2</sub>CO<sub>3</sub> (189 mg, 1.37 mmol). The product was purified  
by trituration from H<sub>2</sub>O, followed by recrystalli-  
zation from MeOH. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.21 (s, 1H);  
10 8.10 (s, 1H); 7.52 (t, J=7.7 Hz, 1H); 7.18-7.31 (m,  
3H); 7.06 (d, J=8.1 Hz, 1H); 5.68 (s, 2H); 4.14 (d,  
J=6.8 Hz, 2H); 2.77 (s, 3H); 1.34 (quint, J=6.4 Hz,  
1H); 0.45-0.60 (m, 4H). MS (ES): m/z 362 (M+H),  
308.

15

2-(2-Amino-9H-purin-6-ylsulfanylmethyl)-3-cyclo-  
propylmethyl-5-methyl-3H-quinazolin-4-one (D-056)

20                   Prepared according to procedure C using 3b  
(280mg, 1.1 mmol), 2-amino-6-mercaptapurine (200 mg,  
1.2 mmol), and K<sub>2</sub>CO<sub>3</sub> (180 mg, 1.3 mmol). The product  
was purified by trituration from MeOH. <sup>1</sup>H NMR (DMSO-  
d<sub>6</sub>) δ: 12.70 (br s, 1H); 7.95 (s, 1H); 7.64 (t,  
25 J=7.8 Hz, 1H); 7.44 (d, J=7.9 Hz, 1H); 7.28 (d,  
J=7.4 Hz, 1H); 6.41 (s, 2H); 4.91 (s, 2H); 4.05 (d,  
J=6.8 Hz, 2H); 2.78 (s, 3H); 1.26-1.43 (m, 1H);  
0.36-0.56 (m, 4H). MS (ES): m/z 394 (M+H), 340.

30 5-Methyl-3-phenethyl-2-(9H-purin-6-ylsulfanyl-  
methyl)-3H-quinazolin-4-one (D-057)

                  Prepared according to procedure C using 3c  
35 (750mg, 2.4 mmol), 6-mercaptapurine monohydrate (442  
mg, 2.6 mmol), and K<sub>2</sub>CO<sub>3</sub> (398 mg, 2.9 mmol). The  
product was purified by trituration from H<sub>2</sub>O. <sup>1</sup>H NMR

- 161 -

(DMSO- $d_6$ )  $\delta$ : 13.61 (s, 1H); 8.71 (s, 1H); 8.48 (s, 1H); 7.65 (t,  $J=7.7$  Hz, 1H); 7.44 (d,  $J=7.9$  Hz, 1H); 7.16-7.35 (m, 6H); 4.89 (s, 2H); 4.29 (br t,  $J=7.9$  Hz, 2H); 3.08 (br t,  $J=7.8$  Hz, 2H); 2.81 (s, 3H).

5 MS (ES):  $m/z$  429 (M+H), 105.

**2-(2-Amino-9H-purin-6-ylsulfanylmethyl)-5-methyl-3-phenethyl-3H-quinazolin-4-one (D-058)**

10

Prepared according to procedure C using 3c (750mg, 2.4 mmol), 2-amino-6-mercaptapurine (435 mg, 2.6 mmol), and  $K_2CO_3$  (398 mg, 2.9 mmol). The product was purified by trituration from  $H_2O$ .  $^1H$  NMR (DMSO- $d_6$ )  $\delta$ : 12.61 (s, 1H); 7.95 (s, 1H); 7.65 (t,  $J=7.7$  Hz, 1H); 7.45 (d,  $J=7.9$  Hz, 1H); 7.14-7.32 (m, 6H); 6.44 (s, 2H); 4.81 (s, 2H); 4.24 (br t,  $J=7.9$  Hz, 2H); 3.04 (br t,  $J=7.8$  Hz, 2H); 2.81 (s, 3H). MS (ES):  $m/z$  444 (M+H), 340.

15

20

**3-Cyclopentyl-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one (D-059)**

25

Prepared according to procedure C using 3d (100mg, 0.36 mmol), 6-mercaptapurine monohydrate (73 mg, 0.43 mmol), and  $K_2CO_3$  (100 mg, 0.72 mmol). The product was purified by recrystallization from MeOH.  $^1H$  NMR (DMSO- $d_6$ )  $\delta$ : 13.62 (br s, 1H); 8.77 (s, 1H); 8.48 (s, 1H); 7.62 (t,  $J=7.7$  Hz, 1H); 7.42 (d,  $J=7.8$  Hz, 2H); 7.26 (d,  $J=7.4$  Hz, 1H); 5.03 (s, 2H); 4.80 (quint,  $J=8.0$  Hz, 1H); 2.76 (s, 3H); 2.12-2.31 (m, 2H); 1.79-2.04 (m, 4H); 1.44-1.58 (m, 2H). MS (ES):  $m/z$  393 (M+H), 325.

30

35

- 162 -

2-(6-Aminopurin-9-ylmethyl)-3-cyclopentyl-5-methyl-3H-quinazolin-4-one (D-060)

5                   Prepared according to procedure C using 3d  
(100mg, 0.36 mmol), adenine (58 mg, 0.43 mmol), and  
K<sub>2</sub>CO<sub>3</sub> (100 mg, 0.72 mmol). The product was purified  
by recrystallization from MeOH. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ:  
8.15 (s, 1H); 8.11 (s, 1H); 7.52 (t, J=7.7 Hz, 1H);  
10 7.16-7.31 (m, 3H); 7.10 (d, J=8.0 Hz, 2H); 5.68 (s,  
2H); 4.78 (quint, J=8.3 Hz, 1H); 2.74 (s, 3H); 2.09-  
2.32 (m, 2H); 1.86-2.04 (m, 2H); 1.68-1.86 (m, 2H);  
1.43-1.67 (m, 2H). MS (ES): m/z 376 (M+H), 308,  
154.

15

3-(2-Chloro-pyridin-3-yl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one (D-061)

20

Prepared according to procedure C using 3e  
(500mg, 1.6 mmol), 6-mercaptapurine monohydrate (289  
mg, 1.7 mmol), and K<sub>2</sub>CO<sub>3</sub> (262 mg, 1.9 mmol). The  
product was purified by trituration from H<sub>2</sub>O. MS  
25 (ES): m/z 436 (M+H), 200.

2-(6-Aminopurin-9-ylmethyl)-3-(2-chloro-pyridin-3-yl)-5-methyl-3H-quinazolin-4-one (D-062)

30

Prepared according to procedure C using 3e  
(500mg, 1.6 mmol), adenine (230 mg, 1.7 mmol), and  
K<sub>2</sub>CO<sub>3</sub> (262 mg, 1.9 mmol). The product was purified  
by trituration from H<sub>2</sub>O. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.59  
35 (dd, J=1.7, 4.8 Hz, 1H); 8.22 (dd, J=1.7, 7.8 Hz,  
1H); 8.025 (s, 1H); 8.017 (s, 1H); 7.60-7.72 (m,  
2H); 7.35 (t, J=8.2 Hz, 2H); 7.22 (s, 2H); 5.12 (d,

- 163 -

J=17.0 Hz, 1H); 5.02 (d, J=17.0 Hz, 1H); 2.72 (s, 3H). MS (ES): m/z 419 (M+H).

5      **3-Methyl-4-[5-methyl-4-oxo-2-(9H-purin-6-ylsulfanylmethyl)-4H-quinazolin-3-yl]-benzoic acid (D-063)**

Prepared according to procedure C using 3f (400mg, 1.17 mmol), 6-mercaptopurine monohydrate (219 mg, 1.29 mmol), and K<sub>2</sub>CO<sub>3</sub> (226 mg, 1.64 mmol). The product was purified by recrystallization from MeOH. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 13.54 (br s, 1H); 8.44 (s, 1H); 8.42 (s, 1H); 7.80 (s, 2H); 7.71 (t, J=7.7 Hz, 1H); 7.59 (d, J=8.6 Hz, 1H); 7.52 (d, J=7.9 Hz, 1H); 7.34 (d, J=7.4 Hz, 1H); 4.46 (d, J=15.4 Hz, 1H); 4.34 (d, J=15.7 Hz, 1H); 3.17 (d, J=4.4 Hz, 1H); 2.73 (s, 3H); 2.17 (s, 3H). MS (ES): m/z 459 (M+H).

20      **3-Cyclopropyl-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one (D-064)**

Prepared according to procedure C using 3a (100mg, 0.40 mmol), 6-mercaptopurine monohydrate (90 mg, 0.53 mmol), and K<sub>2</sub>CO<sub>3</sub> (97 mg, 0.7 mmol). The product was purified by trituration from H<sub>2</sub>O. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.69 (d, J=0.8 Hz, 1H); 8.47 (s, 1H); 7.57 (d, J=7.9 Hz, 1H); 7.37 (d, J=8.1 Hz, 1H); 7.23 (d, J=7.3 Hz, 1H); 5.08 (s, 2H); 3.06-3.18 (m, 1H); 2.74 (s, 3H); 1.14-1.36 (m, 2H); 0.92-1.06 (m, 2H).



- 164 -

2-(6-Aminopurin-9-ylmethyl)-3-cyclopropyl-5-methyl-3H-quinazolin-4-one (D-065)

5                   Prepared according to procedure C using 3a  
(100mg, 0.40 mmol), adenine (94 mg, 0.7 mmol), and  
K<sub>2</sub>CO<sub>3</sub> (121 mg, 0.88 mmol). The product was purified  
by trituration from H<sub>2</sub>O. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.19  
(d, J=0.9 Hz, 1H); 8.09 (d, J=1.0 Hz, 1H); 7.48 (t,  
10 J=7.8 Hz, 1H); 7.13-7.29 (m, 3H); 7.04 (d, J=8.1 Hz,  
1H); 5.74 (s, 2H); 3.00-3.13 (m, 1H); 2.73 (s, 3H);  
1.18-1.38 (m, 2H); 0.94-1.09 (m, 2H).

15                   5-Methyl-3-(4-nitro-benzyl)-2-(9H-purin-6-ylsul-  
fanylmethyl)-3H-quinazolin-4-one (D-066)

                  Prepared according to procedure C using 3g  
(200mg, 0.58 mmol), 6-mercaptapurine monohydrate  
20 (148 mg, 0.87 mmol), and K<sub>2</sub>CO<sub>3</sub> (160 mg, 1.16 mmol).  
The product was purified by trituration from MeOH. <sup>1</sup>H  
NMR (DMSO-d<sub>6</sub>) δ: 13.44 (br s, 1H); 8.50 (s, 1H);  
8.31 (s, 1H); 8.03 (d, J=8.6 Hz, 2H); 7.58 (t, J=7.9  
Hz, 1H); 7.37 (d, J=8.3 Hz, 3H); 7.22 (d, J=7.5 Hz,  
25 1H); 5.44 (s, 2H); 4.70 (s, 2H); 2.66 (s, 3H). MS  
(ES): m/z 460 (M+H).

30                   3-Cyclohexyl-5-methyl-2-(9H-purin-6-ylsulfanyl-  
methyl)-3H-quinazolin-4-one (D-067)

                  Prepared according to procedure C using 3h  
(150mg, 0.52 mmol), 6-mercaptapurine monohydrate (97  
mg, 0.57 mmol), and K<sub>2</sub>CO<sub>3</sub> (86 mg, 0.62 mmol). The  
35 product was purified by trituration from MeOH. <sup>1</sup>H NMR  
(DMSO-d<sub>6</sub>) δ: 13.66 (br s, 1H); 8.82 (s, 1H); 8.50  
(s, 1H); 7.62 (t, J=7.7 Hz, 1H); 7.42 (d, J=8.0 Hz,

- 165 -

1H); 7.26 (d, J=7.3 Hz, 1H); 5.01 (s, 2H); 4.11 (br s, 1H); 2.75 (s, 3H); 2.38-2.65 (m, 2H); 1.58-1.90 (m, 4H); 1.37-1.57 (m, 1H); 0.71-1.26 (m, 3H). MS (ES): m/z 407 (M+H), 325.

5

2-(6-Aminopurin-9-ylmethyl)-3-cyclohexyl-5-methyl-3H-quinazolin-4-one (D-068)

10 Prepared according to procedure C using 3h (150mg, 0.52 mmol), adenine (77 mg, 0.57 mmol), and K<sub>2</sub>CO<sub>3</sub> (86 mg, 0.62 mmol). The product was purified by trituration from MeOH. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.15 (s, 2H); 7.54 (t, J=7.9 Hz, 1H); 7.06-7.35 (m, 4H);  
15 5.65 (s, 2H); 4.09 (br s, 1H); 2.73 (s, 3H); 1.41-1.90 (m, 6H); 0.99-1.34 (m, 4H). MS (ES): m/z 390 (M+H), 308.

20 2-(2-Amino-9H-purin-6-ylsulfanylmethyl)-3-cyclohexyl-5-methyl-3H-quinazolin-4-one (D-069)

Prepared according to procedure C using 3h (150mg, 0.52 mmol), 2-amino-6-mercaptapurine (95 mg, 0.57 mmol), and K<sub>2</sub>CO<sub>3</sub> (86 mg, 0.62 mmol). The  
25 product was purified by reversed-phase HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220 ). MS (ES): m/z 422 (M+H),  
30 340, 170.

5-Methyl-3-(E-2-phenyl-cyclopropyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one (D-070)

35

Prepared according to procedure C using 3i and 6-mercaptapurine monohydrate). The product was

- 166 -

purified by reversed-phase HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220 ). MS (ES): m/z 441.


5

The following Methods 1 and 2 were used as the HPLC analyses for the following compounds:

HPLC Method 1. Column: 2 x 50 mm C18  
10 Luna column (from Phenomenex), flow rate: 0.3 mL/min, UV detection at 214 and 254 nm. Initial conditions: 2% solvent B in solvent A; t=3 min, 20% Solvent B; t=6 min, 80% Solvent B, where Solvent A=water with 0.05% formic acid and Solvent  
15 B=acetonitrile with 0.05% formic acid.


HPLC Method 2. Column: 2 x 50 mm C18  
Luna column (from Phenomenex), flow rate: 0.3 mL/min, UV detection at 214 and 254 nm. Initial  
20 conditions: 10-100% solvent B in solvent A over 6 min, where Solvent A=water and Solvent B=acetonitrile.

Compounds D-070A and D-070B were prepared  
25 as follows:




30

- 167 -



5


Intermediate (1s)



10

15

Intermediate (2s)




20

25

D-070A

H<sub>2</sub>N

- 168 -



2-(6-Aminopurin-9-ylmethyl)-3-(2-benzyloxyphenyl)-5-methyl-3H-quinazolin-4-one (D-070A)


Using procedure A, 6-methylantranilic acid and 2-benzyloxyaniline were converted to Intermediate (1s). Using procedure B, Intermediate (1s) was converted to Intermediate (2s). Using procedure C, Intermediate (2s) was converted to D-070A. Retention time using HPLC Method 2: 4.7 min. LRMS (ES pos.) m/z = 490 (M+1).

2-(6-Aminopurin-9-ylmethyl)-3-(2-hydroxyphenyl)-5-methyl-3H-quinazolin-4-one (D-070B)

A mixture of D-070A (35 mg, 0.07 mmol) and Pd(OH)<sub>2</sub> (20% by wt. on C, 50 mg) in ethanol (5 mL) was shaken 24 hours under 40 psi of hydrogen gas. The catalyst was removed by filtration through a 0.22 um cellulose acetate membrane (Corning), and the filtrate was concentrated *in vacuo* to afford the solid product (10 mg). Retention time using HPLC Method 2: 3.6 min. LRMS (ES pos.) m/z = 400 (M+1).

- 169 -


Compounds D-070C through D-070F were prepared according to the following scheme.



- 170 -

2-(6-Aminopurin-9-ylmethyl)-5-methyl-3-{2-(2-(1-methylpyrrolidin-2-yl)-ethoxy)-phenyl}-3H-quinazolin-4-one (D-070C)

5



10


A mixture of D-070B (30 mg, 0.075 mmol), 2-(2-chloroethyl)-1-methylpyrrolidine hydrochloride (28 mg, 0.15 mmol), and potassium carbonate (50 mg, 0.36 mmol) in DMF (0.3 mL) was stirred at 80°C for 16 hours. The solvent was removed *in vacuo*, then the residue was dissolved in DMSO/water (1 mL) and purified by HPLC in two portions (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 2-50% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, 0.05% TFA in all solvents, detector at 220). Appropriate fractions were concentrated *in vacuo*, then concentrated twice from 1 N HCl (1 mL) to yield the final product as the HCl salt (10 mg). Retention time using HPLC method 1: 4.7 min. LRMS (ES pos.)  $m/z=511$  (M+1).

25

- 171 -

2-(6-Aminopurin-9-ylmethyl)-3-(2-(3-dimethylamino-propoxy)-phenyl)-5-methyl-3H-quinazolin-4-one  
(D-070D)

5



10

A mixture of D-070B (30 mg, 0.075 mmol), 2-chloroethyl)-1-methylpyrrolidine hydrochloride (28 mg, 0.15 mmol), and potassium carbonate (50 mg, 0.36 mmol) in DMF (0.3 mL) was stirred at 80°C for 16 h. The solvent was removed in vacuo, then the residue was dissolved in DMSO/water (1 mL) and purified by HPLC in two portions (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 2-50% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, 0.05% TFA in all solvents, detector at 2201). Appropriate fractions were concentrated in vacuo, then concentrated twice from 1 N HCl (1 mL) to yield the final product as the HCl salt (14 mg). Retention time using HPLC Method 1: 4.5 min. LRMS (ES pos.) m/z=485 (M+1).

25

2-(6-Aminopurin-9-ylmethyl)-5-methyl-3-(2-prop-2-ynyloxyphenyl)-3H-quinazolin-4-one (D-070E)

30



35


A mixture of D-070B (20 mg, 0.05 mmol), propargyl chloride (0.025 mL, 0.33 mmol), and potassium carbonate (14 mg, 0.1 mmol) in DMF (0.3 mL) was stirred at 80°C for 16 hours. The reaction mixture



- 172 -

was cooled to room temperature, treated with water (5 mL), and the resulting dark brown precipitate was collected by filtration. The crude solid was dissolved in 0.6 mL DMSO and purified by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 2201). Appropriate fractions were concentrated *in vacuo* to yield the final product as a white solid (4 mg.). Retention time using HPLC Method 2: 4.1 min. LRMS (ES pos.)  $m/z=438$  (M+1).

2-{2-(2-(6-Aminopurin-9-ylmethyl)-5-methyl-4-oxo-4H-quinazolin-3-yl)-phenoxy}-acetamide (D-070F)



20

A mixture of D-070B (20 mg, 0.05 mmol), 2-chloroacetamide (14 mg, 0.15 mmol), and potassium carbonate (21 mg, 0.15 mmol) in DMF (0.3 mL) was stirred at 80°C for 16 hours. The reaction mixture was cooled to room temperature, treated with water (5 mL), and the resulting precipitate was collected by filtration. Retention time using HPLC Method 2: 3.4 min. LRMS (ES pos.)  $m/z=457$  (M+1).

30


Additional compounds of the present invention were prepared by the following synthetic procedures.

- 173 -

Additional compounds of the invention follow, together with the synthetic route to compounds D-071 to D-118.


5      Procedure D

10



4a


15



5a

20


25



4b

- 174 -

5




5b

## Procedure E

10

15




5a

20


25

30



- 175 -


5



5b


10

15



20

25




5c

30


- 176 -

5




10

15



20




25




- 177 -

5



10


15



20

5e



25




30



- 178 -

**Procedure F**

4b



6a

**Procedure D:** A mixture of amide 4a or 4b, Fmoc-glycyl-chloride, and glacial acetic acid was heated to 120°C for 1 to 4 hours. The resulting mixture was concentrated in vacuo and purified by flash chromatography to provide the protected, cyclized amine. This material was combined with 10

- 179 -

equivalents octanethiol and a catalytic amount of DBU in THF and stirred at ambient temp until consumption of starting material was indicated by LCMS. The reaction was poured directly onto a flash column (equilibrated in  $\text{CH}_2\text{Cl}_2$ ) and eluted with 0-5% MeOH/ $\text{CH}_2\text{Cl}_2$  to provide the free amine, 5a or 5b. Compound 5c was prepared in an analogous manner using (+) Fmoc-alanyl-chloride in place of Fmoc-glycyl chloride.

10

**Procedure D1:** Amide 4a was admixed with Cbz- $\alpha$ -aminobutyric acid OSu, DIEA, DMAP (cat.), and toluene, then stirred at reflux for 3 days. The resulting mixture was purified by flash chromatography (EtOAc/hexanes) to provide the protected, cyclized amine. The amine was dissolved in EtOH with a catalytic amount of Pd/C, and allowed to stir at ambient temperature under  $\text{H}_2$  until a complete reaction was indicated by LCMS. The reaction mixture was filtered, and the filtrate concentrated in *vacuo*. Purification by flash chromatography (MeOH/ $\text{CH}_2\text{Cl}_2$ ) provided the free amine 5d.

15

20

**Procedure D2:** Amide 4a was combined with Boc-Serine(OBn)-OSu, DIEA, DMAP (cat.), and toluene, then stirred at reflux for 4 days. The resulting mixture was purified by flash chromatography (EtOAc/hexanes) to provide the protected, cyclized amine. The product was dissolved in a mixture of TFA and  $\text{CH}_2\text{Cl}_2$ , and allowed to stir at ambient temperature until a complete reaction was indicated by LCMS. The reaction was concentrated in *vacuo* and

25

30




- 180 -

purified by flash chromatography (MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to provide free amine 5e.

5                   **Procedure E:** Compounds 5(a-e), the appropriate 6-chloropurine or 6-bromopurine, and DIEA were combined with EtOH or DMF in a small vial and heated to 80°C. The reaction was monitored regularly by LCMS and purified as stated.

10                   **Procedure F:** A mixture of amide 4b, acetoxyacetyl chloride, and glacial acetic acid was heated to 120°C and stirred for 2 hours. The cooled reaction was filtered and the solids washed with CH<sub>2</sub>Cl<sub>2</sub> to provide the cyclized acetate as a white  
15 solid. This material was combined with K<sub>2</sub>CO<sub>3</sub> in aqueous methanol and stirred for one hour, then concentrated in vacuo. The resulting solids were triturated from H<sub>2</sub>O to provide 6a as a white solid.



20


3-(2-Chlorophenyl)-5-fluoro-2-[(9H-purin-6-ylamino)-methyl]-3H-quinazolin-4-one (D-072)

25

Prepared according to procedure E using 5b (50 mg, 0.165 mmol) and 6-chloropurine (26 mg, 0.165 mmol) in 1 mL EtOH. After 5 days, reaction purified by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min,

- 181 -


10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220λ). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ: 12.99 (br s, 1H); 8.14 (br s, 1H); 8.12 (s, 1H); 7.85 (dt, J=5.7, 8.1 Hz., 1H); 7.68-7.79 (m, 3H); 7.57 (t, J=6.2 Hz., 1H); 7.57 (d, J=7.7 Hz., 1H); 7.50 (d, J=8.1 Hz., 1H); 7.35 (dd, J=8.4, 10.7 Hz., 1H); 4.15-4.55 (m, 2H). MS (ES): m/z 422 (M+H), 211.



10 2-[(2-Amino-9H-purin-6-ylamino)methyl]-3-(2-chlorophenyl)-5-fluoro-3H-quinazolin-4-one (D-074)

Prepared according to procedure E using 5b (50 mg, 0.165 mmol) and 2-amino-6-chloropurine (28 mg, 0.165 mmol) in 1 mL EtOH. After 5 days, reaction purified by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220λ). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ: 12.13 (br s, 1H); 7.86 (dt, J=5.6, 8.2 Hz., 1H); 7.76-7.83 (m, 2H); 7.68 (br s, 1H); 7.61 (t, J=5.7 Hz., 1H); 7.61 (d, J=7.2 Hz., 1H); 7.53 (d, J=8.2 Hz., 1H); 7.35 (dd, J=8.2, 10.9 Hz., 1H); 5.66 (br s, 2 H); 4.16-4.50 (m, 1H); 4.09 (q, J=5.3 Hz., 2H). MS (ES): m/z 437 (M+H), 219.

- 182 -




**5-Methyl-2-[(9H-purin-6-ylamino)methyl]-3-o-tolyl-3H-quinazolin-4-one (D-071)**

5

Prepared according to procedure E using 6-chloropurine (11 mg, 0.072 mmol) and 5a (20 mg, 0.072 mmol). After 5 days, the reaction was quenched with water and the resulting suspension filtered. The solids were purified by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220 $\lambda$ ). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  $\delta$ : 12.98 (br s, 1H); 8.14 (br s, 1H); 8.10 (s, 1H); 7.58-7.79 (m, 2H); 7.37-7.48 (m, 4H); 7.26-7.36 (m, 2H); 3.93-4.39 (m, 2H); 2.75 (s, 3H); 2.18 (s, 3H). MS (ES): *m/z* 398 (M+H), 199.

10


15



- 183 -

2-[(2-Amino-9H-purin-6-ylamino)methyl]-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-073)

5                   Prepared according to procedure E using 5a  
(189 mg, 0.677 mmol) and 2-amino-6-chloropurine (115  
mg, 0.677) in 3 mL EtOH. After 3 days, the reaction  
was filtered to remove excess purine and the fil-  
trate purified by HPLC (C18 Luna column, 4.6 x 250  
10 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15  
min, 100% acetonitrile at 18 min, detector at 220λ)  
to provide 7 mg of the product as the TFA salt. <sup>1</sup>H  
NMR (DMSO-d<sub>6</sub>) δ: 8.88 (br s, 1H); 8.21 (s, 1H);  
7.71 (t, J=7.7 Hz., 1H); 7.45-7.56 (m, 2H); 7.38-  
15 7.44 (m, 3H); 7.35 (d, J=7.5 Hz., 1H); 7.30 (br s,  
1H); 4.40 (dd, J=4.5, 17.5 Hz., 1H); 4.27 (dd,  
J=5.3, 17.4 Hz., 1H); 2.75 (s, 3H); 2.09 (s, 3H).  
MS (ES): m/z 413 (M+H), 207, 163.




20

2-[(2-Fluoro-9H-purin-6-ylamino)methyl]-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-076)

25                   Prepared according to procedure E using 5a  
(20 mg, 0.072 mmol) and 2-fluoro-6-chloropurine (16  
mg, 0.094 mmol) in 1 mL EtOH. After 18 hours, the  
reaction was purified by HPLC (C18 Luna column, 4.6  
x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over

- 184 -


15 min, 100% acetonitrile at 18 min, detector at 220λ) and subsequently recrystallized from EtOH to provide 14 mg of the product as a yellow solid. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ: 13.12 (br s, 1H); 8.40 (br s, 1H);  
5 8.15 (s, 1H); 7.66 (t, J=7.7 Hz, 1H); 7.35-7.49 (m, 4H); 7.31 (d, J=7.2 Hz., 1H); 4.00-4.22 (m, 2H); 3.17 (s, 1H); 2.74 (s, 3H); 2.18 (s, 3H). MS (ES): m/z 416 (M+H), 208.



10 (2-Chlorophenyl)-dimethylamino-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one (D-075)

15 D-015 (100 mg, 0.228 mmol) was combined with ammonium hydroxide (28-30%, 1 mL) in DMF (2 mL) and heated to 80°C. After 2 days, the reaction was purified by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100%  
20 acetonitrile at 18 min, detector at 220λ) to provide the product as a yellow solid, ~2mg. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ: 13.52 (br s, 1H); 8.46 (s, 1H); 8.42 (s, 1H); 7.69 (dd, J=2.1, 7.3 Hz, 1H); 7.62 (dd, J=1.6, 7.6 Hz., 1H); 7.61 (t, J=8.0 Hz., 1H); 7.37-7.48 (m, 2H); 7.05 (d, J=7.9 Hz., 1H); 6.96 (d, J=7.8 Hz., 1H); 4.32-4.45 (m, 2H); 2.80 (s, 6H). MS (ES): m/z  
25 464 (M+H), 232.

- 185 -




5        5-(2-Benzyloxyethoxy)-3-(2-chlorophenyl)-2-(9H-  
purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
         (D-078)

10                To a solution of 2-benzyloxyethanol (0.3  
mL) in DMF (1.0 mL) was added NaH (50 mg, 2.08  
mmol). After stirring for 5 minutes, 0.5 mL was  
added to a solution of D-015 (50 mg, 0.114 mmol)  
in anhydrous DMF (0.75 mL). The reaction was heated  
to 50°C and stirred for 3 days. Purification by  
15        HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-  
75% acetonitrile/water over 15 min, 100% aceto-  
nitrile at 18 min, detector at 220λ) provided the  
product as a heterogenous solid, 150 µg. MS (ES):  
m/z 571 (M+H), 481.

20

- 186 -




6-Aminopurine-9-carboxylic acid 3-(2-chlorophenyl)-  
5-fluoro-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl  
ester (D-079)

5

To a solution of 3b (20 mg, 0.066 mmol) in  
CH<sub>2</sub>Cl<sub>2</sub> (500  $\mu$ L) at 0°C was added phosgene (2M/tolu-  
ene, 36  $\mu$ L, 0.072 mmol), followed by adenine (10 mg,  
0.072 mmol), and DIEA (25  $\mu$ L, 0.145 mmol). The re-  
action was allowed to attain ambient temperature and  
stir for 8 days. Purification by HPLC (C18 Luna  
column, 4.6 x 250 mm, 4.7 mL/min, 10-75% aceto-  
nitrile/water over 15 min, 100% acetonitrile at 18  
min, detector at 220 $\lambda$ ) provided the product as a  
mixture. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$ : 11.04 (br s, 1H); 8.61  
(s, 1H); 8.40 (s, 1H); 7.85-7.95 (m, 1H); 7.76 (dd,  
J=5.4, 9.6 Hz, 1H); 7.70-7.78 (m, 1H); 7.52-7.63 (m,  
3H); 7.38 (dt, J=8.3, 10.6 Hz., 1H); 4.76-4.89 (m,  
2H). MS (ES): m/z 466 (M+H), 331, 305.

20


- 187 -



- 5        N-[3-(2-Chlorophenyl)-5-fluoro-4-oxo-3,4-dihydro-  
quinazolin-2-ylmethyl]-2-(9H-purin-6-ylsulfanyl)-  
acetamide (D-077)
- 10                    (9H-Purin-6-ylsulfanyl)-acetic acid (63  
mg, 0.296 mmol), 5b (108 mg, 0.355 mmol), EDC (68  
mg, 0.355 mmol), HOBT (48 mg, 0.355 mmol), DIEA (62  
μL, 0.355 mmol), and DMF (1 mL) were combined in a  
15 flask and stirred at ambient temperature for one  
hour. The reaction was diluted with EtOAc (20 mL)  
and washed with dilute brine (2 x 13 mL). The or-  
ganic phase was concentrated in vacuo and chroma-  
tographed in 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to provide the 91 mg of  
the product as a viscous, peach foam. <sup>1</sup>H NMR (DMSO-  
20 d<sub>6</sub>) δ: 12.88 (br s, 1H); 8.72 (s, 1H); 8.62 (t,  
J=5.0 Hz, 1H); 8.49 (s, 1H); 7.88 (dt, J=5.6, 8.2  
Hz, 1H); 7.73-7.78 (m, 1H); 7.67-7.72 (m, 1H); 7.57-  
7.65 (m, 2H); 7.38 (d, J=8.1 Hz., 1H); 7.36 (dd,  
J=8.3, 11.1 Hz., 1H); 4.11-4.24 (m, 2H); 3.96 (dd,  
25 J=5.0, 17.4 Hz, 1H); 3.78 (dd, J=5.2, 17.4 Hz, 1H).  
MS (ES): m/z 496 (M+H), 248.



- 188 -



2-[1-(2-Fluoro-9H-purin-6-ylamino)ethyl]-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-080)


5

Prepared according to procedure E using 5c (50 mg, 0.17 mmol) and 2-fluoro-6-chloropurine (35 mg, 0.204 mmol) in 1.2 mL EtOH. Purification by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220λ) provided two atropisomers as white solids. Data for one of these follows: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.48 (br d, J=6.4 Hz, 1H); 8.17 (s, 1H); 7.69 (t, J=7.8 Hz, 1H); 7.53 (d, J=7.8 Hz, 1H); 7.44 (d, J=7.8 Hz, 2H); 7.33 (d, J=7.2 Hz, 2H); 7.07 (br t, J=7.2 Hz, 1H); 4.80 (br t, J=6.8 Hz, 1H); 2.74 (s, 3H); 2.09 (s, 3H); 1.38 (d, J=6.7 Hz, 3H). MS (ES): m/z 430 (M+H), 215.

10


15

- 189 -




**5-Methyl-2-[1-(9H-purin-6-ylamino)ethyl]-3-o-tolyl-3H-quinazolin-4-one (D-081)**

5                    Prepared according to procedure E using 5c  
                      (50 mg, 0.17 mmol) and 6-chloropurine (32 mg, 0.204  
                      mmol) in 1.2 mL EtOH. Purification by HPLC (C18  
                      Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75%  
                      acetonitrile/water over 15 min, 100% acetonitrile at  
 10                   18 min, detector at 220λ) provided two atropisomers  
                      as yellow solids. Data for one of these follows: <sup>1</sup>H  
                      NMR (DMSO-*d*<sub>6</sub>) δ: 8.39 (br s, 1H); 8.34 (s, 1H);  
                      8.18 (s, 1H); 7.71 (t, J=7.7 Hz, 1H); 7.56 (d, J=7.9  
                      Hz, 1H); 7.49 (d, J=6.9 Hz, 1H); 7.28-7.43 (m, 3H);  
 15                   7.20 (br s, 1H); 5.06 (br s, 1H); 2.73 (s, 3H); 2.04  
                      (s, 3H); 1.51 (d, J=6.6 Hz, 3H). MS (ES): m/z 412  
                      (M+H), 206.



20


- 190 -



2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-081a)

Synthesized according to procedure E using compound 5c (61 mg, 0.209 mmol) and 2-amino-6-chloropurine (43 mg, 0.251 mmol) in 1 mL EtOH. Purification by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220 nm) provided a white solid consisting of a mixture of two atropisomers (in brackets in nmr). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.93-9.02 (m, 1H); [8.19 (s), 8.15 (s), 1H]; [7.76 (t, J=8.1 Hz), 7.73 (t, J=7.8 Hz), 1H]; [7.64 (d, J=7.9 Hz), 7.57 (d, J=8.0 Hz), 1H]; [7.50 (d, J=7.7 Hz), 7.45 (d, J=7.8 Hz), 1H]; 7.29-7.40 (m, 3H); [7.23 (t, J=7.5 Hz), 7.15-7.22 (m), 1H]; [7.09 (t, J=7.5 Hz), 6.98 (d, J=7.3 Hz), 1H]; [5.28 (dd, J=7.2, 8.0 Hz), 4.96 (pent, J=7.0 Hz), 1H]; 2.75 (s, 3H); [2.10 (s), 1.84 (s), 3H]; [1.51 (d, J=6.6 Hz), 1.39 (d, J=6.7 Hz), 3H]. MS (ES): m/z 427 (M + H), 214.


- 191 -



5-Methyl-2-(1-(9H-purin-6-ylamino)propyl)-3-o-tolyl-3H-quinazolin-4-one (D-081b)

Synthesized according to procedure E using compound 5d (100 mg, 0.325 mmol) and 6-bromopurine (78 mg, 0.390 mmol) in 2 mL EtOH. Purification by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220 nm) provided two atropisomers as yellow solids. Data for one isomer follows: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.64 (br s, 1H); 8.44 (s, 1H); 8.27 (s, 1H); 7.72 (t, J=7.7 Hz, 1H); 7.56 (d, J=8.0 Hz, 1H); 7.50 (d, J=6.6 Hz, 1H); 7.34-7.44 (m, 2H); 7.35 (d, J=7.4 Hz, 1H); 7.18-7.27 (m, 1H); 4.85-5.01 (m, 1H); 2.73 (s, 3H); 2.04-2.19 (m, 1H); 1.99 (s, 3H); 1.78-1.91 (m, 1H); 0.79 (t, J=7.0 Hz, 3H). MS (ES): m/z 426 (M+H), 213.


- 192 -



2-(1-(2-Fluoro-9H-purin-6-ylamino)propyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-081c)

Synthesized according to procedure E using compound 5d (100 mg, 0.325 mmol) and 2-fluoro-6-chloropurine (78 mg, 0.455 mmol) in 2 mL EtOH. Purification by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220 nm) provided two atropisomers as off-white solids. Data for one isomer: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.46 (br d, J=7.1 Hz, 1H); 8.20 (s, 1H); 7.71 (t, J=7.7 Hz, 1H); 7.55 (d, J=7.9 Hz, 1H); 7.45 (d, J=7.3 Hz, 1H); 7.28-7.37 (m, 3H); 7.00 (t, J=7.3 Hz, 1H); 4.66 (q, J=6.7 Hz, 1H); 2.74 (s, 3H); 2.10 (s, 3H); 1.65-1.95 (m, 2H); 0.80 (t, J=7.1 Hz, 3H). MS (ES): m/z 444 (M+H), 222.


- 193 -



2-[1-(2-Amino-9H-purin-6-ylamino)propyl]-5-methyl-3-o-tolyl-3H-quinazolin-4-one (Compound 081d)

Synthesized according to procedure E using compound 5d (100 mg, 0.325 mmol) and 2-amino-6-bromopurine (104 mg, 0.488 mmol) in 2 mL EtOH. Purification by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220 nm) provided an off-white solid consisting of a mixture of two atropisomers (in brackets in nmr). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.89 (br d, J=7.8 Hz, 1H); [8.20 (s), 8.17 (s), 1H]; 7.75 (q, J=7.6 Hz, 1H); [7.62 (d, J=7.9 Hz); 7.57 (d, J=7.8 Hz), 1H]; 7.48 (t, J=7.3 Hz, 1H); 7.25-7.43 (m, 4H); 7.15 (br s, 1H); 7.02-7.12 (m, 1H); [5.03-5.15 (m), 4.77-4.87 (m), 1H]; 2.74 (s, 3H); [2.11 (s), 1.83 (s), 3H]; 1.65-2.19 (m, 2H); [0.83 (t, J=7.3 Hz), 0.80 (t, J=7.5 Hz), 3H]. MS (ES): m/z 441 (M + H), 221.

- 194 -



2-[2-Benzyloxy-1-(9H-purin-6-ylamino)ethyl]-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-081e)


Synthesized according to procedure E using compound 5e (212 mg, 0.413 mmol) and 6-bromopurine (107 mg, 0.537 mmol) in 2 mL EtOH. Purification by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 2201) provided two atropisomers as brown solids. Data for one isomer: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.45-8.63 (m, 1H); 8.35-8.44 (m, 1H); 8.27 (s, 1H); 7.75 (t, J=7.7 Hz, 1H); 7.59 (d, J=7.8 Hz, 1H); 7.30-7.44 (m, 3H); 7.21-7.30 (m, 4H); 7.13-7.19 (m, 2H); 6.95-7.07 (m 1H); 5.35-5.45 (m, 1H); 5.14-5.26 (m, 1H); 4.43 (s, 2H); 3.94-4.04 (m, 1H); 3.67 (dd, J=6.0, 9.4 Hz, 1H); 2.74 (s, 3H); 2.01 (s, 3H). MS (ES): m/z 518 (M+H), 410.

The following compounds of the present invention (D-082 through D-109) were prepared as outlined in Procedure C, using 2-chloromethyl-5-methyl-3-o-tolyl-3H-quinazolin-4-one (10 mg), the appropriate nucleophile XH (20 mg, excess), and

- 195 -

potassium carbonate (10 mg) in DMF (0.25 mL). The reaction mixture was stirred 16 h at room temperature, quenched with water, and the crude solid product was collected by filtration and air dried. The crude material was dissolved in 0.5 mL of DMSO and purified by reversed-phase HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220 $\lambda$ ). Appropriate fractions were concentrated *in vacuo* to yield the final products.

2-(6-Dimethylaminopurin-9-ylmethyl)-5-methyl-3-*o*-tolyl-3*H*-quinazolin-4-one (D-082)




Yield: 8.1 mg.  $^1\text{H}$  NMR (300 MHz,  $d_6$ -DMSO)  $\delta$ : 8.13 (s, 1H), 8.11 (s, 1H), 7.60 (t,  $J=7.8$  Hz, 1H), 7.54-7.38 (m, 4H), 7.30 (d,  $J=7.4$  Hz, 1H), 7.20 (d,  $J=8.1$  Hz, 1H), 5.11 (d,  $J=17.4$  Hz, 1H), 4.76 (d,  $J=17.4$  Hz, 1H), 3.33 (s, 6H), 2.73 (s, 3H), 2.20 (s, 3H). LRMS (ES pos.)  $m/z$  = 426 (M+1).



- 196 -

5-Methyl-2-(2-methyl-6-oxo-1,6-dihydro-purin-7-ylmethyl)-3-o-tolyl-3H-quinazolin-4-one (D-083)




10

Yield: 3.3 mg. <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 12.06 (s, 1H), 8.12 (s, 1H), 7.60 (t, J=7.8 Hz, 1H), 7.55-7.38 (m, 4H), 7.30 (d, J=7.4 Hz, 1H), 7.15 (d, J=7.9 Hz, 1H), 5.26 (d, J=17.4 Hz, 1H), 4.94 (d, J=17.4 Hz, 1H), 2.73 (s, 3H), 2.32 (s, 3H), 2.24 (s, 3H).

Alkylation at purine N, assigned arbitrarily based on downfield shift of methylene protons due to the carbonyl group. LRMS (ES pos.) m/z = 413 (M+1).

5-Methyl-2-(2-methyl-6-oxo-1,6-dihydro-purin-9-ylmethyl)-3-o-tolyl-3H-quinazolin-4-one (D-084)




30

Purified from same reaction mixture as D-083. Yield: 3.6 mg. <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) 12.17 (s, 1H), 7.96 (s, 1H), 7.63 (t, J=7.8 Hz, 1H), 7.57-7.39 (m, 4H), 7.32 (d, J=7.4 Hz, 1H), 7.26 (d, J=8.1 Hz, 1H), 5.08 (d, J=17.2 Hz, 1H), 4.70 (d, J=17.2 Hz, 1H), 2.73 (s, 3H), 2.27 (s, 3H), 2.17 (s, 3H). LRMS (ES pos.) m/z = 413 (M+1).

40


- 197 -

2-(Amino-dimethylaminopurin-9-ylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-085)



Yield: 6.7 mg. <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 7.66 (s, 1H), 7.61 (d, J=7.8 Hz, 1H), 7.55-7.40 (m, 4H), 7.32-7.26 (m, 2H), 6.74 (s, 2H), 4.94 (d, J=17.2 Hz, 1H), 4.63 (d, J=17.2 Hz, 1H), 4.63 (d, J=17.2 Hz, 1H), 2.97 (s, 6H), 2.73 (s, 3H), 2.17 (s, 3H), 2.08 (s, 3H). LRMS (ES pos.) m/z = 441 (M+1).

2-(2-Amino-9H-purin-6-ylsulfanylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-086)




Yield: 9.5 mg. <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 12.54 (s, 1H), 7.89 (s, 1H), 7.69 (t, J=7.8 Hz, 1H), 7.51 (d, J=8.0 Hz, 1H), 7.51 (d, J=8.0 Hz, 1H), 7.43 (t, J=3.9 Hz, 1H), 7.34 = 7.26 (m, 4H), 6.16 (s, 2H), 4.32 (AB quartet, J<sub>AB</sub>=14.8 Hz, Δn=23.7), 2.74 (s, 3H), 2.09 (s, 3H). LRMS (ES pos.) m/z = 430 (M+1).

- 198 -

2-(4-Amino-1,3,5-triazin-2-ylsulfanylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-087)

5




10

Yield: 5.8 mg.  $^1\text{H}$  NMR (300 MHz,  $d_6$ -DMSO)  $\delta$ : 8.10 (s, 1H), 7.70 (t,  $J=7.8$  Hz, 1H), 7.58 (s, 1H), 7.52 (d,  $J=8.0$  Hz, 1H), 7.48-7.26 (m, 6H), 4.08 (s, 2H), 2.73 (s, 3H), 2.09 (s, 3H). LRMS (ES pos.)  $m/z$  = 391 (M+1).

15

5-Methyl-2-(7-methyl-7H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one (D-088)

20



25


30

Yield: 3.1 mg.  $^1\text{H}$  NMR (300 MHz,  $d_6$ -DMSO)  $\delta$ : 8.52 (s, 1H), 8.49 (s, 1H), 7.70 (t,  $J=7.8$  Hz, 1H), 7.50 (d,  $J=7.8$  Hz, 1H), 7.45 (d,  $J=7.1$  Hz, 1H), 7.35-7.20 (m, 4H), 4.41 (AB quartet,  $J_{AB}=15.3$  Hz,  $\Delta\nu=19.2$  Hz), 4.08 (s, 3H), 2.73 (s, 3H), 2.12 (s, 3H). LRMS (ES pos.)  $m/z$  = 406 (M+1).

35


- 199 -

5-Methyl-2-(2-oxo-1,2-dihydro-pyrimidin-4-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one (D-089)



Yield: 2.4 mg. <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 11.49 (s, 1H), 7.70 (t, J=7.8 Hz, 1H), 7.60 (brt, J=6.0 Hz, 1H), 7.53-7.48 (m, 2H), 7.46-7.28 (m, 4H), 6.31 (d, J=6.7 Hz, 1H), 4.05 (s, 2H), 2.73 (s, 3H), 2.12 (s, 3H). LRMS (ES pos.) m/z = 391 (M+1).

20 5-Methyl-2-purin-7-ylmethyl-3-o-tolyl-3H-quinazolin-4-one (D-090)




<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 9.04 (s, 1H), 8.97 (s, 1H), 8.48 (s, 1H), 7.65-7.54 (m, 2H), 7.53-7.39 (m, 3H), 7.31 (d, J=7.4 Hz, 1H), 7.13 (d, J=8.0 Hz, 1H), 5.31 (d, J=17.6 Hz, 1H), 5.16 (d, J=17.6 Hz, 1H), 2.73 (s, 3H), 2.09 (s, 3H). Alkylation at purine N7 was determined by NOE enhancement between the purine 6-position proton and methylene protons on the linker between the purine and quinazolinone groups. LRMS (ES pos.) m/z = 383 (M+1).

40

- 200 -

5-Methyl-2-purin-9-ylmethyl-3-o-tolyl-3H-quinazolin-4-one (D-091)




10

From same reaction that produced D-090. <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 9.17 (s, 1H), 8.86 (s, 1H), 8.55 (s, 1H), 7.59 (t, J=7.8 Hz, 1H), 7.55-7.42 (m, 4H), 7.30 (d, J=7.4 Hz, 1H), 7.13 (d, J=8.0 Hz, 1H), 5.26 (d, J=17.5 Hz, 1H), 4.92 (d, J=17.5 Hz, 1H), 2.73 (s, 3H), 2.19 (s, 3H). Alkylation at purine N9 suggested by the lack of NOE enhancement between purine 6-position protons and the linker methylene protons. LRMS (ES pos.) m/z = 383 (M+1).

20

5-Methyl-2-(9-methyl-9H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one (D-092)



30


35

<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 8.52 (s, 1H), 8.42 (s, 1H), 7.69 (t, J=7.7 Hz, 1H), 7.50 (d, J=8.0 Hz, 1H), 7.44 (d, J=7.6 Hz, 1H), 7.36-7.27 (m, 4H), 4.38 (AB quartet, J<sub>AB</sub>=15.5 Hz, Δν=21.0 Hz), 3.80 (s, 3H), 2.73 (s, 3H), 2.12 (s, 3H). LRMS (ES pos.) m/z = 429 (M+1).

40


- 201 -

2-(2,6-Diamino-pyrimidin-4-ylsulfanylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-093)



<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 7.70 (t, J=7.7 Hz, 1H), 7.54 (d, J=8.0 Hz, 1H), 7.45-7.27 (m, 5H), 6.22 (br s, 1H), 5.80 (br s, 1H), 3.99 (AB quartet, J<sub>AB</sub>=14.6 Hz, Δν=26.9 Hz, 2H), 2.73 (s, 3H), 2.08 (s, 3H).  
LRMS (ES pos.) m/z = 405 (M+1).

5-Methyl-2-(5-methyl-[1,2,4]triazolo[1,5-a]pyrimidin-7-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one (D-094)




30

<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 8.57 (s, 1H), 7.73 (t, J=7.8 Hz, 1H), 7.55-7.35 (m, 4H), 7.18 (s, 1H), 4.27 (s, 2H), 2.74 (s, 3H), 2.55 (s, 3H), 2.08 (s, 3H).  
35 LRMS (ES pos.) m/z = 429 (M+1).

- 202 -


5-Methyl-2-(2-methylsulfanyl-9H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one (D-095)



<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 13.30 (s, 1H), 8.29 (s, 1H), 7.72 (t, J=7.8 Hz, 1H), 7.54 (d, J=7.8 Hz, 1H), 7.47 (d, J=6.3 Hz, 1H), 7.38-7.26 (m, 4H), 4.34 (AB quartet, J<sub>AB</sub>=16.1 Hz, Δν=23.6 Hz, 2H), 2.74 (s, 3H), 2.32 (s, 3H), 2.10 (s, 3H). LRMS (ES pos.) m/z = 461 (M+1).

20

2-(2-Hydroxy-9H-purin-6-ylsulfanylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-096)




<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 8.08 (s, 1H), 7.69 (t, J=7.8 Hz, 1H), 7.50 (brd, J=t.8 Hz, 2H), 7.33-7.50 (m, 4H), 4.28 (AB quartet, J<sub>AB</sub>=15.5 Hz, Δν=21.3 Hz, 2H), 2.74 (s, 3H), 2.12 (s, 3H). LRMS (ES pos.) m/z = 431 (M+1).

- 203 -

5-Methyl-2-(1-methyl-1*H*-imidazol-2-ylsulfanylmethyl)-3-*o*-tolyl-3*H*-quinazolin-4-one (D-097)

5




10

<sup>1</sup>H NMR (300 MHz, *d*<sub>6</sub>-DMSO) δ: 7.69 t, *J*=7.8 Hz, 1H), 7.46-7.37 (m, 5H), 7.32 (d, *J*=7.3 Hz, 1H), 7.20 (d, *J*=1.0 Hz, 1H), 6.48 (d, *J*=1.0 Hz), 3.83 (AB quartet, *J*<sub>AB</sub>=15.0 Hz, Δ*ν*=18.8 Hz, 1H), 3.55 (s, 3H), 2.73 (s, 3H), 2.09 (s, 3H). LRMS (ES pos.) *m/z* = 364 (*M*+1).

15

5-Methyl-3-*o*-tolyl-2-(1*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-3*H*-quinazolin-4-one (D-098)

20



25

<sup>1</sup>H NMR (300 MHz, *d*<sub>6</sub>-DMSO) δ: 13.98 (s, 1H), 8.47 (s, 1H), 7.70 (t, *J*=7.8 Hz, 1H), 7.49 (d, *J*=7.9 Hz, 1H), 7.44-7.31 (m, 5H), 4.04 (AB quartet, *J*<sub>AB</sub>=15.5 Hz, Δ*ν*=19.1 Hz, 1H), 2.74 (s, 3H), 2.10 (s, 3H). LRMS (ES pos.) *m/z* = 364 (*M*+1).

30




- 204 -

2-(2-Amino-6-chloro-purin-9-ylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-099)

5

X =




10

LRMS (ES pos.) 432 (M+1).

2-(6-Aminopurin-7-ylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-100)

15

X =




20

<sup>1</sup>H NMR (300 MHz, *d*<sub>6</sub>-DMSO) δ: 8.19 (s, 3H), 7.66 (t, J=7.8 Hz, 1H), 7.59-7.43 (m, 5H), 7.34 9d, J=7.4 Hz, 1H), 7.23 (d, J=8.0 Hz, 1H), 6.90 (s, 2H), 5.21 (AB quartet, *J*<sub>AB</sub>=17.4 Hz, Δ*v*=22.1 Hz, 2H), 2.72 (s, 3H), 1.93 (s, 3H). Alkylation at purine N7 was confirmed by NOE enhancements between the following protons:

30 1) Exocyclic amine and methylene protons; 2) Exocyclic amine and toluyl methyl protons. LRMS (ES pos.) *m/z* = 398 (M+1).

- 205 -

2-(7-Amino-1,2,3-triazolo[4,5-d]pyrimidin-3-yl-  
methyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one  
(D-101)




10

<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 8.43 (br s, 1H), 8.19 (s, 1H), 8.10 (br s, 1H), 7.62 (t, J=7.8 Hz, 1H), 7.49-7.28 (m, 5H), 7.22 (d, J=8.1 Hz, 1H), 5.49 (d, J=17.0 Hz, 1H), 5.19 (d, J=17.0 Hz, 1H), 2.73 (s, 3H), 2.11 (s, 3H). Alkylation at purine N7 determined by similarity to nmr spectrum of D-030. LRMS (ES, pos.) m/z = 399 (M+1).

20

2-(7-Amino-1,2,3-triazolo[4,5-d]pyrimidin-1-yl-  
methyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one  
(D-102)




30

From same reaction mixture as D-101. <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 8.27 (s, 1H), 8.20 (br s, 1H), 8.05 (br s, 1H), 7.70 (t, J=7.8 Hz, 1H), 7.47-7.26 (m, 6H), 5.61 (AB quartet, J<sub>AB</sub>=16.0 Hz, Δν=20.7 Hz, 2H), 2.75 (s, 3H), 1.98 (s, 3H). Alkylation at purine N7 determined by similarity to nmr spectrum of D-100. LRMS (ES pos.) m/z = 399 (M+1).

40

- 206 -


2-(6-Amino-9H-purin-2-ylsulfanylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-103)



10

<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 12.62 (s, 1H), 7.93 (s, 1H), 7.69 (t, J=7.7 Hz, 1H), 7.51 (d, J=8.1 Hz, 1H), 7.42 (dd, J=7.6, 1.7 Hz, 1H), 7.35-7.15 (m, 6H), 4.12 (AB quartet, J<sub>AB</sub>=14.5 Hz, Δν=18.2 Hz, 2H), 2.73 (s, 3H), 2.10 (s, 3H). LRMS (ES pos.) m/z = 430 (M+1).

2-(2-Amino-6-ethylamino-pyrimidin-4-ylsulfanylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-104)



25

30


<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 7.70 (t, J=7.8 Hz, 1H), 7.53 (d, J=8.0 Hz, 1H), 7.44-7.31 (m, 5H), 6.69 (br s, 1H), 5.83, (br s, 2H), 5.61 (s, 1H), 4.03 (d, J=14.6 Hz, 1H), 3.95 (d, J=14.6 Hz, 1H), 3.22-3.11 (m, 2H), 2.73 (s, 3H), 2.08 (s, 3H), 1.06 (t, J=7.1 Hz, 3H). LRMS (ES pos.) m/z = 433 (M+1).

35

- 207 -

2-(3-Amino-5-methylsulfanyl-1,2,4-triazol-1-yl-  
methyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one  
(D-105)

5



10


Yield: 5.0 mg. <sup>1</sup>H NMR (300 MHz, d<sub>4</sub>-MeOH) δ: 7.67  
(t, J=7.8 Hz, 1H), 7.55-7.37 (m, 4H), 7.35-7.27 (m,  
2H), 4.77 (d, J=17.1 Hz, 1H), 4.60 (d, J=17.1 Hz,  
1H), 2.80 (s, 3H), 2.43 (s, 3H), 2.14 (s, 3H). LRMS  
(ES pos.) m/z = 393 (M+1).

15

20

2-(5-Amino-3-methylsulfanyl-1,2,4-triazol-1-  
ylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one  
(D-106)

25




30

Yield: 0.6 mg. Purified from same reaction mixture  
as D-105. <sup>1</sup>H NMR (300 MHz, d<sub>4</sub>-MeOH) δ: 7.67 (t,  
J=7.8 Hz, 1H), 7.50-7.24 (m, 6H), 4.83 (d, J=16.5  
Hz, 1H), 4.70 (d, J=16.5 Hz, 1H), 2.79 (s, 3H), 2.47  
(s, 3H), 2.14 (s, 3H). LRMS (ES pos.) m/z = 393  
(M+1).

35

- 208 -


5-Methyl-2-(6-methylaminopurin-9-ylmethyl)-3-o-  
tolyl-3H-quinazolin-4-one (D-107)



10

Yield: 5.0 mg <sup>1</sup>H NMR (300 MHz, d<sub>4</sub>-MeOH) δ: 8.17  
(s, 1H), 8.03 (s, 1H), 7.54-7.43 (m 4H), 7.31-7.23  
15 (m, 2H), 5.14 (d, J=17.5 Hz, 1H), 4.90 (d, J=17.5  
Hz, 1H), 3.14 (br s, 3H), 2.79 (s, 3H), 2.22 (s,  
3H). LRMS (ES pos.) m/z = 412 (M+1).

20 2-(6-Benzylaminopurin-9-ylmethyl)-5-methyl-3-o-  
tolyl-3H-quinazolin-4-one (D-108)




30

Yield: 6.7 mg. <sup>1</sup>H NMR (300 MHz, d<sub>4</sub>-MeOH) δ: 8.13  
(s, 1H), 8.04 (s, 1H), 7.58 (t, J=7.8 Hz, 1H), 7.51-  
35 7.21 (m, 11H), 5.15 (d, J=17.5 Hz, 1H), 4.91 (d,  
J=17.5 Hz, 1H), 4.83 (s, 2H, under H<sub>2</sub>O Peak), 2.79  
(s, 3H), 2.22 (s, 3H). LRMS (ES pos.) m/z = 488  
(M+1).

- 209 -

2-(2,6-Diaminopurin-9-ylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one (D-109)

5




10

Doubled the amounts of all reactants. Yield: 14 mg. <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 8.53 (br s, 2H), 8.01 (s, 1H), 7.64 (t, J=7.8 Hz, 1H), 7.53-7.40 (m, 4H), 7.33 (d, J=7.4 Hz, 1H), 7.27 (d, J=7.9 Hz, 1H), 4.96 (d, J=17.5 Hz, 1H), 4.64 (d, J=17.5 Hz, 1H), 2.74 (s, 3H), 2.17 (s, 3H). LRMS (ES pos.) m/z = 413 (M+1).

20

Compounds D-110 through D-115 of the following general structure were prepared from the following Intermediates E-1 through E-3.




25

- 210 -

## Intermediate E-1.

5-Methyl-2-(9H-purin-6-ylsulfanylmethyl)-3,1-benzoxazin-4-one

5



Intermediate E-1


10                    **Step 1.** A suspension of 6-methylantranilic acid (2 g, 13.2 mmol) in chloroacetyl chloride (12 mL, large excess) was stirred at 115°C in a sealed vial for 30 min. The resulting solution was cooled to room temperature and treated with ether (~5 mL). After cooling at 4°C overnight, the resulting tan precipitate was collected by filtration, washed with ether, and dried in vacuo to yield the chloro intermediate (1.39 g, 50%). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ: 7.67 (t, J=7.8 Hz, 1H), 7.46 (d, J=7.9 Hz, 1H), 7.35 (d, J=7.6 Hz, 1H), 4.39 (s, 2H), 2.81 (s, 3H). LRMS (ES pos.) m/z = 210, (M+1).

20                    **Step 2.** A mixture of the chloro intermediate (50 mg, 0.25 mmol), 6-mercaptopurine monohydrate (43 mg, 0.25 mmol), and potassium carbonate (25 mg, 0.25 mmol) in dry DMF (0.5 mL) was stirred at room temperature for 30 min. The mixture was

- 211 -

poured into ethyl acetate (20 mL) and all insoluble material was filtered off and discarded. The filtrate was concentrated *in vacuo* to remove all ethyl acetate, and the residue was treated with ether, resulting in a light orange precipitate. The precipitate was collected by filtration, washed with ether, and dried *in vacuo* to afford Intermediate E-1 (41 mg, 51%).  $^1\text{H}$  NMR (300 MHz,  $d_6$ -DMSO)  $\delta$ : 8.64 (s, 1H), 8.39 (s, 1H), 7.73 (t,  $J=7.8$  Hz, 1H), 7.44-7.37 (m, 2H), 4.69 (s, 2H), 2.69 (s, 3H). LRMS (ES pos.)  $m/z$  = 326 (M+1).

## Intermediate E-2




A solution of 2-nitroacetanilide (1.0 g, 5.6 mmol) in EtOH was purged with nitrogen, treated with  $\text{Pd}(\text{OH})_2$  (20% by wt. on C, 200 mg, cat.), and shaken for 2 h under  $\text{H}_2$  (20 psi). The catalyst was removed by filtration through a 0.22  $\mu\text{m}$  cellulose acetate membrane (Corning), and the filtrate was concentrated *in vacuo* to afford the white crystalline solid product (800 mg, 96%).  $^1\text{H}$  NMR (300 MHz,  $d_6$ -DMSO)  $\delta$ : 9.12 (s, 1H), 7.14 (dd,  $J=7.8, 1.3$  Hz, 1H), 6.88 (dt,  $J=7.6, 1.5$  Hz, 1H), 6.70 (dd,  $J=8.0, 1.3$  Hz, 1H), 6.52 (dt,  $J=7.5, 1.4$  Hz, 1H), 4.85 (br s, 2H), 2.03 (s, 3H). LRMS (ES pos.)  $m/z$  = 151 (M+1).



- 212 -

## Intermediate E-3



A mixture of 2-fluoro-nitrobenzene (1.41 g, 10 mmol) and NaHCO<sub>3</sub> in EtOH (20 mL) was treated with (N,N,N'-trimethyl)-1,2-diaminoethane (1.1 g, 11 mmol) and was stirred 16 h at 80°C. Solvent was removed in vacuo, residue was treated with 0.1 M NaOH (120 mL), and the mixture was extracted with ethyl acetate (2 x 50 mL). The organic layers were combined and washed with 20 mL of water (1x) and brine (2x), dried with sodium sulfate, and concentrated in vacuo to an orange liquid (2.2 g, 100%; ESMS: m/z = 224, M+1).

This intermediate was dissolved in EtOH, the solution was purged with nitrogen, treated with Pd(OH)<sub>2</sub> (20% by wt. on C, 180 mg, cat.), and shaken for 2 h under H<sub>2</sub> (50 psi). The catalyst was removed by filtration through a 0.22 μm cellulose acetate membrane (Corning), and the filtrate was concentrated in vacuo to afford the red liquid product E-3 (1.8 g, 95%). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ: 8.64 (s, 1H), 7.03 (dd, J=8.3, 1.4 Hz, 1H), 6.91 (ddd, J=7.6, 7.2, 1.4 Hz, 1H), 6.73-6.67 (m, 2H), 4.20 (br s, 2H), 2.95 (t, J=6.7 Hz, 2H), 2.68 (s, 3H), 2.41 (t, J=6.7 Hz, 1H), 2.26 (s, 6H). LRMS (ES pos.) m/z=194 (M+1).

- 213 -

Compounds D-110 through D-115 were prepared as follows:

5-Methyl-2-(9H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one (D-110)



A mixture of Intermediate E-1 (40 mg) and o-toluidine (0.3 mL, large excess) was warmed at 100°C in a sealed vial for 16 h. The reaction mixture was cooled, treated with 1N HCl (2 mL) and ether (2 mL), and the resulting gray precipitate was collected by filtration, washed with ether, and air dried (19 mg crude). The crude solid was dissolved in 0.5 mL DMSO and purified by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220λ). Appropriate fractions were concentrated in vacuo to yield the final product as a white solid (4 mg). <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 13.52 (s, 1H), 8.47 (s, 1H), 8.43 (s, 1H), 7.69 (t, J=7.8 Hz, 1H), 7.50 (d, J=7.9 Hz, 1H), 7.46-7/43 (m, 1H), 7.37-7.25 (m, 4H), 4.37 (AB quartet, J<sub>AB</sub>=15.4 Hz, Δν=22.4 Hz, 2H), 2.74 (s, 3H), 2.12 (s, 3H). LRMS (ES pos.) m/z = 415 (M+1).

- 214 -

3-Isobutyl-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-quinazolin-4-one (D-111)

5




10 A mixture of Intermediate E-1 (40 mg) and isobutyl-  
amine (0.4 mL, large excess) was warmed at 120°C in  
a sealed vial for 16 h. Excess isobutylamine was  
allowed to evaporate, residue was dissolved in 1 mL  
DMSO and purified in two portions by HPLC (C18 Luna  
15 column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetoni-  
trile/water over 15 min, 100% acetonitrile at 18  
min, detector at 220λ). Appropriate fractions were  
concentrated in vacuo to yield the final product as  
a white solid (4 mg). <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ:  
20 13.75 (br s, 1H), 8.73 (s, 1H), 8.50 (s, 1H), 7.63  
(t, J=7.7 Hz, 1H), 7.42 (d, J=8.0 Hz, 1H), 7.28 (d,  
J=7.3 Hz, 1H), 4.96 (s, 2H), 4.00 (d, J=7.5 Hz, 2H),  
2.77 (s, 3H), 2.30-2.15 (m, 1H), 0.98 (d, J=6.7 Hz,  
1H). LRMS (ES pos.) m/z = 381 (M+1).

25

- 215 -

*N*-{2-[5-Methyl-4-oxo-2-(9*H*-purin-6-ylsulfanylmethyl)-4*H*-quinazolin-3-yl]-phenyl}-acetamide  
(D-112)

5



10

A mixture of Intermediate E-1 (80 mg, 0.25 mmol) and Intermediate E-2 (75 mg, 0.5 mmol, 2 eq) was warmed until melted in a sealed vial using a heat gun. The reaction mixture was triturated with ether and the solids were collected by filtration. The crude material was dissolved in 1 mL DMSO and purified in two portions by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220Å). Appropriate fractions were concentrated *in vacuo* to yield the final product as a white solid. <sup>1</sup>H NMR (300 MHz, *d*<sub>6</sub>-DMSO) δ: 13.52 (s, 1H), 9.52 (s, 1H), 8.48 (s, 3H), 8.42 (s, 3H), 8.02 (d, *J*=8.0 Hz, 1H), 7.69 (t, *J*=7.8 Hz, 1H), 7.51 (d, *J*=7.9 Hz, 1H), 7.45-7.37 (m, 2H), 7.31 (d, *J*=7.3 Hz, 1H), 7.19 (t, *J*=7.5 Hz, 1H), 4.38 (s, 2H), 2.74 (s, 3H), 1.93 (s, 3H). LRMS (ES pos.) *m/z* = 458 (*M*+1).

25

- 216 -

5-Methyl-3-(E-2-methyl-cyclohexyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one (D-113)

5



10


A mixture of Intermediate E-1 (80 mg, 0.25 mmol) and trans-2-methyl-1-aminocyclohexane (0.25 mL, large excess) was warmed in a sealed vial at 100°C for 16 h. The reaction mixture was triturated with ether and the solids were collected by filtration. The crude material was dissolved in 0.5 mL DMSO and purified by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220λ). Appropriate fractions were concentrated in vacuo to yield the final product as a white solid (1.5 mg). <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 13.5 (br s, 1H), 8.82 (s, 1H), 8.51 (s, 1H), 7.63 (t, J=7.7 Hz, 1H), 7.43 (d, J=7.9 Hz, 1H), 7.27 (d, J=7.4 Hz, 1H), 5.11 (d, J=14.5 Hz, 1H), 3.78-3.69 (m, 1H), 2.73 (s, 3H), 2.55-2.40 (m, 3H), 1.88-1.46 (m, 4H), 1.31-1.11 (m, 1H), 0.90-0.65 (m, 1H), 0.74 (d, J=6.7 Hz, 3H). LRMS (ES pos.) m/z = 421 (M+1).

30

- 217 -

2-[5-Methyl-4-oxo-2-(9H-purin-6-ylsulfanylmethyl)-  
4H-quinazolin-3-yl]-benzoic acid (D-114)

5




10 A mixture of Intermediate E-1 (80 mg, 0.25 mmol)  
methyl anthranilate (0.25 mL, large excess) was  
warmed in a sealed vial at 100°C for 16 h. The  
reaction mixture was triturated with ether and the  
solids were collected by filtration. The crude  
15 material was dissolved in 0.5 mL DMSO and purified  
by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min,  
10-75% acetonitrile/water over 15 min, 100% aceto-  
nitrile at 18 min, detector at 220λ). Appropriate  
fractions were concentrated in vacuo to yield the  
20 final product as a white solid (8 mg). <sup>1</sup>H NMR (300  
MHz, d<sub>6</sub>-DMSO) δ: 13.51 (s, 1H), 8.51 (s, 1H), 8.42  
(s, 1H), 8.11 (dd, J=7.4, 1.1 Hz, 1H), 7.88 (dt,  
J=7.7, 1.4 Hz, 1H), 7.70 (d, J=8.0 Hz, 1H), 7.57 (t,  
J=7.2 Hz, 1H), 7.49-7.35 (m, 3H), 4.58 (d, J=15.5  
25 Hz, 1H), 4.35 (d, J=15.5 Hz, 1H), 2.44 (s, 3H).  
LRMS (ES pos.) m/z = 445 (M+1).

- 218 -

3-{2-[(2-Dimethylamino-ethyl)-methyl-amino]-phenyl}-  
5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-  
quinazolin-4-one (D-115)

5




10

A mixture of Intermediate E-1 (40 mg, 0.25 mmol)  
Intermediate E-3 (0.2 mL, large excess) was warmed  
in a sealed vial at 100°C for 16 h. The reaction  
mixture was triturated with ether and the solids  
were collected by filtration. The crude material  
was dissolved in 1 mL DMSO and purified by HPLC in  
two portions (C18 Luna column, 4.6 x 250 mm, 4.7  
mL/min, 10-75% acetonitrile/water over 15 min, 100%  
acetonitrile at 18 min, 0.05% TFA in all solvents,  
detector at 220λ). Appropriate fractions were  
concentrated in vacuo to yield the final product as  
the TFA salt (11 mg). <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ:  
13.4 (br s, 1H), 9.27 (s, 1H), 8.52 (s, 1H), 8.44  
(s, 1H), 7.72 (t, J=7.8 Hz, 1H), 7.53 (d, J=7.9 Hz,  
1H), 7.40-7.33 (m, 4H), 7.10-7.04 (m, 1H), 4.42 (s,  
3H), 3.5 (m, 2H), 3.23-3.03 (m, 3H), 2.75 (s, 3H),  
2.68-2.56 (m, 8H). LRMS (ES pos.) m/z = 501 (M+1).

- 219 -

Compounds D-116 through D-118 were prepared as follows:



5

3-(2-Chlorophenyl)-5-methoxy-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one (D-116)

10 (R = Me, X = O)

A mixture of D-015 (25 mg) in 0.5 M NaOMe (2 mL in MeOH; large excess) was stirred at  $50^\circ C$  for 16 h in a sealed vial. The reaction mixture was cooled to room temperature, treated with water (5 mL), and the resulting precipitate was collected by filtration, washed with water, and air dried. The crude material was dissolved in 0.5 mL DMSO and purified by HPLC (C18 Luna column, 4.6 x 250 mm, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220 nm). Appropriate fractions were concentrated in vacuo to yield the final product as a white solid (5.3 mg).  $^1H$  NMR (300 MHz,  $d_6$ -DMSO)  $\delta$ : 13.52 (s, 1H), 8.48 (s, 1H), 8.44 (br s, 1H), 7.77 (t, J=8.2

15

20

25



- 220 -

Hz, 1H), 7.71-7.60 (m, 2H), 7.51-7.34 (m, 2H), 7.23 (d, J=8.2 Hz, 1H), 7.10 (d, J=8.4 Hz, 1H), 4.39 (AB quartet,  $J_{AB}$ =5.2 Hz,  $\Delta\nu$ =23.2 Hz, 2H), 3.85 (s, 3H). LRMS (ES positive) m/z = 451 (M+1).

5

3-(2-Chlorophenyl)-5-(2-morpholin-4-yl-ethylamino)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one (D-117)

10




15

A mixture of D-015 (25 mg) and 4-(aminoeth-2-yl)-morpholine (650 mg, large excess) was stirred at 50°C for 16 h. The crude reaction mixture was purified by HPLC (C18 Luna column, 4.6 x 250 mm, 4.7 mL/min, 10-75% acetonitrile/water over 15 min, 100% acetonitrile at 18 min, detector at 220λ). Appropriate fractions were concentrated *in vacuo* to yield the final product. <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-acetone) δ: 8.57 (br s, 1H), 8.47 (s, 1H), 8.37 (s, 1H), 7.72 (dd, J=7.7, 1.6 Hz, 1H), 7.65 (dd, J=8.0, 1.2 Hz, 1H), 7.57 (t, J=8.1 Hz, 1H), 7.49 (dt, J=7.7, 1.6 Hz, 1H), 7.40 (dt, J=7.7, 1.5 Hz, 1H), 6.86 (d, J=7.4 Hz, 1H), 6.82 (d, J=8.3 Hz, 1H), 4.55 (d, J=15.0 Hz, 1H), 4.42 (d, J=15.1 Hz, 1H), 4.05-3.90 (m, 4H), 3.90 (t, J=6.9 Hz, 2H), 3.75-3.4 (m, 4H), 3.54 (t, J=6.9 Hz, 2H). LRMS (ES positive) m/z = 549 (M+1).

35

- 221 -

3-Benzyl-5-methoxy-2-(9H-purin-6-ylsulfanylmethyl)-  
3H-quinazolin-4-one (D-118)



A mixture of D-043 (25 mg) in 0.5 M NaOMe (2 mL in  
20 MeOH; large excess) was stirred at 50°C for 16 h in  
a sealed vial. The reaction mixture was treated  
with 1 N HCl (1 mL) and aliquots of this solution  
(0.5 mL each) were purified by HPLC (C18 Luna  
25 trile/water over 15 min, 100% acetonitrile at 18  
min, detector at 220λ). Appropriate fractions were  
concentrated in vacuo to yield the final product as  
a white solid (6.6 mg). <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ:  
13.57 (s, 1H), 8.60 (s, 1H), 8.45 (s, 1H), 7.72 (t,  
30 J=8.1 Hz, 1H), 7.42-7.30 (m, 2H), 7.30-7.19 (m, 3H),  
7.15 (d, J=8.0 Hz, 1H), 7.06 (d, J=8.3 Hz, 1H), 5.43  
(s, 2H), 4.80 (s, 2H), 3.87 (s, 3H). LRMS (ES  
positive) m/z = 431 (M+1).

- 222 -

Compound D-999 (comparative)

3-(2-Chlorophenyl)-2-(1H-pyrazolo[3,4-d]pyrimidin-4-ylsulfanylmethyl)-3H-quinazolin-4-one

5

An analog compound, 3-(2-chlorophenyl)-2-(1H-pyrazolo[3,4-d]pyrimidin-4-ylsulfanylmethyl)-3H-quinazolin-4-one, also was synthesized generally in accordance with the described methods, except that a 4-mercapto-1H-pyrazolo[3,4-d]pyrimidine was substituted for the mercaptopurine in the final step.

10

#### EXAMPLE 11

#### Biochemical Assays of PI3K Potency and Selectivity

15

##### A. Biochemical Assay using 20 $\mu$ M ATP

Using the method described in Example 2, above, compounds of the invention were tested for inhibitory activity and potency against PI3K $\delta$ , and for selectivity for PI3K $\delta$  versus other Class I PI3K isozymes. In Table 2, IC<sub>50</sub> values ( $\mu$ M) are given for PI3K $\alpha$  ("Alpha"), PI3K $\beta$  ("Beta"), PI3K $\gamma$  ("Gamma"), and PI3K $\delta$  ("Delta"). To illustrate selectivity of the compounds, the ratios of the IC<sub>50</sub> values of the compounds for PI3K $\alpha$ , PI3K $\beta$ , and PI3K $\gamma$  relative to PI3K $\delta$  are given, respectively, as "Alpha/Delta Ratio," "Beta/Delta Ratio," and "Gamma/Delta Ratio."

20

25

The initial selectivity assays were done identically to the selectivity assay protocol in Example 2, except using 100  $\mu$ L Ecoscint for radio-label detection. Subsequent selectivity assays were done similarly using the same 3X substrate stocks except they contained 0.05 mCi/mL  $\gamma$ [<sup>32</sup>P]ATP and 3 mM

30

- 223 -

PIP<sub>2</sub>. Subsequent selectivity assays also used the same 3X enzyme stocks, except they now contained 3 nM of any given PI3K isoform.

For all selectivity assays, the test  
5 compounds were weighed out and dissolved into 10-50 mM stocks in 100% DMSO (depending on their respective solubilities) and stored at -20°C. Compounds were thawed (to room temperature or 37°C), diluted to 300 µM in water from which a 3-fold dilution  
10 series into water was done. From these dilutions, 20 µL was added into the assay wells alongside water blanks used for the enzyme (positive) control and the no enzyme (background) control. The rest of the assay was essentially done according to the selectivity assay protocol in Example 2.  
15

For those cases in which the greatest concentration used in the assay, i.e., 100 µM, did not inhibit activity of the enzyme by at least 50%, the table recites the percent activity remaining at that  
20 concentration (i.e., at 100 µM). In these cases, the true activity ratio(s) for the compounds cannot be calculated, since one of the required IC<sub>50</sub> values is missing. However, to provide some insight into the characteristics of these compounds, a hypothetical activity ratio is calculated using 100 µM substituted for the missing value. In such cases, the  
25 selectivity ratio must in fact be greater than the hypothetical value, and this is indicated by use of a greater than (>) symbol.

Table 2

Compound	Alpha IC <sub>50</sub>	Beta IC <sub>50</sub>	Delta IC <sub>50</sub>	Gamma IC <sub>50</sub>	Alpha/Delta Ratio	Beta/Delta Ratio	Gamma/Delta Ratio
D-000	86%	74%	0.33	7.7	>302	>302	23
D-001	83%	45	68		>1.5	0.66	
D-002	88%	78%	44		>2.3	>2.3	
D-003	92	53%	4		22	>24	
D-004	93%	89%	64		>2	>1.6	
D-005	89%	46	0.8		>121	56	
D-006	78%	6	0.15		>652	38	
D-007	82%	30	0.16		>619	188	
D-008	82%	68	1.2		>85	57	
D-009	82	6	0.12		683	50	
D-010	48	11	0.06	0.70	800	183	12
D-011	72%	55	0.10	1.0	>1,000	550	10
D-012	69%	11	0.17		>588	65	
D-013	71%	13	0.05	2.1	>2,000	260	42
D-014	63%	3.6	0.06	0.56	>1,667	60	9.3
D-015	65%	69%	0.21	3.6	>480	>480	17
D-016	91%	81%	40		>2.5	>3	

Table 2

Compound	Alpha IC <sub>50</sub>	Beta IC <sub>50</sub>	Delta IC <sub>50</sub>	Gamma IC <sub>50</sub>	Alpha/Delta Ratio	Beta/Delta Ratio	Gamma/Delta Ratio
D-017	89%	108%	12		>8	>8	
D-018	88%	93%	4.2		>24	>24	
D-019	67	105	7		10	15	
D-020	69%	69%	1.9		>53	>53	
D-021	100	110	1.6		62	68	
D-022	81%	110	0.8	40	>125	137.50	50
D-023	83%	91%	26		>4	>3.9	
D-024	100	76%	2.6		38	>38	
D-025	73%	61%	0.11	1.5	>909	>909	14
D-026	68%	54%	0.08	1.7	>1,250	>1,250	21
D-027	59%	58	0.6		>169	97	
D-028	67%	13	0.18		>556.	69	
D-029	49	3.0	0.06		882	54	
D-030	50	5	0.07		758	70	
D-031	74	10	0.12		>833	83	
D-034	19	11	0.15		131	74	
D-035	9	3	0.05		199	65	

Table 2


Compound	Alpha IC <sub>50</sub>	Beta IC <sub>50</sub>	Delta IC <sub>50</sub>	Gamma IC <sub>50</sub>	Alpha/Delta Ratio	Beta/Delta Ratio	Gamma/Delta Ratio
D-036	63%	31	0.4		>226	69	
D-037	64%	80	0.8		>125	100	
D-039	77%	66%	0.9	38	>111	>111	42
D-038	77%	63%	0.6	60	>167	>170	100
D-040	77%	64%	1.7		>61	>61	
D-041	67%	65%	4		>25	>25	
D-042	70%	25	3		>32	8	
D-043	83%	77%	2.1		>47	>47	
D-044	105	61	4.2		25	15	
D-045	98%	74%	7.6		>13	>13	
D-046	64%	95	9		>11	11	
D-047	30	9	0.09	0.5	333	100	5.6
D-048	70	14	0.16		449	90	
D-049	110%	30	1.0		>100	30	
D-050	99%	41	1.6		>63	26	
D-051	89%	57%	3.3		>31	>31	
D-052	0.7	69%	8		0.09	>13	

- 227 -

Table 2

Compound	Alpha IC <sub>50</sub>	Beta IC <sub>50</sub>	Delta IC <sub>50</sub>	Gamma IC <sub>50</sub>	Alpha/Delta Ratio	Beta/Delta Ratio	Gamma/Delta Ratio
D-121	69%	70%	0.48		>211	>211	
D-999	105	71%	47	60	2.2	2.1	1.3
LY294002	1.2	0.4	0.23		5.3	1.7	

1) Compound D-121 is 3-phenyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one





- 228 -

B. Biochemical Assay using 200  $\mu$ M ATP

In Part A, above, compounds of the invention were tested to establish their  $IC_{50}$  for inhibition of the alpha, beta, delta, and gamma isoforms of PI3K using 20  $\mu$ M ATP. A further screen was performed to establish the  $IC_{50}$  for inhibition of the four PI3K isoforms at a final concentration of 200  $\mu$ M ATP, 10-fold greater, and substantially closer to the normal physiological concentration of ATP in cells. This selectivity protocol is identical to that described above, except the 3X stock ATP concentration was 600  $\mu$ M. Data from this assay are summarized in Table 3, below. The observed sensitivity to ATP concentration suggests that these PI3K $\delta$  inhibitor compounds act as ATP competitors.

Table 3

Compound	Alpha IC <sub>50</sub>	Beta IC <sub>50</sub>	Delta IC <sub>50</sub>	Gamma IC <sub>50</sub>	Alpha/Delta Ratio	Beta/Delta Ratio	Gamma/Delta Ratio
D-000	91±1%	84±2%	2±1	35±35	91	84	18
D-005	104%	82%	11	91%	20	16	17
D-006	104±1%	44±5	0.92±0.1	87±33	226	48	95
D-007	92±11%	72±12	0.73±0.2	88±4	252	99	121
D-009	70%	18	0.7	53	200	26	76
D-010	74±18%	33±4	0.23±0.2	6±3	658	144	27
D-011	88±4%	105±35	0.25±0.2	61±70	700	420	244
D-012	70±4%	108±4	1.3±0.4	50±0	107	83	38
D-013	117±8%	73±24%	0.51±0.6	12±1	461	289	24
D-014	100±6%	13±0	0.5±0.4	5±3	398	26	10
D-015	95±22%	81±3%	1.1±0.5	83±37%	180	154	160
D-019	100%	100	30	33	7	3	1
D-022	88%	101%	4.2	60%	42	48	29
D-025	89±11%	77±6%	0.32±0.3	7.8±3	556	478	24
D-026	83±1%	77±8%	0.38±0.2	13±10	443	411	34
D-027	74%	110	4	60	37	28	15
D-028	100%	81%	1.6	29	125	101	18
D-029	110±12%	34±4	0.34±0.08	13±0.7	653	101	37

Table 3

Compound	Alpha IC <sub>50</sub>	Beta IC <sub>50</sub>	Delta IC <sub>50</sub>	Gamma IC <sub>50</sub>	Alpha/Delta Ratio	Beta/Delta Ratio	Gamma/Delta Ratio
D-030	95±11%	80±14	0.53±0.05	31±10	362	152	59
D-031	87±10%	137±23	0.2±0.01	155±60	903	707	802
D-034	92±11%	103±4	1.2±0.3	34±1	153	85	28
D-035	95±6	34±6	0.49±0.1	6.8±1	193	69	14
D-036	99%	73%	4.1	72	48	36	18
D-037	112%	58%	3.5	45	64	33	13
D-038	69%	74%	1.8	55	77	82	31
D-039	85%	65%	2.6	57%	65	50	44
D-047	81%	30	0.2	4.5	810	150	23
D-048	90±57	95±7	1.4±0.9	123±40	67	70	91
D-121	71%	62%	0.9	61%	158	138	136
D-999	62%	71%	75	90	2	2	1
LY294002	23±5	3.7±2	2.1±1.5	29±13	11	2	13

- 231 -

EXAMPLE 12Cell-Based Assay Data for  
Inhibitors of PI3K $\delta$  Activity

5

Using the methods described in Examples 3-  
5, above, compounds of the invention were tested for  
inhibitory activity and potency in assays of stim-  
10 ulated B and T cell proliferation, neutrophil (PMN)  
migration, and neutrophil (PMN) elastase release.  
Data from these assays are set forth in Table 4,  
below. In Table 4, the values shown are effective  
concentrations of the compound ( $EC_{50}$ ;  $\mu M$ ). Where no  
15 value is given, no assay was performed.

- 232 -

Table 4				
Compound	Mouse BCR Stim (EC <sub>50</sub> )	Mouse TCE Stim (EC <sub>50</sub> )	Human PMN Elastase (EC <sub>50</sub> )	Human PMN Migration (EC <sub>50</sub> )
D-000	0.9±0.4	5.5±4	2.2±2	1-5
D-003	3.9	5.7		
D-005	0.7±0.1	3.9	4.3±1	
D-006	0.2±0.1	5.3	0.3±0.1	
D-007	0.3±0.1	4.2	0.4	
D-008	1.0			
D-009	0.3±0.2		10.5	
D-010	0.2±0.1		0.3±0.3	
D-011	0.3±0.1		0.9±0.7	
D-012	0.3±0.2		0.3	
D-013	1.4			
D-014	0.2±0.1	4.3		
D-015	1.2±0.2	1.8	1.3±0.4	2.0
D-019	0.9±0.01	0.9		
D-021	1.8	3.5		
D-022	1.8	2.3		
D-024			2.9	
D-025	0.3±0.1	4.4±0.6	0.3±0.2	0.3±0.3
D-026	0.3±0.1	3.5	0.2±0.2	0.3±0.3
D-027	>2		2	
D-028	0.4±0.2		1	
D-029	0.1±0.03	3.4±2	0.5±0.6	0.3
D-030	0.1±0.1	6	0.4±0.5	0.2
D-031	0.2±0.1		0.7±0.1	
D-034	0.6±0.4			
D-035	0.2±0.1	2.9±0.7	0.3±0.1	
D-036	0.9±0.04	4.1	5.5±5	0.2
D-037	1.2±0.4		1.3±0.4	2.0
D-038	1.4±0.1	2.9	5	

- 233 -

Table 4				
Compound	Mouse BCR Stim (EC <sub>50</sub> )	Mouse TCE Stim (EC <sub>50</sub> )	Human PMN Elastase (EC <sub>50</sub> )	Human PMN Migration (EC <sub>50</sub> )
D-039	0.9±0.1		5	
D-043	1.4	2.6		
D-045			9.0	
D-047	0.3±0.1		0.5±0.2	
D-048	0.4±0.2	5	0.9±0.2	
D-049	2.0	6.3	5.0	
D-121	1.4			
D-999	3.1±0.7	5.9	>20	1
LY294002	0.9±0.5			

- 234 -

EXAMPLE 13Assay of Inhibitors of PI3K $\delta$   
Activity in Cancer Cells

5

The effect of compounds of the invention on cancer cell proliferation was evaluated by testing one of the compounds against a panel of Chronic Myeloid Leukemia (CML) cell lines, including KU812, RWLeu4, K562, and MEG-01.

The inhibitory activity of the compound (D-000, dissolved in DMSO) was determined as follows. The tested compound was added in a series of concentrations (0.001  $\mu$ M to 20  $\mu$ M) to 96-well microtiter plates with cells (1000 to 5000 cells/well). Plates were incubated for five days at 37°C during which the control cultures without test compound were able to undergo at least two cell-division cycles. Cell growth was measured by incorporation of [ $^3$ H]-thymidine for eighteen hours added at days three, four, and five. Cells were transferred to a filter, washed and the radioactivity counted using a Matrix 96 beta counter (Packard). The percentage of cell growth was measured as follows:

30

- 235 -

$$\% \text{ Cell growth} = \frac{\begin{array}{l} \text{(average counts of cells} \\ \text{incubated with a given} \\ \text{inhibitor concentration)} \\ \times 100 \end{array}}{\begin{array}{l} \text{(average counts of the} \\ \text{cells without inhibitor)} \end{array}}$$

The EC<sub>50</sub> value in these experiments was determined by the concentration of the test compound that resulted in a radioactivity count 50% lower than that obtained using the control without inhibitor. The D-000 compound exhibited inhibitory activity with an EC<sub>50</sub> of approximately 2 µM for the KU812 and RWLeu4 lines. The compound was not found to exhibit an effect in the K562 and MEG-01 lines.

PI3Kδ inhibitors of the invention appear to inhibit CML cell growth and therefore could be useful in the treatment of benign or malignant tumors. PI3Kδ expression has been demonstrated so far mostly in cells of hematopoietic origin. However, it could be present in a broader variety of proliferating cells. Therefore, the compounds of the invention could be used to induce tumor regression and to prevent the formation of tumor metastasis in both leukemia and solid tumor or in proliferation of nontumoral origin. In addition, the compounds could be used both alone and in combination with other pharmacologically active compounds or in combination with radiation as a sensitizing agent.



- 236 -


EXAMPLE 14Measurement of elastase exocytosis  
in mouse air pouch lavage

5

The effect of D-030 on leukocyte influx and neutrophil elastase exocytosis in animal models was tested. The six-day air pouch model is an *in vivo* inflammation model that histologically resembles a joint synovium. A lining of organized mononuclear cells and fibroblasts develops that closely resembles a synovial cavity. The model represents an "acute" model of a chronic disease (e.g., rheumatoid arthritis). This model allows for the *in vivo* evaluation of agents to block cellular influx into the air pouch under the influence of an inflammatory stimulus.

20

25



30

The test was performed as follows: on day zero, groups of rats were shaved and 10 ml of air was injected subcutaneously on the back of each, forming a pouch. On day three, 10 ml of air was

- 237 -

reinjected. Six hours prior to TNF challenge on day six, one group of rats (n=6) received D-030 (100 mg/kg in PEG 400 vehicle) orally, and another group (n=12) received vehicle alone orally. Six hours following dosing, the air pouches of both groups received 2.5 ng of TNF. Twelve hours following dosing, the pouches were washed with saline, and the resulting lavage fluid was analyzed for leukocyte counts and neutrophil elastase activity. In addition, blood was drawn to determine the levels of D-030 in circulation. The results were as follows: rats that received D-030 for twelve hours had an average of 8.7  $\mu$ M of compound in circulation and had an 82% reduction in total leukocytes in the lavage fluid compared to vehicle controls. Reductions in specific leukocyte counts were as follows: neutrophils (90%), eosinophils (66%), and lymphocytes (70%). Quantitation of neutrophil elastase showed that D-030-treated rats had elastase levels that were somewhat reduced (15%) versus vehicle controls.

In another test, an area of the mouse back was shaved using clippers, and an air pouch was created by injecting 3 ml air subcutaneously. On day three, the air injection was repeated. On day six, the animals were dosed with either D-030 (32 mg/kg in LABRAFIL®) or LABRAFIL® only one hour before and two hours after challenge with TNF- $\alpha$  (0.5 ng in 1 ml PBS), or PBS only. PBS is phosphate buffered saline. Four hours after TNF challenge, the animals were anesthetized and the pouches were lavaged with 2 mL of 0.9% saline with 2 mM EDTA. The lavages were centrifuged at 14,000 rpm in a microcentrifuge. Fifty microliters of the super-

- 238 -

natant was used to measure elastase exocytosis according to the procedure described above.

5 As shown in Figure 9, TNF challenge induced a high level of elastase exocytosis compared to PBS challenged animals. However, when the TNF challenged animals were treated with D-030, a significant decrease in the elastase activity in the air pouch lavages was observed.


10 All publications and patent documents cited in this specification are incorporated herein by reference for all that they disclose.

15 While the present invention has been described with specific reference to certain preferred embodiments for purposes of clarity and understanding, it will be apparent to the skilled artisan that further changes and modifications can be practiced within the scope of the invention as it is defined in the claims set forth below. Accordingly, no  
20 limitations should be placed on the invention other than those specifically recited in the claims.

- 239 -

## WHAT IS CLAIMED IS:

1. A method of disrupting leukocyte function comprising contacting leukocytes with a compound having a structure



wherein A is an optionally substituted monocyclic or bicyclic ring system containing at least two nitrogen atoms, and at least one ring of the system is aromatic;

X is selected from the group consisting of  $C(R^b)_2$ ,  $CH_2CHR^b$ , and  $CH=C(R^b)$ ;

Y is selected from the group consisting of null, S, SO,  $SO_2$ , NH, O,  $C(=O)$ ,  $OC(=O)$ ,  $C(=O)O$ , and  $NHC(=O)CH_2S$ ;

$R^1$  and  $R^2$ , independently, are selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl, aryl, heteroaryl, halo,  $NHC(=O)C_{1-3}$ alkylene $N(R^a)_2$ ,  $NO_2$ ,  $OR^a$ ,  $CF_3$ ,  $OCF_3$ ,  $N(R^a)_2$ , CN,  $OC(=O)R^a$ ,  $C(=O)R^a$ ,  $C(=O)OR^a$ ,  $arylOR^b$ , Het,  $NR^aC(=O)C_{1-3}$ alkylene $C(=O)OR^a$ ,  $arylOC_{1-3}$ alkylene $N(R^a)_2$ ,  $arylOC(=O)R^a$ ,  $C_{1-4}$ alkylene $C(=O)OR^a$ ,  $OC_{1-4}$ alkylene $C(=O)OR^a$ ,  $C_{1-4}$ alkylene $OC_{1-4}$ alkylene $C(=O)OR^a$ ,  $C(=O)NR^aSO_2R^a$ ,  $C_{1-4}$ alkylene $N(R^a)_2$ ,  $C_{2-6}$ alkenylene $N(R^a)_2$ ,  $C(=O)NR^aC_{1-4}$ alkylene $OR^a$ ,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $OC_{2-4}$ alkylene $N(R^a)_2$ ,  $OC_{1-4}$ alkylene $CH(OR^b)CH_2N(R^a)_2$ ,  $OC_{1-4}$ alk-

- 240 -

yleneHet,  $\text{OC}_{2-4}\text{alkyleneOR}^a$ ,  $\text{OC}_{2-4}\text{alkyleneNR}^a\text{C}(=\text{O})\text{OR}^a$ ,  $\text{NR}^a\text{C}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{NR}^a\text{C}(=\text{O})\text{R}^a$ ,  $\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{N}(\text{SO}_2\text{C}_{1-4}\text{alkyl})_2$ ,  $\text{NR}^a(\text{SO}_2\text{C}_{1-4}\text{alkyl})$ ,  $\text{SO}_2\text{N}(\text{R}^a)_2$ ,  $\text{OSO}_2\text{CF}_3$ ,  $\text{C}_{1-3}\text{alkylenearyl}$ ,  $\text{C}_{1-4}\text{alkyleneHet}$ ,  $\text{C}_{1-6}\text{alkyleneOR}^b$ ,  $\text{C}_{1-3}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{C}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{NHC}(=\text{O})\text{C}_{1-3}\text{alkylenearyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{3-8}\text{heterocycloalkyl}$ ,  $\text{arylOC}_{1-3}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{arylOC}(=\text{O})\text{R}^b$ ,  $\text{NHC}(=\text{O})\text{C}_{1-3}\text{alkyleneC}_{3-8}\text{heterocycloalkyl}$ ,  $\text{NHC}(=\text{O})\text{C}_{1-3}\text{alkyleneHet}$ ,  $\text{OC}_{1-4}\text{alkyleneOC}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^b$ ,  $\text{C}(=\text{O})\text{C}_{1-4}\text{alkyleneHet}$ , and  $\text{NHC}(=\text{O})\text{haloC}_{1-6}\text{alkyl}$ ;

or  $\text{R}^1$  and  $\text{R}^2$  are taken together to form a 3- or 4-membered alkylene or alkenylene chain component of a 5- or 6-membered ring, optionally containing at least one heteroatom;

$\text{R}^3$  is selected from the group consisting of optionally substituted hydrogen,  $\text{C}_{1-6}\text{alkyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{3-8}\text{heterocycloalkyl}$ ,  $\text{C}_{1-4}\text{alkylenecycloalkyl}$ ,  $\text{C}_{2-6}\text{alkenyl}$ ,  $\text{C}_{1-3}\text{alkylenearyl}$ ,  $\text{arylC}_{1-3}\text{alkyl}$ ,  $\text{C}(=\text{O})\text{R}^a$ ,  $\text{aryl}$ ,  $\text{heteroaryl}$ ,  $\text{C}(=\text{O})\text{OR}^a$ ,  $\text{C}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{C}(=\text{S})\text{N}(\text{R}^a)_2$ ,  $\text{SO}_2\text{R}^a$ ,  $\text{SO}_2\text{N}(\text{R}^a)_2$ ,  $\text{S}(=\text{O})\text{R}^a$ ,  $\text{S}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{C}(=\text{O})\text{NR}^a\text{C}_{1-4}\text{alkyleneOR}^a$ ,  $\text{C}(=\text{O})\text{NR}^a\text{C}_{1-4}\text{alkyleneHet}$ ,  $\text{C}(=\text{O})\text{C}_{1-4}\text{alkylenearyl}$ ,  $\text{C}(=\text{O})\text{C}_{1-4}\text{alkyleneheteroaryl}$ ,  $\text{C}_{1-4}\text{alkylenearyl}$  substituted with one or more of  $\text{SO}_2\text{N}(\text{R}^a)_2$ ,  $\text{N}(\text{R}^a)_2$ ,  $\text{C}(=\text{O})\text{OR}^a$ ,  $\text{NR}^a\text{SO}_2\text{CF}_3$ ,  $\text{CN}$ ,  $\text{NO}_2$ ,  $\text{C}(=\text{O})\text{R}^a$ ,  $\text{OR}^a$ ,  $\text{C}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ , and  $\text{OC}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{C}_{1-4}\text{alkyleneheteroaryl}$ ,  $\text{C}_{1-4}\text{alkyleneHet}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{C}_{1-4}\text{alkylenearyl}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{C}_{1-4}\text{alkyleneheteroaryl}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{Het}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{C}_{1-4}\text{alkyleneOR}^a$ ,  $\text{C}_{1-4}\text{alkyleneNR}^a\text{C}(=\text{O})\text{R}^a$ ,  $\text{C}_{1-4}\text{alkyleneOC}_{1-4}\text{alkyleneOR}^a$ ,  $\text{C}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^a$ , and  $\text{C}_{1-4}\text{alkyleneOC}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^a$ ;

$\text{R}^a$  is selected from the group consisting of hydrogen,  $\text{C}_{1-6}\text{alkyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{3-8}\text{heterocyclo-}$

- 241 -

alkyl,  $C_{1-3}$ alkyleneN(R<sup>c</sup>)<sub>2</sub>, aryl, arylC<sub>1-3</sub>alkyl,  
C<sub>1-3</sub>alkylenearyl, heteroaryl, heteroarylC<sub>1-3</sub>alkyl, and  
C<sub>1-3</sub>alkyleneheteroaryl;

or two R<sup>a</sup> groups are taken together to form  
a 5- or 6-membered ring, optionally containing at  
least one heteroatom;

R<sup>b</sup> is selected from the group consisting of  
hydrogen, C<sub>1-6</sub>alkyl, heteroC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenehet-  
eroC<sub>1-3</sub>alkyl, arylheteroC<sub>1-3</sub>alkyl, aryl, heteroaryl,  
arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl,  
and C<sub>1-3</sub>alkyleneheteroaryl;

R<sup>c</sup> is selected from the group consisting of  
hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, aryl, and hetero-  
aryl;

Het is a 5- or 6-membered heterocyclic  
ring, saturated or partially or fully unsaturated,  
containing at least one heteroatom selected from the  
group consisting of oxygen, nitrogen, and sulfur,  
and optionally substituted with C<sub>1-4</sub>alkyl or C(=O)OR<sup>a</sup>;

and pharmaceutically acceptable salts and  
solvates,

in an amount sufficient to inhibit phos-  
phatidylinositol 3-kinase delta activity in said  
leukocytes.

- 242 -

2'. The method according to claim 1 wherein the compound is selected from the group consisting of

2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-6,7-dimethoxy-3H-quinazolin-4-one

2-(6-aminopurin-o-ylmethyl)-6-bromo-3-(2-chlorophenyl)-3H-quinazolin-4-one

2-(6-aminopurin-o-ylmethyl)-3-(2-chlorophenyl)-7-fluoro-3H-quinazolin-4-one

2-(6-aminopurin-9-ylmethyl)-6-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one

2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-5-fluoro-3H-quinazolin-4-one

2-(6-aminopurin-o-ylmethyl)-5-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one

2-(6-aminopurin-9-ylmethyl)-3-(2-chlorophenyl)-5-methyl-3H-quinazolin-4-one

2-(6-aminopurin-9-ylmethyl)-8-chloro-3-(2-chlorophenyl)-3H-quinazolin-4-one

2-(6-aminopurin-9-ylmethyl)-3-biphenyl-2-yl-5-chloro-3H-quinazolin-4-one

5-chloro-2-(9H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one

5-chloro-3-(2-fluorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one

2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-fluorophenyl)-3H-quinazolin-4-one

3-biphenyl-2-yl-5-chloro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one

5-chloro-3-(2-methoxyphenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one

3-(2-chlorophenyl)-5-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one

- 243 -

3-(2-chlorophenyl)-6,7-dimethoxy-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
6-bromo-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
3-(2-chlorophenyl)-8-trifluoromethyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-benzo[g]quinazolin-4-one  
6-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
8-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
3-(2-chlorophenyl)-7-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
3-(2-chlorophenyl)-7-nitro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
3-(2-chlorophenyl)-6-hydroxy-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
5-chloro-3-(2-chlorophenyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
3-(2-chlorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
3-(2-chlorophenyl)-6,7-difluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
3-(2-chlorophenyl)-6-fluoro-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
2-(6-aminopurin-9-ylmethyl)-3-(2-isopropylphenyl)-5-methyl-3H-quinazolin-4-one  
2-(6-aminopurin-9-ylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one  
3-(2-fluorophenyl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one



- 244 -

2-(6-aminopurin-9-ylmethyl)-5-chloro-3-o-tolyl-3H-quinazolin-4-one  
2-(6-aminopurin-9-ylmethyl)-5-chloro-3-(2-methoxyphenyl)-3H-quinazolin-4-one  
2-(2-amino-9H-purin-6-ylsulfanylmethyl)-3-cyclopropyl-5-methyl-3H-quinazolin-4-one  
3-cyclopropylmethyl-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
2-(6-aminopurin-9-ylmethyl)-3-cyclopropylmethyl-5-methyl-3H-quinazolin-4-one  
2-(2-amino-9H-purin-6-ylsulfanylmethyl)-3-cyclopropylmethyl-5-methyl-3H-quinazolin-4-one  
5-methyl-3-phenethyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
2-(2-amino-9H-purin-6-ylsulfanylmethyl)-5-methyl-3-phenethyl-3H-quinazolin-4-one  
3-cyclopentyl-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
2-(6-aminopurin-9-ylmethyl)-3-cyclopentyl-5-methyl-3H-quinazolin-4-one  
3-(2-chloropyridin-3-yl)-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
2-(6-aminopurin-9-ylmethyl)-3-(2-chloropyridin-3-yl)-5-methyl-3H-quinazolin-4-one  
3-methyl-4-[5-methyl-4-oxo-2-(9H-purin-6-ylsulfanylmethyl)-4H-quinazolin-3-yl]-benzoic acid  
3-cyclopropyl-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
2-(6-aminopurin-9-ylmethyl)-3-cyclopropyl-5-methyl-3H-quinazolin-4-one  
5-methyl-3-(4-nitrobenzyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one

- 245 -

3-cyclohexyl-5-methyl-2-(9H-purin-6-ylsulfanyl)-  
methyl)-3H-quinazolin-4-one  
2-(6-aminopurin-9-ylmethyl)-3-cyclohexyl-5-methyl-  
3H-quinazolin-4-one  
2-(2-amino-9H-purin-6-ylsulfanylmethyl)-3-cyclo-  
hexyl-5-methyl-3H-quinazolin-4-one  
5-methyl-3-(E-2-phenylcyclopropyl)-2-(9H-purin-6-  
ylsulfanylmethyl)-3H-quinazolin-4-one  
3-(2-chlorophenyl)-5-fluoro-2-[(9H-purin-6-ylamino)-  
methyl]-3H-quinazolin-4-one  
2-[(2-amino-9H-purin-6-ylamino)methyl]-3-(2-chloro-  
phenyl)-5-fluoro-3H-quinazolin-4-one  
5-methyl-2-[(9H-purin-6-ylamino)methyl]-3-o-tolyl-  
3H-quinazolin-4-one  
2-[(2-amino-9H-purin-6-ylamino)methyl]-5-methyl-3-o-  
tolyl-3H-quinazolin-4-one  
2-[(2-fluoro-9H-purin-6-ylamino)methyl]-5-methyl-3-  
o-tolyl-3H-quinazolin-4-one  
(2-chlorophenyl)-dimethylamino-(9H-purin-6-ylsulfan-  
ylmethyl)-3H-quinazolin-4-one  
5-(2-benzyloxyethoxy)-3-(2-chlorophenyl)-2-(9H-  
purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
6-aminopurine-9-carboxylic acid 3-(2-chlorophenyl)-  
5-fluoro-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl  
ester  
N-[3-(2-chlorophenyl)-5-fluoro-4-oxo-3,4-dihydro-  
quinazolin-2-ylmethyl]-2-(9H-purin-6-ylsulfanyl)-  
acetamide  
2-[1-(2-fluoro-9H-purin-6-ylamino)ethyl]-5-methyl-3-  
o-tolyl-3H-quinazolin-4-one  
5-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-3-o-tolyl-  
3H-quinazolin-4-one

- 246 -

2-(6-dimethylaminopurin-9-ylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one  
5-methyl-2-(2-methyl-6-oxo-1,6-dihydro-purin-7-ylmethyl)-3-o-tolyl-3H-quinazolin-4-one  
5-methyl-2-(2-methyl-6-oxo-1,6-dihydro-purin-9-ylmethyl)-3-o-tolyl-3H-quinazolin-4-one  
2-(amino-dimethylaminopurin-9-ylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one  
2-(2-amino-9H-purin-6-ylsulfanylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one  
2-(4-amino-1,3,5-triazin-2-ylsulfanylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one  
5-methyl-2-(7-methyl-7H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one  
5-methyl-2-(2-oxo-1,2-dihydro-pyrimidin-4-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one  
5-methyl-2-purin-7-ylmethyl-3-o-tolyl-3H-quinazolin-4-one  
5-methyl-2-purin-9-ylmethyl-3-o-tolyl-3H-quinazolin-4-one  
5-methyl-2-(9-methyl-9H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one  
2-(2,6-Diamino-pyrimidin-4-ylsulfanylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one  
5-methyl-2-(5-methyl-[1,2,4]triazolo[1,5-a]pyrimidin-7-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one  
5-methyl-2-(2-methylsulfanyl-9H-purin-6-ylsulfanylmethyl)-3-o-tolyl-3H-quinazolin-4-one  
2-(2-hydroxy-9H-purin-6-ylsulfanylmethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one

- 247 -

5-methyl-2-(1-methyl-1*H*-imidazol-2-ylsulfanyl-  
methyl)-3-*o*-tolyl-3*H*-quinazolin-4-one  
5-methyl-3-*o*-tolyl-2-(1*H*-[1,2,4]triazol-3-ylsulfan-  
ylmethyl)-3*H*-quinazolin-4-one  
2-(2-amino-6-chloro-purin-9-ylmethyl)-5-methyl-3-*o*-  
tolyl-3*H*-quinazolin-4-one  
2-(6-aminopurin-7-ylmethyl)-5-methyl-3-*o*-tolyl-3*H*-  
quinazolin-4-one  
2-(7-amino-1,2,3-triazolo[4,5-*d*]pyrimidin-3-yl-  
methyl)-5-methyl-3-*o*-tolyl-3*H*-quinazolin-4-one  
2-(7-amino-1,2,3-triazolo[4,5-*d*]pyrimidin-1-yl-  
methyl)-5-methyl-3-*o*-tolyl-3*H*-quinazolin-4-one  
2-(6-amino-9*H*-purin-2-ylsulfanylmethyl)-5-methyl-3-  
*o*-tolyl-3*H*-quinazolin-4-one  
2-(2-amino-6-ethylamino-pyrimidin-4-ylsulfanyl-  
methyl)-5-methyl-3-*o*-tolyl-3*H*-quinazolin-4-one  
2-(3-amino-5-methylsulfanyl-1,2,4-triazol-1-yl-  
methyl)-5-methyl-3-*o*-tolyl-3*H*-quinazolin-4-one  
2-(5-amino-3-methylsulfanyl-1,2,4-triazol-1-  
ylmethyl)-5-methyl-3-*o*-tolyl-3*H*-quinazolin-4-one  
5-methyl-2-(6-methylaminopurin-9-ylmethyl)-3-*o*-  
tolyl-3*H*-quinazolin-4-one  
2-(6-benzylaminopurin-9-ylmethyl)-5-methyl-3-*o*-  
tolyl-3*H*-quinazolin-4-one  
2-(2,6-diaminopurin-9-ylmethyl)-5-methyl-3-*o*-tolyl-  
3*H*-quinazolin-4-one  
5-methyl-2-(9*H*-purin-6-ylsulfanylmethyl)-3-*o*-tolyl-  
3*H*-quinazolin-4-one  
3-isobutyl-5-methyl-2-(9*H*-purin-6-ylsulfanylmethyl)-  
3*H*-quinazolin-4-one  
*N*-{2-[5-Methyl-4-oxo-2-(9*H*-purin-6-ylsulfanyl-  
methyl)-4*H*-quinazolin-3-yl]-phenyl}-acetamide

- 248 -


5-methyl-3-(E-2-methyl-cyclohexyl)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
2-[5-methyl-4-oxo-2-(9H-purin-6-ylsulfanylmethyl)-4H-quinazolin-3-yl]-benzoic acid  
3-{2-[(2-dimethylaminoethyl)methylamino]phenyl}-5-methyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
3-(2-chlorophenyl)-5-methoxy-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
3-(2-chlorophenyl)-5-(2-morpholin-4-yl-ethylamino)-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
3-benzyl-5-methoxy-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one  
2-(6-aminopurin-9-ylmethyl)-3-(2-benzyloxyphenyl)-5-methyl-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-3-(2-hydroxyphenyl)-5-methyl-3H-quinazolin-4-one;  
2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one;  
5-methyl-2-[1-(9H-purin-6-ylamino)propyl]-3-o-tolyl-3H-quinazolin-4-one;  
2-(1-(2-fluoro-9H-purin-6-ylamino)propyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one;  
2-(1-(2-amino-9H-purin-6-ylamino)propyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one;  
2-(2-benzyloxy-1-(9H-purin-6-ylamino)ethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-5-methyl-3-{2-(2-(1-methylpyrrolidin-2-yl)-ethoxy)-phenyl}-3H-quinazolin-4-one;  
2-(6-aminopurin-9-ylmethyl)-3-(2-(3-dimethylamino-propoxy)-phenyl)-5-methyl-3H-quinazolin-4-one;

- 249 -

2-(6-aminopurin-9-ylmethyl)-5-methyl-3-(2-prop-2-ynyloxyphenyl)-3H-quinazolin-4-one; and  
2-{2-[1-(6-aminopurin-9-ylmethyl)-5-methyl-4-oxo-4H-quinazolin-3-yl]-phenoxy}-acetamide.

- 250 -

3. A method of inhibiting kinase activity of a phosphatidylinositol 3-kinase delta polypeptide comprising contacting the polypeptide with a compound having a structure



wherein A is an optionally substituted monocyclic or bicyclic ring system containing at least two nitrogen atoms, and at least one ring of the system is aromatic;

X is selected from the group consisting of  $C(R^b)_2$ ,  $CH_2CHR^b$ , and  $CH=C(R^b)$ ;

Y is selected from the group consisting of null, S, SO,  $SO_2$ , NH, O,  $C(=O)$ ,  $OC(=O)$ ,  $C(=O)O$ , and  $NHC(=O)CH_2S$ ;

$R^1$  and  $R^2$ , independently, are selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl, aryl, heteroaryl, halo,  $NHC(=O)C_{1-3}$ alkylene $N(R^a)_2$ ,  $NO_2$ ,  $OR^a$ ,  $CF_3$ ,  $OCF_3$ ,  $N(R^a)_2$ , CN,  $OC(=O)R^a$ ,  $C(=O)R^a$ ,  $C(=O)OR^a$ ,  $arylOR^b$ , Het,  $NR^aC(=O)C_{1-3}$ alkylene $C(=O)OR^a$ ,  $arylOC_{1-3}$ alkylene $N(R^a)_2$ ,  $arylOC(=O)R^a$ ,  $C_{1-4}$ alkylene $C(=O)OR^a$ ,  $OC_{1-4}$ alkylene $C(=O)OR^a$ ,  $C_{1-4}$ alkylene $OC_{1-4}$ alkylene $C(=O)OR^a$ ,  $C(=O)NR^aSO_2R^a$ ,  $C_{1-4}$ alkylene $N(R^a)_2$ ,  $C_{2-6}$ alkenylene $N(R^a)_2$ ,  $C(=O)NR^aC_{1-4}$ alkylene $OR^a$ ,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $OC_{2-4}$ alkylene $N(R^a)_2$ ,  $OC_{1-4}$ alkylene $CH(OR^b)CH_2N(R^a)_2$ ,  $OC_{1-4}$ -

- 251 -

alkyleneHet,  $\text{OC}_{2-4}\text{alkyleneOR}^a$ ,  $\text{OC}_{2-4}\text{alkyleneNR}^a\text{C}(=\text{O})\text{OR}^a$ ,  $\text{NR}^a\text{C}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{NR}^a\text{C}(=\text{O})\text{R}^a$ ,  $\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{N}(\text{SO}_2\text{C}_{1-4}\text{alkyl})_2$ ,  $\text{NR}^a(\text{SO}_2\text{C}_{1-4}\text{alkyl})$ ,  $\text{SO}_2\text{N}(\text{R}^a)_2$ ,  $\text{OSO}_2\text{CF}_3$ ,  $\text{C}_{1-3}\text{alkylenearyl}$ ,  $\text{C}_{1-4}\text{alkyleneHet}$ ,  $\text{C}_{1-6}\text{alkyleneOR}^b$ ,  $\text{C}_{1-3}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{C}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{NHC}(=\text{O})\text{C}_{1-3}\text{alkylenearyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{3-8}\text{heterocycloalkyl}$ ,  $\text{arylOC}_{1-3}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{arylOC}(=\text{O})\text{R}^b$ ,  $\text{NHC}(=\text{O})\text{C}_{1-3}\text{alkyleneC}_{3-8}\text{heterocycloalkyl}$ ,  $\text{NHC}(=\text{O})\text{C}_{1-3}\text{alkyleneHet}$ ,  $\text{OC}_{1-4}\text{alkyleneOC}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^b$ ,  $\text{C}(=\text{O})\text{C}_{1-4}\text{alkyleneHet}$ , and  $\text{NHC}(=\text{O})\text{haloC}_{1-6}\text{alkyl}$ ;

or  $\text{R}^1$  and  $\text{R}^2$  are taken together to form a 3- or 4-membered alkylene or alkenylene chain component of a 5- or 6-membered ring, optionally containing at least one heteroatom;

$\text{R}^3$  is selected from the group consisting of optionally substituted hydrogen,  $\text{C}_{1-6}\text{alkyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{3-8}\text{heterocycloalkyl}$ ,  $\text{C}_{1-4}\text{alkylenecycloalkyl}$ ,  $\text{C}_{2-6}\text{alkenyl}$ ,  $\text{C}_{1-3}\text{alkylenearyl}$ ,  $\text{arylC}_{1-3}\text{alkyl}$ ,  $\text{C}(=\text{O})\text{R}^a$ ,  $\text{aryl}$ ,  $\text{heteroaryl}$ ,  $\text{C}(=\text{O})\text{OR}^a$ ,  $\text{C}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{C}(=\text{S})\text{N}(\text{R}^a)_2$ ,  $\text{SO}_2\text{R}^a$ ,  $\text{SO}_2\text{N}(\text{R}^a)_2$ ,  $\text{S}(=\text{O})\text{R}^a$ ,  $\text{S}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{C}(=\text{O})\text{NR}^a\text{C}_{1-4}\text{alkyleneOR}^a$ ,  $\text{C}(=\text{O})\text{NR}^a\text{C}_{1-4}\text{alkyleneHet}$ ,  $\text{C}(=\text{O})\text{C}_{1-4}\text{alkylenearyl}$ ,  $\text{C}(=\text{O})\text{C}_{1-4}\text{alkyleneheteroaryl}$ ,  $\text{C}_{1-4}\text{alkylenearyl}$  optionally substituted with one or more of halo,  $\text{SO}_2\text{N}(\text{R}^a)_2$ ,  $\text{N}(\text{R}^a)_2$ ,  $\text{C}(=\text{O})\text{OR}^a$ ,  $\text{NR}^a\text{SO}_2\text{CF}_3$ ,  $\text{CN}$ ,  $\text{NO}_2$ ,  $\text{C}(=\text{O})\text{R}^a$ ,  $\text{OR}^a$ ,  $\text{C}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ , and  $\text{OC}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{C}_{1-4}\text{alkyleneheteroaryl}$ ,  $\text{C}_{1-4}\text{alkyleneHet}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{C}_{1-4}\text{alkylenearyl}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{C}_{1-4}\text{alkyleneheteroaryl}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{Het}$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{N}(\text{R}^a)_2$ ,  $\text{C}_{1-4}\text{alkyleneOR}^a$ ,  $\text{C}_{1-4}\text{alkyleneNR}^a\text{C}(=\text{O})\text{R}^a$ ,  $\text{C}_{1-4}\text{alkyleneOC}_{1-4}\text{alkyleneOR}^a$ ,  $\text{C}_{1-4}\text{alkyleneN}(\text{R}^a)_2$ ,  $\text{C}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^a$ , and  $\text{C}_{1-4}\text{alkyleneOC}_{1-4}\text{alkyleneC}(=\text{O})\text{OR}^a$ ;

$\text{R}^a$  is selected from the group consisting of hydrogen,  $\text{C}_{1-6}\text{alkyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{3-8}\text{heterocyclo-}$



- 252 -

alkyl,  $C_{1-3}$ alkylene $N(R^c)_2$ , aryl, aryl $C_{1-3}$ alkyl,  $C_{1-3}$ -alkylenearyl, heteroaryl, heteroaryl $C_{1-3}$ alkyl, and  $C_{1-3}$ alkyleneheteroaryl;

or two  $R^a$  groups are taken together to form a 5- or 6-membered ring, optionally containing at least one heteroatom;

$R^b$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl, hetero $C_{1-3}$ alkyl,  $C_{1-3}$ alkylenehetero $C_{1-3}$ alkyl, arylhetero $C_{1-3}$ alkyl, aryl, heteroaryl, aryl $C_{1-3}$ alkyl, heteroaryl $C_{1-3}$ alkyl,  $C_{1-3}$ alkylenearyl, and  $C_{1-3}$ alkyleneheteroaryl;


$R^c$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl, aryl, and heteroaryl;

Het is a 5- or 6-membered heterocyclic ring, saturated or partially or fully unsaturated, containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted with  $C_{1-4}$ alkyl or  $C(=O)OR^a$ ;

and pharmaceutically acceptable salts and solvates thereof.

- 253 -

4. A compound having a general structural formula



wherein A is an optionally substituted monocyclic or bicyclic ring system containing at least two nitrogen atoms, and at least one ring of the system is aromatic;

X is selected from the group consisting of  $C(R^b)_2$ ,  $CH_2CHR^b$ , and  $CH=C(R^b)$ ;

Y is selected from the group consisting of null, S, SO,  $SO_2$ , NH, O,  $C(=O)$ ,  $OC(=O)$ ,  $C(=O)O$ , and  $NHC(=O)CH_2S$ ;

$R^1$  and  $R^2$ , independently, are selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl, aryl, heteroaryl, halo,  $NHC(=O)C_{1-3}$ alkylene $N(R^a)_2$ ,  $NO_2$ ,  $OR^a$ ,  $CF_3$ ,  $OCF_3$ ,  $N(R^a)_2$ , CN,  $OC(=O)R^a$ ,  $C(=O)R^a$ ,  $C(=O)OR^a$ ,  $arylOR^b$ , Het,  $NR^aC(=O)C_{1-3}$ alkylene $C(=O)OR^a$ ,  $arylOC_{1-3}$ alkylene $N(R^a)_2$ ,  $arylOC(=O)R^a$ ,  $C_{1-4}$ alkylene $C(=O)OR^a$ ,  $OC_{1-4}$ alkylene $C(=O)OR^a$ ,  $C_{1-4}$ alkylene $OC_{1-4}$ alkylene $C(=O)OR^a$ ,  $C(=O)NR^aSO_2R^a$ ,  $C_{1-4}$ alkylene $N(R^a)_2$ ,  $C_{2-6}$ alkenylene $N(R^a)_2$ ,  $C(=O)NR^aC_{1-4}$ alkylene $OR^a$ ,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $OC_{2-4}$ alkylene $N(R^a)_2$ ,  $OC_{1-4}$ alkylene $CH(OR^b)CH_2N(R^a)_2$ ,  $OC_{1-4}$ alkyleneHet,  $OC_{2-4}$ alkylene $OR^a$ ,  $OC_{2-4}$ alkylene $NR^aC(=O)OR^a$ ,  $NR^aC_{1-4}$ alkylene $N(R^a)_2$ ,  $NR^aC(=O)R^a$ ,  $NR^aC(=O)N(R^a)_2$ ,  $N(SO_2C_{1-4}alkyl)_2$ ,  $NR^a(SO_2C_{1-4}alkyl)$ ,  $SO_2N(R^a)_2$ ,  $OSO_2CF_3$ ,  $C_{1-3}$ alkylenearyl,  $C_{1-4}$ alkyleneHet,  $C_{1-6}$ alkylene $OR^b$ ,

- 254 -

$C_{1-3}$ alkylene $N(R^a)_2$ ,  $C(=O)N(R^a)_2$ ,  $NHC(=O)C_1-$   
 $C_3$ alkylenearyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocycloalkyl,  
 aryl $OC_{1-3}$ alkylene $N(R^a)_2$ , aryl $OC(=O)R^b$ ,  $NHC(=O)-$   
 $C_{1-3}$ alkylene $C_{3-8}$ heterocycloalkyl,  $NHC(=O)C_{1-3}$ alkylene-  
 Het,  $OC_{1-4}$ alkylene $OC_{1-4}$ alkylene $C(=O)OR^b$ ,  
 $C(=O)C_{1-4}$ alkyleneHet, and  $NHC(=O)haloC_{1-6}$ alkyl;

or  $R^1$  and  $R^2$  are taken together to form a  
 3- or 4-membered alkylene or alkenylene chain  
 component of a 5- or 6-membered ring, optionally  
 containing at least one heteroatom;

$R^3$  is selected from the group consisting of  
 optionally substituted hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cyclo-  
 alkyl,  $C_{3-8}$ heterocycloalkyl,  $C_{1-4}$ alkylenecycloalkyl,  
 $C_{2-6}$ alkenyl,  $C_{1-3}$ alkylenearyl, aryl $C_{1-3}$ alkyl,  $C(=O)R^a$ ,  
 aryl, heteroaryl,  $C(=O)OR^a$ ,  $C(=O)N(R^a)_2$ ,  $C(=S)N(R^a)_2$ ,  
 $SO_2R^a$ ,  $SO_2N(R^a)_2$ ,  $S(=O)R^a$ ,  $S(=O)N(R^a)_2$ ,  $C(=O)NR^aC_{1-4}-$   
 alkylene $OR^a$ ,  $C(=O)NR^aC_{1-4}$ alkyleneHet,  $C(=O)C_{1-4}$ alkyl-  
 enearyl,  $C(=O)C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkylenearyl  
 optionally substituted with one or more of halo,  
 $SO_2N(R^a)_2$ ,  $N(R^a)_2$ ,  $C(=O)OR^a$ ,  $NR^aSO_2CF_3$ , CN,  $NO_2$ ,  $C(=O)R^a$ ,  
 $OR^a$ ,  $C_{1-4}$ alkylene $N(R^a)_2$ , and  $OC_{1-4}$ alkylene $N(R^a)_2$ ,  
 $C_{1-4}$ alkyleneheteroaryl,  $C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkylene-  
 $C(=O)C_{1-4}$ alkylenearyl,  $C_{1-4}$ alkylene $C(=O)C_{1-4}$ alkylene-  
 heteroaryl,  $C_{1-4}$ alkylene $C(=O)Het$ ,  $C_{1-4}$ alkylene $C(=O)-$   
 $N(R^a)_2$ ,  $C_{1-4}$ alkylene $OR^a$ ,  $C_{1-4}$ alkylene $NR^aC(=O)R^a$ ,  
 $C_{1-4}$ alkylene $OC_{1-4}$ alkylene $OR^a$ ,  $C_{1-4}$ alkylene $N(R^a)_2$ ,  
 $C_{1-4}$ alkylene $C(=O)OR^a$ , and  $C_{1-4}$ alkylene $OC_{1-4}$ alkylene-  
 $C(=O)OR^a$ ;

$R^a$  is selected from the group consisting of  
 hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocyclo-  
 alkyl,  $C_{1-3}$ alkylene $N(R^c)_2$ , aryl, aryl $C_{1-3}$ alkyl,  
 $C_{1-3}$ alkylenearyl, heteroaryl, heteroaryl $C_{1-3}$ alkyl, and  
 $C_{1-3}$ alkyleneheteroaryl;

- 255 -

or two  $R^a$  groups are taken together to form a 5- or 6-membered ring, optionally containing at least one heteroatom;


$R^b$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl, hetero $C_{1-3}$ alkyl,  $C_{1-3}$ alkylenehetero $C_{1-3}$ alkyl, arylhetero $C_{1-3}$ alkyl, aryl, heteroaryl, aryl $C_{1-3}$ alkyl, heteroaryl $C_{1-3}$ alkyl,  $C_{1-3}$ alkylenearyl, and  $C_{1-3}$ alkyleneheteroaryl;

$R^c$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl, aryl, and heteroaryl;





Het is a 5- or 6-membered heterocyclic ring, saturated or partially or fully unsaturated, containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted with  $C_{1-4}$ alkyl or  $C(=O)OR^a$ ;

and pharmaceutically acceptable salts and solvates thereof,


with the provisos that if X-Y is  $CH_2S$ , then  $R^3$  is different from



- 256 -



- 257 -







and if X-Y is CH<sub>2</sub>S, then R<sup>3</sup> is different from -CH<sub>2</sub>CH(OH)CH<sub>2</sub>OH substituted phenyl.

5. The compound of claim 4 wherein X is selected from the group consisting of CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, CH=CH, CH(CH<sub>3</sub>), CH(CH<sub>2</sub>CH<sub>3</sub>), CH<sub>2</sub>CH(CH<sub>3</sub>), and C(CH<sub>3</sub>)<sub>2</sub>.


6. The compound of claim 5 wherein Y is selected from the group consisting of null, S, and NH.

- 258 -


7. The compound of claim 5 wherein the A ring system is selected from the group consisting of




- 259 -



, and




8. The compound of claim 7 wherein the A ring system is substituted with one to three substituents selected from the group consisting of  $N(R^a)_2$ , halo,  $C_{1-3}$ alkyl,  $S(C_{1-3}$ alkyl),  $OR^a$ , and






- 260 -

9. The compound of claim 8 wherein the A ring system is substituted with one to three substituents selected from the group consisting of  $\text{NH}_2$ ,  $\text{NH}(\text{CH}_3)$ ,  $\text{N}(\text{CH}_3)_2$ ,  $\text{NHCH}_2\text{C}_6\text{H}_5$ ,  $\text{NH}(\text{C}_2\text{H}_5)$ ,  $\text{Cl}$ ,  $\text{F}$ ,  $\text{CH}_3$ ,  $\text{SCH}_3$ ,  $\text{OH}$ , and



10. The compound of claim 5 wherein  $\text{R}^1$  and  $\text{R}^2$ , independently, selected from the group consisting of hydrogen,  $\text{OR}^a$ , halo,  $\text{C}_{1-6}$ alkyl,  $\text{CF}_3$ ,  $\text{NO}_2$ ,  $\text{N}(\text{R}^a)_2$ ,  $\text{NR}^a\text{C}_{1-3}$ alkylene $\text{N}(\text{R}^a)_2$ , and  $\text{OC}_{1-3}$ alkylene $\text{OR}^a$ . Specific substituents include, but are not limited to,  $\text{H}$ ,  $\text{OCH}_3$ ,  $\text{Cl}$ ,  $\text{Br}$ ,  $\text{F}$ ,  $\text{CH}_3$ ,  $\text{CF}_3$ ,  $\text{NO}_2$ ,  $\text{OH}$ ,  $\text{N}(\text{CH}_3)_2$ ,







and  $\text{O}(\text{CH}_2)_2\text{OCH}_2\text{C}_6\text{H}_5$ , or  $\text{R}^1$  and  $\text{R}^2$  are taken together to form a five- or six-membered ring.

- 261 -





11. The compound of claim 5 wherein  $R^3$  is selected from the group consisting of  $C_{1-6}$ alkyl, aryl, heteroaryl,  $C_{3-8}$ cycloalkyl,  $C_{3-8}$ heterocycloalkyl,  $C(=O)OR^a$ ,  $C_{1-4}$ alkyleneHet,  $C_{1-4}$ alkylenecycloalkyl,  $C_{1-4}$ alkylenearyl,  $C_{1-4}$ alkyleneC(=O) $C_{1-4}$ alkylenearyl,  $C_{1-4}$ alkyleneC(=O) $OR^a$ ,  $C_{1-4}$ alkyleneC(=O) $N(R^a)_2$ ,  $C_{1-4}$ alkyleneC(=O)Het,  $C_{1-4}$ alkylene $N(R^a)_2$ , and  $C_{1-4}$ alkylene $NR^aC(=O)R^a$ .

- 262 -


12. The compound of claim 5 wherein  $R^3$  is selected from the group consisting of  $OR^a$ ,  $C_{1-6}$ alkyl, aryl, heteroaryl,  $NO_2$ ,  $N(R^a)_2$ ,  $NR^aC(=O)R^a$ ,  $C(=O)OC_2H_5$ ,  $CH_2CH(CH_3)_2$ ,



- 263 -




, and



- 264 -

13. The compound of claim 4 wherein R<sup>3</sup> is substituted with a substituent selected from the group consisting of halo, OR<sup>a</sup>, C<sub>1-6</sub>alkyl, aryl, heteroaryl, NO<sub>2</sub>, N(R<sup>a</sup>)<sub>2</sub>, NR<sup>a</sup>SO<sub>2</sub>CF<sub>3</sub>, NR<sup>a</sup>C(=O)R<sup>a</sup>, C(=O)OR<sup>a</sup>, SO<sub>2</sub>N(R<sup>a</sup>)<sub>2</sub>, CN, C(=O)R<sup>a</sup>, C<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, OC<sub>1-4</sub>alkyleneC=CR<sup>a</sup>, OC<sub>1-4</sub>alkyleneC(=O)N(R<sup>a</sup>)<sub>2</sub>, OC<sub>1-4</sub>alkylenearyl, OC<sub>1-4</sub>alkyleneheteroaryl, OC<sub>1-4</sub>alkyleneHet, OC<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>, and N(R<sup>a</sup>)C<sub>1-4</sub>alkyleneN(R<sup>a</sup>)<sub>2</sub>.

14. The compound of claim 4 wherein R<sup>3</sup> is substituted with a substituent selected from the group consisting of Cl, F, CH<sub>3</sub>, CH(CH<sub>3</sub>)<sub>2</sub>, OH, OCH<sub>3</sub>, OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, O(CH<sub>2</sub>)<sub>3</sub>N(CH<sub>3</sub>)<sub>2</sub>, OCH<sub>2</sub>C≡CH, OCH<sub>2</sub>C(=O)NH<sub>2</sub>, C<sub>6</sub>H<sub>5</sub>, NO<sub>2</sub>, NH<sub>2</sub>, NHC(=O)CH<sub>3</sub>, CO<sub>2</sub>H, and N(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, and



- 265 -

15. The compound of claim 4 selected from the group consisting of:

2-(6-aminopurin-9-ylmethyl)-3-(2-benzyloxyphenyl)-5-methyl-3H-quinazolin-4-one;

2-(6-aminopurin-9-ylmethyl)-3-(2-hydroxyphenyl)-5-methyl-3H-quinazolin-4-one;

2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one;

5-methyl-2-[1-(9H-purin-6-ylamino)propyl]-3-o-tolyl-3H-quinazolin-4-one;

2-(1-(2-fluoro-9H-purin-6-ylamino)propyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one;

2-(1-(2-amino-9H-purin-6-ylamino)propyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one;

2-(2-benzyloxy-1-(9H-purin-6-ylamino)ethyl)-5-methyl-3-o-tolyl-3H-quinazolin-4-one;

2-(6-aminopurin-9-ylmethyl)-5-methyl-3-{2-(2-(1-methylpyrrolidin-2-yl)-ethoxy)-phenyl}-3H-quinazolin-4-one;

2-(6-aminopurin-9-ylmethyl)-3-(2-(3-dimethylamino-propoxy)-phenyl)-5-methyl-3H-quinazolin-4-one;

2-(6-aminopurin-9-ylmethyl)-5-methyl-3-(2-prop-2-ynyloxyphenyl)-3H-quinazolin-4-one; and

2-{2-(1-(6-aminopurin-9-ylmethyl)-5-methyl-4-oxo-4H-quinazolin-3-yl)-phenoxy}-acetamide.

FIGURE 1




FIGURE 2






FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7





FIGURE 8






FIGURE 9

## SEQUENCE LISTING

<110> SADHU, Chanchal et al.

<120> INHIBITORS OF HUMAN PHOSPHATIDYLINOSITOL 3-KINASE DELTA

<130> 27866/36170C

<140> To be determined

<141> 2001-10-19

<150> 09/841,341

<151> 2001-04-24

<150> 60/199,655

<151> 2000-04-25

<150> 60/238,057

<151> 2000-10-25

<160> 6

<170> PatentIn version 3.0

<210> 1

<211> 5220

<212> DNA

<213> Human p110delta complete cDNA

<220>

<221> CDS

<222> (196)..(3327)



```

<400> 1
cagtcgctcc gagcgccgc gagcagagcc gccagccct gtcagctgcg ccgggacgat      60
aaggagtcag gccagggcgg gatgacactc attgattcta aagcatcttt aatctgccag      120
gcggaggggg ctttctgggt ctttcttggga ctattccaga gaggacaact gtcattctggg      180
aagtaacaac gcagg atg ccc cct ggg gtg gac tgc ccc atg gaa ttc tgg      231
          Met Pro Pro Gly Val Asp Cys Pro Met Glu Phe Trp
          1          5          10
acc aag gag gag aat cag agc gtt gtg gtt gac ttc ctg ctg ccc aca      279
Thr Lys Glu Glu Asn Gln Ser Val Val Val Asp Phe Leu Leu Pro Thr
          15          20          25
ggg gtc tac ctg aac ttc cct gtg tcc cgc aat gcc aac ctc agc acc      327
Gly Val Tyr Leu Asn Phe Pro Val Ser Arg Asn Ala Asn Leu Ser Thr
          30          35          40
atc aag cag ctg ctg tgg cac cgc gcc cag tat gag ccg ctc ttc cac      375
Ile Lys Gln Leu Leu Trp His Arg Ala Gln Tyr Glu Pro Leu Phe His
          45          50          55          60
atg ctc agt ggc ccc gag gcc tat gtg ttc acc tgc atc aac cag aca      423
Met Leu Ser Gly Pro Glu Ala Tyr Val Phe Thr Cys Ile Asn Gln Thr
          65          70          75
gcg gag cag caa gag ctg gag gac gag caa cgg cgt ctg tgt gac gtg      471
Ala Glu Gln Gln Glu Leu Glu Asp Glu Gln Arg Arg Leu Cys Asp Val
          80          85          90
cag ccc ttc ctg ccc gtc ctg cgc ctg gtg gcc cgt gag ggc gac cgc      519
Gln Pro Phe Leu Pro Val Leu Arg Leu Val Ala Arg Glu Gly Asp Arg
          95          100          105
gtg aag aag ctc atc aac tca cag atc agc ctc ctc atc ggc aaa ggc      567
Val Lys Lys Leu Ile Asn Ser Gln Ile Ser Leu Leu Ile Gly Lys Gly
          110          115          120
ctc cac gag ttt gac tcc ttg tgc gac cca gaa gtg aac gac ttt cgc      615
Leu His Glu Phe Asp Ser Leu Cys Asp Pro Glu Val Asn Asp Phe Arg
          125          130          135          140
gcc aag atg tgc caa ttc tgc gag gag gcg gcc gcc cgc cgg cag cag      663
Ala Lys Met Cys Gln Phe Cys Glu Glu Ala Ala Arg Arg Gln Gln
          145          150          155
ctg ggc tgg gag gcc tgg ctg cag tac agt ttc ccc ctg cag ctg gag      711
Leu Gly Trp Glu Ala Trp Leu Gln Tyr Ser Phe Pro Leu Gln Leu Glu
          160          165          170
ccc tcg gct caa acc tgg ggg cct ggt acc ctg cgg ctc ccg aac cgg      759
Pro Ser Ala Gln Thr Trp Gly Pro Gly Thr Leu Arg Leu Pro Asn Arg
          175          180          185
gcc ctt ctg gtc aac gtt aag ttt gag ggc agc gag gag agc ttc acc      807
Ala Leu Leu Val Asn Val Lys Phe Glu Gly Ser Glu Glu Ser Phe Thr
          190          195          200
ttc cag gtg tcc acc aag gac gtg ccg ctg gcg ctg atg gcc tgt gcc      855
Phe Gln Val Ser Thr Lys Asp Val Pro Leu Ala Leu Met Ala Cys Ala
          205          210          215          220

```

ctg cgg aag aag gcc aca gtg ttc cgg cag ccg ctg gtg gag cag ccg Leu Arg Lys Lys Ala Thr Val Phe Arg Gln Pro Leu Val Glu Gln Pro	903
225 230 235	
gaa gac tac acg ctg cag gtg aac ggc agg cat gag tac ctg tat ggc Glu Asp Tyr Thr Leu Gln Val Asn Gly Arg His Glu Tyr Leu Tyr Gly	951
240 245 250	
aac tac ccg ctc tgc cag ttc cag tac atc tgc agc tgc ctg cac agt Asn Tyr Pro Leu Cys Gln Phe Tyr Ile Cys Ser Cys Leu His Ser	999
255 260 265	
ggg ttg acc cct cac ctg acc atg gtc cat tcc tcc tcc atc ctc gcc Gly Leu Thr Pro His Leu Thr Met Val His Ser Ser Ser Ile Leu Ala	1047
270 275 280	
atg cgg gat gag cag agc aac cct gcc ccc cag gtc cag aaa ccg cgt Met Arg Asp Glu Gln Ser Asn Pro Ala Pro Gln Val Gln Lys Pro Arg	1095
285 290 295 300	
gcc aaa cca cct ccc att cct gcg aag aag cct tcc tct gtg tcc ctg Ala Lys Pro Pro Pro Ile Pro Ala Lys Lys Pro Ser Ser Val Ser Leu	1143
305 310 315	
tgg tcc ctg gag cag ccg ttc cgc atc gag ctc atc cag ggc agc aaa Trp Ser Leu Glu Gln Pro Phe Arg Ile Glu Leu Ile Gln Gly Ser Lys	1191
320 325 330	
gtg aac gcc gac gag cgg atg aag ctg gtg gtg cag gcc ggg ctt ttc Val Asn Ala Asp Glu Arg Met Lys Leu Val Val Gln Ala Gly Leu Phe	1239
335 340 345	
cac ggc aac gag atg ctg tgc aag acg gtg tcc agc tgc gag gtg agc His Gly Asn Glu Met Leu Cys Lys Thr Val Ser Ser Ser Glu Val Ser	1287
350 355 360	
gtg tgc tgc gag ccc gtg tgg aag cag cgg ctg gag ttc gac atc aac Val Cys Ser Glu Pro Val Trp Lys Gln Arg Leu Glu Phe Asp Ile Asn	1335
365 370 375 380	
atc tgc gac ctg ccc cgc atg gcc cgt ctc tgc ttt gcg ctg tac gcc Ile Cys Asp Leu Pro Arg Met Ala Arg Leu Cys Phe Ala Leu Tyr Ala	1383
385 390 395	
gtg atc gag aaa gcc aag aag gct cgc tcc acc aag aag aag tcc aag Val Ile Glu Lys Ala Lys Lys Ala Arg Ser Thr Lys Lys Lys Ser Lys	1431
400 405 410	
aag gcg gac tgc ccc att gcc tgg gcc aac ctc atg ctg ttt gac tac Lys Ala Asp Cys Pro Ile Ala Trp Ala Asn Leu Met Leu Phe Asp Tyr	1479
415 420 425	
aag gac cag ctt aag acc ggg gaa cgc tgc ctc tac atg tgg ccc tcc Lys Asp Gln Leu Lys Thr Gly Glu Arg Cys Leu Tyr Met Trp Pro Ser	1527
430 435 440	
gtc cca gat gag aag ggc gag ctg ctg aac ccc acg ggc act gtg cgc Val Pro Asp Glu Lys Gly Glu Leu Leu Asn Pro Thr Gly Thr Val Arg	1575
445 450 455 460	
agt aac ccc aac acg gat agc gcc gct gcc ctg ctc atc tgc ctg ccc Ser Asn Pro Asn Thr Asp Ser Ala Ala Ala Leu Leu Ile Cys Leu Pro	1623
465 470 475	

gag gtg gcc ccg cac ccc gtg tac tac ccc gcc ctg gag aag atc ttg Glu Val Ala Pro His Pro Val Tyr Tyr Pro Ala Leu Glu Lys Ile Leu 480 485 490	1671
gag ctg ggg cga cac agc gag tgt gtg cat gtc acc gag gag gag cag Glu Leu Gly Arg His Ser Glu Cys Val His Val Thr Glu Glu Glu Gln 495 500 505	1719
ctg cag ctg cgg gaa atc ctg gag cgg cgg ggg tct ggg gag ctg tat Leu Gln Leu Arg Glu Ile Leu Glu Arg Arg Gly Ser Gly Glu Leu Tyr 510 515 520	1767
gag cac gag aag gac ctg gtg tgg aag ctg cgg cat gaa gtc cag gag Glu His Glu Lys Asp Leu Val Trp Lys Leu Arg His Glu Val Gln Glu 525 530 535 540	1815
cac ttc ccg gag gcg cta gcc cgg ctg ctg ctg gtc acc aag tgg aac His Phe Pro Glu Ala Leu Ala Arg Leu Leu Leu Val Thr Lys Trp Asn 545 550 555	1863
aag cat gag gat gtg gcc cag atg ctc tac ctg ctg tgc tcc tgg ccg Lys His Glu Asp Val Ala Gln Met Leu Tyr Leu Leu Cys Ser Trp Pro 560 565 570	1911
gag ctg ccc gtc ctg agc gcc ctg gag ctg cta gac ttc agc ttc ccc Glu Leu Pro Val Leu Ser Ala Leu Glu Leu Leu Asp Phe Ser Phe Pro 575 580 585	1959
gat tgc cac gta ggc tcc ttc gcc atc aag tcg ctg cgg aaa ctg acg Asp Cys His Val Gly Ser Phe Ala Ile Lys Ser Leu Arg Lys Leu Thr 590 595 600	2007
gac gat gag ctg ttc cag tac ctg ctg cag ctg gtg cag gtg ctc aag Asp Asp Glu Leu Phe Gln Tyr Leu Leu Gln Val Gln Val Leu Lys 605 610 615 620	2055
tac gag tcc tac ctg gac tgc gag ctg acc aaa ttc ctg ctg gac cgg Tyr Glu Ser Tyr Leu Asp Cys Glu Leu Thr Lys Phe Leu Leu Asp Arg 625 630 635	2103
gcc ctg gcc aac cgc aag atc ggc cac ttc ctt ttc tgg cac ctc cgc Ala Leu Ala Asn Arg Lys Ile Gly His Phe Leu Phe Trp His Leu Arg 640 645 650	2151
tcc gag atg cac gtg ccg tcg gtg gcc ctg cgc ttc ggc ctc atc ctg Ser Glu Met His Val Pro Ser Val Ala Leu Arg Phe Gly Leu Ile Leu 655 660 665	2199
gag gcc tac tgc agg ggc agc acc cac cac atg aag gtg ctg atg aag Glu Ala Tyr Cys Arg Gly Ser Thr His His Met Lys Val Leu Met Lys 670 675 680	2247
cag ggg gaa gca ctg agc aaa ctg aag gcc ctg aat gac ttc gtc aag Gln Gly Glu Ala Leu Ser Lys Leu Lys Ala Leu Asn Asp Phe Val Lys 685 690 695 700	2295
ctg agc tct cag aag acc ccc aag ccc cag acc aag gag ctg atg cac Leu Ser Ser Gln Lys Thr Pro Lys Pro Gln Thr Lys Glu Leu Met His 705 710 715	2343
ttg tgc atg cgg cag gag gcc tac cta gag gcc ctc tcc cac ctg cag	2391

Leu Cys Met Arg Gln Glu Ala Tyr Leu Glu Ala Leu Ser His Leu Gln	
720 725 730	
tcc cca ctc gac ccc agc acc ctg ctg gct gaa gtc tgc gtg gag cag	2439
Ser Pro Leu Asp Pro Ser Thr Leu Leu Ala Glu Val Cys Val Glu Gln	
735 740 745	
tgc acc ttc atg gac tcc aag atg aag ccc ctg tgg atc atg tac agc	2487
Cys Thr Phe Met Asp Ser Lys Met Lys Pro Leu Trp Ile Met Tyr Ser	
750 755 760	
aac gag gag gca ggc agc ggc ggc agc gtg ggc atc atc ttt aag aac	2535
Asn Glu Glu Ala Gly Ser Gly Gly Ser Val Gly Ile Ile Phe Lys Asn	
765 770 775 780	
ggg gat gac ctc cgg cag gac atg ctg acc ctg cag atg atc cag ctc	2583
Gly Asp Asp Leu Arg Gln Asp Met Leu Thr Leu Gln Met Ile Gln Leu	
785 790 795	
atg gac gtc ctg tgg aag cag gag ggg ctg gac ctg agg atg acc ccc	2631
Met Asp Val Leu Trp Lys Gln Glu Gly Leu Asp Leu Arg Met Thr Pro	
800 805 810	
tat ggc tgc ctc ccc acc ggg gac cgc aca ggc ctc att gag gtg gta	2679
Tyr Gly Cys Leu Pro Thr Gly Asp Arg Thr Gly Leu Ile Glu Val Val	
815 820 825	
ctc cgt tca gac acc atc gcc aac atc caa ctc aac aag agc aac atg	2727
Leu Arg Ser Asp Thr Ile Ala Asn Ile Gln Leu Asn Lys Ser Asn Met	
830 835 840	
gca gcc aca gcc gcc ttc aac aag gat gcc ctg ctc aac tgg ctg aag	2775
Ala Ala Thr Ala Ala Phe Asn Lys Asp Ala Leu Leu Asn Trp Leu Lys	
845 850 855 860	
tcc aag aac ccg ggg gag gcc ctg gat cga gcc att gag gag ttc acc	2823
Ser Lys Asn Pro Gly Glu Ala Leu Asp Arg Ala Ile Glu Glu Phe Thr	
865 870 875	
ctc tcc tgt gct ggc tat tgt gtg gcc aca tat gtg ctg ggc att ggc	2871
Leu Ser Cys Ala Gly Tyr Cys Val Ala Thr Tyr Val Leu Gly Ile Gly	
880 885 890	
gat cgg cac agc gac aac atc atg atc cga gag agt ggg cag ctg ttc	2919
Asp Arg His Ser Asp Asn Ile Met Ile Arg Glu Ser Gly Gln Leu Phe	
895 900 905	
cac att gat ttt ggc cac ttt ctg ggg aat ttc aag acc aag ttt gga	2967
His Ile Asp Phe Gly His Phe Leu Gly Asn Phe Lys Thr Lys Phe Gly	
910 915 920	
atc aac cgc gag cgt gtc cca ttc atc ctc acc tat gac ttt gtc cat	3015
Ile Asn Arg Glu Arg Val Pro Phe Ile Leu Thr Tyr Asp Phe Val His	
925 930 935 940	
gtg att cag cag ggg aag act aat aat agt gag aaa ttt gaa cgg ttc	3063
Val Ile Gln Gln Gly Lys Thr Asn Asn Ser Glu Lys Phe Glu Arg Phe	
945 950 955	
cgg ggc tac tgt gaa agg gcc tac acc atc ctg cgg cgc cac ggg ctt	3111
Arg Gly Tyr Cys Glu Arg Ala Tyr Thr Ile Leu Arg Arg His Gly Leu	
960 965 970	

```

ctc ttc ctc cac ctc ttt gcc ctg atg cgg gcg gca ggc ctg cct gag      3159
Leu Phe Leu His Leu Phe Ala Leu Met Arg Ala Ala Gly Leu Pro Glu
      975                      980                      985

ctc agc tgc tcc aaa gac atc cag tat ctc aag gac tcc ctg gca ctg      3207
Leu Ser Cys Ser Lys Asp Ile Gln Tyr Leu Lys Asp Ser Leu Ala Leu
      990                      995                      1000

ggg aaa aca gag gag gag gca ctg aag cac ttc cga gtg aag ttt      3252
Gly Lys Thr Glu Glu Glu Ala Leu Lys His Phe Arg Val Lys Phe
      1005                      1010                      1015

aac gaa gcc ctc cgt gag agc tgg aaa acc aaa gtg aac tgg ctg      3297
Asn Glu Ala Leu Arg Glu Ser Trp Lys Thr Lys Val Asn Trp Leu
      1020                      1025                      1030

gcc cac aac gtg tcc aaa gac aac agg cag tagtggctcc tcccagccct      3347
Ala His Asn Val Ser Lys Asp Asn Arg Gln
      1035                      1040

gggcccaga ggaggcggct gcgggtcgtg gggaccaagc acattggtcc taaaggggct      3407

gaagagcctg aactgcacct aacgggaaag aaccgacatg gctgcctttt gtttacactg      3467

gttattttatt tatgacttga aatagtttaa ggagctaaac agccataaac ggaaacgcct      3527

ccttcattca gcggcggtgc tgggcccccc gaggctgcac ctggctctcg gctgaggatt      3587

gtcaccctcaa gtcttccagc tgggtggatct gggcccagca aagactgttc tcctcccag      3647

ggaaccttct tcccaggcct cccgccagac tgectgggtc ctggcgccctg gcggtcacct      3707

ggtgcctact gtccgacagg atgcctcgat cctcgtgcga cccaccctgt gtatcctccc      3767

tagactgagt tctggcagct ccccgaggca gccgggggtac cctctagatt cagggatgct      3827

tgctctccac ttttcaagtg ggtcttgggt acgagaattc cctcatcttt ctctactgta      3887

aagtgatttt gtttgcaggt aagaaaataa tagatgactc accacacctc tacggctggg      3947

gagatcaggc ccagcccat aaaggagaat ctacgctgggt cctcaggacg tgtaaagag      4007

atctgggcct catgtagctc accccgggtc cgcataagg caaaagcagg tcagaagcga      4067

atactctgcc attatctcaa aaatcttttt tttttttttt ttgagatggg gtcttctctc      4127

gttgcccagg ctggagtgcg gtggtgcaat cttgggtcac tgtaacctcc gcctcccagg      4187

ttcaagtgat tcttcttgcc tcagcctcct gagtagctgg gattacaggt gtgcaccacc      4247

cgtaccagc taatttttgt attttagtag agacgggggt ttcacatgt tggctgggct      4307

ggtctcgaac tcctgacctc aggtgatcca cccgcctgag cctcccaaag tgctgggatt      4367

acaggcatga gccaccagc cgggccact ctgccattgt ctaagccacc tctgaaagca      4427

ggttttaaca aaaggatgag gccagaactc ttccagaacc atcacctttg ggaacctgct      4487

gtgagagtgc tgaggtagca gaagtgtgag aacgaggggg cgtgctggga tctttctctc      4547

tgactatact tagtttgaaa tgggtcaggc ttagtcttaa gcctccaaag gcctggattt      4607

gagcagcttt agaaatgcag gttctagggc ttctcccagc cttcagaagc caactaactc      4667

```

```

tgcagatggg gctaggactg tgggctttta gcagcccaca ggtgaccta acatatcagg 4727
ccatggactc aggacctgcc cggatgatgct gttgatttct caaaggctctt ccaaaactca 4787
acagagccag aagtagccgc ccgctcagcg gctcagggtgc cagctctgtt ctgattcacc 4847
aggggtccgt cagtagtcat tgccaccgc ggggcacctc cctggccaca cgctgttcc 4907
cagcaagtgc tgaaactcac tagaccgtct gcctgtttcg aaatggggaa agccgtgcgt 4967
gcgcgttatt tatttaagtg cgctgtgtg cgcggtgtg ggagcacact ttgcaaagcc 5027
acagcgtttc tggttttggg tgtacagtct tgtgtgcctg gcgagaagaa tattttctat 5087
ttttttaagt catttcatgt ttctgtctgg ggaaggcaag ttagttaagt atcactgatg 5147
tgggttgaga ccagcactct gtgaaacctt gaaatgagaa gtaaaggcag atgaaaagaa 5207
aaaaaaaaa aaa 5220

```

<210> 2

<211> 1044

<212> PRT

<213> Human p110delta protein

<400> 2

```

Met Pro Pro Gly Val Asp Cys Pro Met Glu Phe Trp Thr Lys Glu Glu
1          5          10          15

```

```

Asn Gln Ser Val Val Val Asp Phe Leu Leu Pro Thr Gly Val Tyr Leu
          20          25          30

```

```

Asn Phe Pro Val Ser Arg Asn Ala Asn Leu Ser Thr Ile Lys Gln Leu
          35          40          45

```

```

Leu Trp His Arg Ala Gln Tyr Glu Pro Leu Phe His Met Leu Ser Gly
          50          55          60

```

```

Pro Glu Ala Tyr Val Phe Thr Cys Ile Asn Gln Thr Ala Glu Gln Gln
65          70          75          80

```

```

Glu Leu Glu Asp Glu Gln Arg Arg Leu Cys Asp Val Gln Pro Phe Leu
          85          90          95

```

```

Pro Val Leu Arg Leu Val Ala Arg Glu Gly Asp Arg Val Lys Lys Leu
          100          105          110

```

```

Ile Asn Ser Gln Ile Ser Leu Leu Ile Gly Lys Gly Leu His Glu Phe
          115          120          125

```

Asp Ser Leu Cys Asp Pro Glu Val Asn Asp Phe Arg Ala Lys Met Cys  
 130 135 140

Gln Phe Cys Glu Glu Ala Ala Ala Arg Arg Gln Gln Leu Gly Trp Glu  
 145 150 155 160

Ala Trp Leu Gln Tyr Ser Phe Pro Leu Gln Leu Glu Pro Ser Ala Gln  
 165 170 175

Thr Trp Gly Pro Gly Thr Leu Arg Leu Pro Asn Arg Ala Leu Leu Val  
 180 185 190

Asn Val Lys Phe Glu Gly Ser Glu Glu Ser Phe Thr Phe Gln Val Ser  
 195 200 205

Thr Lys Asp Val Pro Leu Ala Leu Met Ala Cys Ala Leu Arg Lys Lys  
 210 215 220

Ala Thr Val Phe Arg Gln Pro Leu Val Glu Gln Pro Glu Asp Tyr Thr  
 225 230 235 240

Leu Gln Val Asn Gly Arg His Glu Tyr Leu Tyr Gly Asn Tyr Pro Leu  
 245 250 255

Cys Gln Phe Gln Tyr Ile Cys Ser Cys Leu His Ser Gly Leu Thr Pro  
 260 265 270

His Leu Thr Met Val His Ser Ser Ser Ile Leu Ala Met Arg Asp Glu  
 275 280 285

Gln Ser Asn Pro Ala Pro Gln Val Gln Lys Pro Arg Ala Lys Pro Pro  
 290 295 300

Pro Ile Pro Ala Lys Lys Pro Ser Ser Val Ser Leu Trp Ser Leu Glu  
 305 310 315 320

Gln Pro Phe Arg Ile Glu Leu Ile Gln Gly Ser Lys Val Asn Ala Asp  
 325 330 335

Glu Arg Met Lys Leu Val Val Gln Ala Gly Leu Phe His Gly Asn Glu  
 340 345 350

Met Leu Cys Lys Thr Val Ser Ser Ser Glu Val Ser Val Cys Ser Glu  
 355 360 365

Pro Val Trp Lys Gln Arg Leu Glu Phe Asp Ile Asn Ile Cys Asp Leu  
 370 375 380

Pro Arg Met Ala Arg Leu Cys Phe Ala Leu Tyr Ala Val Ile Glu Lys  
 385 390 395 400  
 Ala Lys Lys Ala Arg Ser Thr Lys Lys Lys Ser Lys Lys Ala Asp Cys  
 405 410 415  
 Pro Ile Ala Trp Ala Asn Leu Met Leu Phe Asp Tyr Lys Asp Gln Leu  
 420 425 430  
 Lys Thr Gly Glu Arg Cys Leu Tyr Met Trp Pro Ser Val Pro Asp Glu  
 435 440 445  
 Lys Gly Glu Leu Leu Asn Pro Thr Gly Thr Val Arg Ser Asn Pro Asn  
 450 455 460  
 Thr Asp Ser Ala Ala Ala Leu Leu Ile Cys Leu Pro Glu Val Ala Pro  
 465 470 475 480  
 His Pro Val Tyr Tyr Pro Ala Leu Glu Lys Ile Leu Glu Leu Gly Arg  
 485 490 495  
 His Ser Glu Cys Val His Val Thr Glu Glu Glu Gln Leu Gln Leu Arg  
 500 505 510  
 Glu Ile Leu Glu Arg Arg Gly Ser Gly Glu Leu Tyr Glu His Glu Lys  
 515 520 525  
 Asp Leu Val Trp Lys Leu Arg His Glu Val Gln Glu His Phe Pro Glu  
 530 535 540  
 Ala Leu Ala Arg Leu Leu Leu Val Thr Lys Trp Asn Lys His Glu Asp  
 545 550 555 560  
 Val Ala Gln Met Leu Tyr Leu Leu Cys Ser Trp Pro Glu Leu Pro Val  
 565 570 575  
 Leu Ser Ala Leu Glu Leu Leu Asp Phe Ser Phe Pro Asp Cys His Val  
 580 585 590  
 Gly Ser Phe Ala Ile Lys Ser Leu Arg Lys Leu Thr Asp Asp Glu Leu  
 595 600 605  
 Phe Gln Tyr Leu Leu Gln Leu Val Gln Val Leu Lys Tyr Glu Ser Tyr  
 610 615 620  
 Leu Asp Cys Glu Leu Thr Lys Phe Leu Leu Asp Arg Ala Leu Ala Asn  
 625 630 635 640



Arg Lys Ile Gly His Phe Leu Phe Trp His Leu Arg Ser Glu Met His  
 645 650 655  
 Val Pro Ser Val Ala Leu Arg Phe Gly Leu Ile Leu Glu Ala Tyr Cys  
 660 665 670  
 Arg Gly Ser Thr His His Met Lys Val Leu Met Lys Gln Gly Glu Ala  
 675 680 685  
 Leu Ser Lys Leu Lys Ala Leu Asn Asp Phe Val Lys Leu Ser Ser Gln  
 690 695 700  
 Lys Thr Pro Lys Pro Gln Thr Lys Glu Leu Met His Leu Cys Met Arg  
 705 710 715 720  
 Gln Glu Ala Tyr Leu Glu Ala Leu Ser His Leu Gln Ser Pro Leu Asp  
 725 730 735  
 Pro Ser Thr Leu Leu Ala Glu Val Cys Val Glu Gln Cys Thr Phe Met  
 740 745 750  
 Asp Ser Lys Met Lys Pro Leu Trp Ile Met Tyr Ser Asn Glu Glu Ala  
 755 760 765  
 Gly Ser Gly Gly Ser Val Gly Ile Ile Phe Lys Asn Gly Asp Asp Leu  
 770 775 780  
 Arg Gln Asp Met Leu Thr Leu Gln Met Ile Gln Leu Met Asp Val Leu  
 785 790 795 800  
 Trp Lys Gln Glu Gly Leu Asp Leu Arg Met Thr Pro Tyr Gly Cys Leu  
 805 810 815  
 Pro Thr Gly Asp Arg Thr Gly Leu Ile Glu Val Val Leu Arg Ser Asp  
 820 825 830  
 Thr Ile Ala Asn Ile Gln Leu Asn Lys Ser Asn Met Ala Ala Thr Ala  
 835 840 845  
 Ala Phe Asn Lys Asp Ala Leu Leu Asn Trp Leu Lys Ser Lys Asn Pro  
 850 855 860  
 Gly Glu Ala Leu Asp Arg Ala Ile Glu Glu Phe Thr Leu Ser Cys Ala  
 865 870 875 880  
 Gly Tyr Cys Val Ala Thr Tyr Val Leu Gly Ile Gly Asp Arg His Ser  
 885 890 895

Asp Asn Ile Met Ile Arg Glu Ser Gly Gln Leu Phe His Ile Asp Phe  
                   900                                  905                                  910

Gly His Phe Leu Gly Asn Phe Lys Thr Lys Phe Gly Ile Asn Arg Glu  
           915                                  920                                  925

Arg Val Pro Phe Ile Leu Thr Tyr Asp Phe Val His Val Ile Gln Gln  
       930                                  935                                  940

Gly Lys Thr Asn Asn Ser Glu Lys Phe Glu Arg Phe Arg Gly Tyr Cys  
   945                                  950                                  955                                  960

Glu Arg Ala Tyr Thr Ile Leu Arg Arg His Gly Leu Leu Phe Leu His  
                   965                                  970                                  975

Leu Phe Ala Leu Met Arg Ala Ala Gly Leu Pro Glu Leu Ser Cys Ser  
                   980                                  985                                  990

Lys Asp Ile Gln Tyr Leu Lys Asp Ser Leu Ala Leu Gly Lys Thr Glu  
           995                                  1000                                  1005

Glu Glu Ala Leu Lys His Phe Arg Val Lys Phe Asn Glu Ala Leu  
       1010                                  1015                                  1020

Arg Glu Ser Trp Lys Thr Lys Val Asn Trp Leu Ala His Asn Val  
       1025                                  1030                                  1035

Ser Lys Asp Asn Arg Gln  
       1040

<210> 3

<211> 69

<212> DNA

<213> 5' Primer for p110beta

<400> 3  
 gatcgaattc ggcgccacca tggactacaa ggacgacgat gacaagtgc tcagtttcat 60  
 aatgcctcc 69

<210> 4

<211> 42

<212> DNA

<213> 3' Primer for p110beta

<400> 4

gatcgcgggcc gcttaagatc tgtagtcttt ccgaactgtg tg

42

<210> 5

<211> 40

<212> DNA

<213> 5' Primer for p110gamma

<400> 5

agaatgcggc cgcatggagc tggagaacta taaacagccc

40

<210> 6

<211> 37

<212> DNA

<213> 3' Primer for p110gamma

<400> 6

cgcggatcct taggetgaat gttctctctcc ttgtttg

37

## INTERNATIONAL SEARCH REPORT

PCT/US 02/27240

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/505 A61K31/53 C07D473/34 C07D473/38 C07D473/24  
 C07D473/16 C07D473/00 C07D473/40 C07D403/12 C07D487/04

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 01 81346 A (ICOS CORP) 1 November 2001 (2001-11-01) page 9 (last paragraph) - page 13 (end); examples; claims ---	1,3,4
X	WO 01 30768 A (BERGNES GUSTAVE ;CHABALA JOHN C (US); FENG BAINIAN (US); FINER JEF) 3 May 2001 (2001-05-03) page 8, line 3 - page 10, line 6; table 3 (page 69/90) ---	1,3,4
X	EP 0 884 310 A (PFIZER PROD INC) 16 December 1998 (1998-12-16) page 4, line 1 - page 4, line 39; example 4 --- -/--	4



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

30 January 2003

Date of mailing of the international search report

24/02/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Wörth, C

## INTERNATIONAL SEARCH REPORT

PCT/US 02/27240

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 900 568 A (PFIZER PROD INC) 10 March 1999 (1999-03-10) page 3, lines 1 and 58; page 4, lines 1 and 47; compounds of formula (F) ---	4
X	JP 55 118918 A (MITSUBISHI ELECTRIC CORP) 12 September 1980 (1980-09-12) compounds QA-1 and QA-2 ---	4
X	JP 55 118917 A (MITSUBISHI ELECTRIC CORP) 12 September 1980 (1980-09-12) compounds paragraph (10) and (11) ---	4
X	JP 56 002322 A (MITSUBISHI ELECTRIC CORP) 12 January 1981 (1981-01-12) compound MQ-1 ---	4
X	US 4 225 489 A (ROLF MEINHARD ET AL) 30 September 1980 (1980-09-30) col. 1, compounds of formula (I); see inter alia examples 1-11 and 70 ---	4
X	GB 1 356 763 A (MERCK & CO INC) 12 June 1974 (1974-06-12) page 2, line 27 -page 2, line 33 ---	4
X	RATHMAN T L ET AL: "Functionalization of 2-methyl-3-(o-tolyl)-4(3H)-quinazolinone and related compounds through carbanion reactions at the 2-methyl group" JOURNAL OF ORGANIC CHEMISTRY, AMERICAN CHEMICAL SOCIETY. EASTON, US, vol. 45, 1980, pages 2169-2176, XP002102009 ISSN: 0022-3263 figure IV; example 14B ---	4
X	AGER I R ET AL: "Synthesis and central nervous system activity of quinazolones related to 2-methyl-3-(o-tolyl)-4(3H)- quinazolinone (methaqualone)" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 20, no. 3, March 1977 (1977-03), pages 379-386, XP002156696 ISSN: 0022-2623 examples 44,45; table IV ---	4
	---	

-/--

## INTERNATIONAL SEARCH REPORT

PCT/US 02/27240

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>CHEMICAL ABSTRACTS, vol. 98, no. 1, 1983  Columbus, Ohio, US;  abstract no. 4514,  ISMAIL ET AL.: "Some reactions of 2-(bromomethyl)-3-phenylquinazolin-4(3H)-one"  XP002229325  compound (II)  abstract  &amp; IND. J. CHEM., SECT. B,  vol. 21b, no. 5, 1982, pages 461-462,</p>	4
X	<p>CHEMICAL ABSTRACTS, vol. 121, no. 9, 1994  Columbus, Ohio, US;  abstract no. 108675e,  PARASHARYA ET AL.: "4(3H)-quinazolones. Part I:  2-Alkyl/arylaminoethyl-3-p-hydroxy/methoxyphenyl-4(3H)-quinazolones"  page 1065; column 1;  XP002229326  RN 156672-76-3  abstract  &amp; J. INST. CHEM.,  vol. 64, no. 5, 1992, pages 184-185,</p>	4
X	<p>CHEMICAL ABSTRACTS, vol. 124, no. 21, 1996  Columbus, Ohio, US;  abstract no. 289434f,  BARAKAT ET AL.: "Synthesis and CNS depressant activity of some new quinazoline derivatives"  page 1334; column 2;  XP002229327  compounds of formula (I)  abstract  &amp; AL-AZHAR J. PHARM. SCI.,  vol. 14, 1994, pages 239-246,</p> <p style="text-align: center;">-/--</p>	4

## INTERNATIONAL SEARCH REPORT

PCT/US 02/27240

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 129, no. 16, 1998 Columbus, Ohio, US; abstract no. 203126a, CHERN ET AL.: "Studies on quinazolines. VII. Reactions of anthranilamide with beta-diketones; new approaches toward the synthesis of tetrahydropyrido(2,1-b)quinazolin-11-one derivatives" page 676; column 2; XP002229328 compounds of formula (II) abstract & CHEM. PHARM. BULL., vol. 46, no. 6, 1998, pages 928-933, ---	4
X	CHEMICAL ABSTRACTS, vol. 131, no. 23, 1999 Columbus, Ohio, US; abstract no. 310612U, EL-FEKY ET AL.: "Novel quinazolinones from 2-cyanomethyl-3-phenyl-4(3H)-quinazoline" page 497; column 2; XP002229329 compound (III) abstract & BOLL. CHIM. PHARM., vol. 137, no. 7, 1998, pages 286-289, ---	4
X	CHEMICAL ABSTRACTS, vol. 106, no. 13, 1987 Columbus, Ohio, US; abstract no. 102196t, EL-FEKY ET AL.: "Synthesis of certain new sulfur-containing quinazolinone derivatives likely to possess CNS depressant action" page 650; column 1; XP002229330 compounds of formula (III) abstract & EGYPT. J. PHARM. SCI., vol. 24, no. 1-4, 1985, pages 39-47, ---	4
X	HASSAN M F ET AL: "REACTIVITY OF 6-ANISYL-5-CYANO-4-OXO-2-THIOXO-1,2,3,4-TE TRAHYDRO- PYRIMIDINE TOWARDS SOME ELECTROPHILES AND NUCLEOPHILES" CHINESE JOURNAL OF CHEMISTRY, XX, XX, vol. 9, no. 3, 1991, pages 262-269, XP000925588 page 264; example 6 --- -/--	4

## INTERNATIONAL SEARCH REPORT

PCT/US 02/27240

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 378 725 A (BONJOUKLIAN ROSANNE ET AL) 3 January 1995 (1995-01-03) the whole document	1,3,4
A	WO 01 53266 A (JACKSON SHAUN ;KENCHE VIJAYA (AU); THOMPSON PHIL (AU); YAIP CINDY) 26 July 2001 (2001-07-26) claims 1,5,10,12	1,3,4



# INTERNATIONAL SEARCH REPORT

PCT/US 02/27240

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 1-3 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 4 (part)  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 4 (part)

Present claim 4 relates to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the methods of claims 1-3 and to the compounds of claims 5-15.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

## INTERNATIONAL SEARCH REPORT

PCT/US 02/27240

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0181346	A	01-11-2001	AU 5566701 A	07-11-2001
			EP 1278748 A2	29-01-2003
			NO 20025104 A	10-12-2002
			WO 0181346 A2	01-11-2001
			US 2002161014 A1	31-10-2002
WO 0130768	A	03-05-2001	AU 1439801 A	08-05-2001
			BR 0015110 A	02-07-2002
			CZ 20021428 A3	13-11-2002
			EP 1226129 A1	31-07-2002
			NO 20021907 A	07-06-2002
			WO 0130768 A1	03-05-2001
			AU 5927001 A	02-01-2002
			WO 0198278 A1	27-12-2001
EP 0884310	A	16-12-1998	BR 9801808 A	21-03-2000
			CA 2240138 A1	09-12-1998
			EP 0884310 A1	16-12-1998
			JP 11012255 A	19-01-1999
EP 0900568	A	10-03-1999	AU 736254 B2	26-07-2001
			AU 8312098 A	18-03-1999
			CA 2246839 A1	05-03-1999
			EP 0900568 A2	10-03-1999
			HU 9802021 A2	28-05-1999
			JP 11158072 A	15-06-1999
			JP 2001316267 A	13-11-2001
			NZ 331741 A	25-08-2000
			US 6136812 A	24-10-2000
			ZA 9808139 A	22-03-2000
JP 55118918	A	12-09-1980	NONE	
JP 55118917	A	12-09-1980	JP 1372998 C	07-04-1987
			JP 61038732 B	30-08-1986
JP 56002322	A	12-01-1981	JP 1294330 C	26-12-1985
			JP 60017375 B	02-05-1985
US 4225489	A	30-09-1980	DE 2644265 A1	06-04-1978
			BE 859180 A1	29-03-1978
			BR 7706506 A	06-06-1978
			CH 634070 A5	14-01-1983
			DE 2660557 C2	30-05-1984
			DK 430577 A	31-03-1978
			ES 462648 A1	16-06-1978
			ES 462758 A1	16-12-1978
			FR 2366337 A1	28-04-1978
			GB 1554057 A	17-10-1979
			JP 1282258 C	27-09-1985
			JP 53044581 A	21-04-1978
			JP 60003302 B	26-01-1985
			JP 1318512 C	29-05-1986
			JP 56081366 A	03-07-1981
			JP 60038421 B	31-08-1985
GB 1356763	A	12-06-1974	DE 2219408 A1	26-10-1972
			FR 2133979 A5	01-12-1972

## INTERNATIONAL SEARCH REPORT

PCT/US 02/27240

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
6B 1356763	A		NL 7204972 A	24-10-1972
			US 3897432 A	29-07-1975
US 5378725	A	03-01-1995	AU 678831 B2	12-06-1997
			AU 6754094 A	27-01-1995
			CA 2128046 A1	20-01-1995
			CN 1111127 A	08-11-1995
			CZ 9401692 A3	15-02-1995
			EP 0635268 A1	25-01-1995
			HU 67667 A2	28-04-1995
			JP 7053370 A	28-02-1995
			NO 942641 A	20-01-1995
			PL 304317 A1	23-01-1995
			ZA 9405103 A	15-01-1996
WO 0153266	A	26-07-2001	AU 3042601 A	31-07-2001
			EP 1257537 A1	20-11-2002
			WO 0153266 A1	26-07-2001